

PENOXsulAM GROUP 2 HERBICIDE

Marble

For Selective Weed Control in Water-Seeded Rice in California.

ACTIVE INGREDIENT:

Penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4] triazolo[1,5c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide.

WT. BY %

0.24%

OTHER INGREDIENTS:

99.76%

TOTAL:

100.00%

Contains 0.24 lb. of active ingredient per 100 lbs. bag.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand this label, find someone to explain it to you in detail.)

**See label booklet for complete First Aid, Precautionary Statements,
Directions For Use, and Storage and Disposal.**

Manufactured For:

Sharda USA LLC



7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 83529-247

EPA Est. No. **DI** 05905-IA-001; **GH** 70815-GA-002; **MA** 83411-MN-001;
MC 89332-GA-001; **SC** 39578-TX-001; **TX** 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: 45 lbs. (20.41 kg)

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For non-emergency resource information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 Monday – Friday 8 am – Noon Pacific Time, (NPIC Web site: www.npic.orst.edu). For Chemical Spill, Leak, Fire or Exposure, call CHEMTREC 800-424-9300 .	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS	
Users should:	<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

Except when treating rice fields as specified in this product label, **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Non-Target Organism Advisory Statement

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Groundwater Advisory Statement

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory Statement

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of penoxsulam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Read all Directions for Use carefully before applying.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

PRODUCT INFORMATION

Marble is a post-flood, post-emergence herbicide for selective control of susceptible grass, broadleaf, and sedge weeds in California rice. Susceptible weeds at the 1- to 3-leaf stage emerged at the time of application will be controlled (refer to **Application Rates and Weeds Controlled** table). Apply **Marble** directly into flooded fields when rice is at a minimum 2 1/2-leaf stage of growth. At this stage, the second leaf is fully expanded or unfurled with the third leaf emerged but not expanded (refer to **Application Timing** section). Average water depth at application must be maintained at 4 - 5 inches, with a minimum water depth of 3 inches in the shallowest part of the field with completely submerged weeds and no exposed soil (refer to **Water Management** section).

Rice crops grown under adverse environmental conditions, including extreme cold or heat, may express temporary crop injury when **Marble** is applied including slight height reduction or root stunting. Any crop stress or environmental factors which decrease plant metabolism and growth may reduce weed control efficacy and crop tolerance. Such effects are transient and **DO NOT** affect yield. Fertilizer top dressing will speed crop recovery.

RESTRICTIONS:

- Apply **Marble** by aerial application only.
- Pre-Harvest Interval: **DO NOT** apply **Marble** within 60 days of rice harvest.
- **DO NOT** apply **Marble** following rainfall until the water depth in checks or basins has been adjusted to meet the correct water depth requirements.
- **DO NOT** apply where runoff or irrigation water may flow directly onto agricultural land other than rice fields.
- **DO NOT** make more than 1 application per year.
- **DO NOT** apply more than 18.5 lbs. (0.044 lb. a.i. penoxsulam) of **Marble** per acre per year.
- **DO NOT** rotate treated land to crops other than rice for 3 months following application.
- **DO NOT** use on wild rice.
- **DO NOT** fish or commercially grow fish, shellfish or crustaceans on treated acres during the year of treatment.
- Chemigation: **DO NOT** apply this product through any type of irrigation system.

PRECAUTIONS:

- **Marble** may not reliably control known ALS-resistant weed biotypes.
- Applications made to greater than 3 leaf weeds, dense weed populations, and/or average water depth greater than 6 inches may result in reduced weed control.
- Static water conditions for 7 days following an application of **Marble** are advised for best results. It is important not to allow the minimum water depth to drop below 3 inches; therefore, adding water may be required. Water must be trickled in or added very slowly.
- For best results, it is advised that water not be released for 14 days following an application of **Marble**. During this time water depth must be maintained or can be increased as rice grows. Early release of water may result in reduced weed control.
- Shallow water conditions can result in reduced weed control as it may promote additional weed flushes and can increase the potential for crop injury.
- Reduced weed control can be expected at field water inlets due to dilution and movement of water.
- The use of **Marble** on rice varieties with known reduced seedling vigor when established and grown under adverse weather conditions is not advised. Use care when using **Marble** on newly released rice varieties until variety tolerance can be determined.
- Application of **Marble** to fields that have been leveled within 12 months prior to application may result in serious rice injury in areas that have been cut or filled. This does not apply to normal annual land planning activities.
- Under poor field preparation practices (i.e., lack of soil aeration, unleveled ground, large clods, excessive crop residue, weeds emerged prior to flooding, etc.), reduced weed control may occur when using **Marble**.
- Cold and/or alternating weather conditions during the planting season can result in prolonged weed seed germination resulting in reduced weed control.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 10 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wing-span for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Aircraft

Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the air-flow in flight.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications

Take precautions to minimize spray drift.

WEED RESISTANCE MANAGEMENT

For resistance management, **Marble** is a Group 2 (acetolactate synthase (ALS) enzyme inhibitor) herbicide. Any weed population may contain or develop plants naturally resistant to **Marble** and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies must be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of **Marble** or other Group 2 (acetolactate synthase (ALS) enzyme inhibitor) herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Sharda USA, LLC at www.shardausa.com.

In addition to the guidance above, registrants are encouraged to incorporate the appropriate elements of Best Management Practices from HRAC and WSSA on the label.

APPLICATION INSTRUCTIONS

Best Management Practices for Aerial Application


Equipment used in the application of **Marble** must be carefully calibrated before use and checked frequently during application to be sure it is working properly and delivering a uniform distribution pattern. Avoid overlaps that will increase **Marble** dosage above specified limits.

Application Timing

Application timing is dependent upon the proper rice growth stage and optimum timing for weed species present. **Marble** must be applied to rice after it reaches a minimum of 2 1/2-leaf stage (**Figure 1**); at this stage the second rice leaf is fully expanded or unfurled with third leaf emerged. **Marble** must be applied when susceptible weeds are from the 1- to 3-leaf stage up to 60 days before harvest (refer to the **Application Rates and Weeds Controlled** table).

It is important that the stage of rice be closely monitored prior to application to determine the correct timing for an application of **Marble**. Under cool growing conditions, rice development can be delayed; therefore, close crop development monitoring is recommended prior to application.

Figure 1



Water Management

Immediately prior to application:

- Average water depth at application must be maintained at 4 - 5 inches, with a minimum water depth of 3 inches in the shallowest part of the field, with completely submerged weeds and no exposed soil.
- If rainfall occurs prior to application, **DO NOT** apply **Marble** until water depth in checks or basins can be adjusted to meet the correct water depth requirements.

First week following application:

- For best results, maintain a static flood for 7 days following application.
- During the static flood period it is important to maintain a minimum water depth of 3 inches. If necessary, water can be slowly added or trickled in to maintain water depth.
- If water is lost from a basin or check, re-flood immediately. Early release of water may result in reduced weed control.
- Failure to maintain the recommended flood depth after an application of **Marble** may result in reduced weed control or crop tolerance.

Second week following application:

- Following the 7-day static flood period it is advised that water not be released for an additional 7 days.
- During this time, water depth must be maintained or increased as rice grows.
- Shallow water conditions can result in reduced weed control as it may promote additional weed flushes.
- If water is lost from a basin or check, re-flood immediately. Early release of water may result in reduced weed control.

At the conclusion of the 2-week treatment period, it is important to manage water flood depth to prevent secondary weed flushes which can occur if water is released or lowered.

Application Rates and Weeds Controlled (For Use in the State of California)

Weeds Controlled ¹ Common Name (Scientific Name)	Application Rate and Stage of Weed Development
Barnyardgrass (<i>Echinochloa crus-galli</i>) California Arrowhead (<i>Sagittaria montevidensis</i> spp., <i>Caclycina</i>) Common Waterplantain (<i>Alisma plantago-aquatica</i>) Ducksalad (<i>Heteranthera</i> spp.) Monochoria (<i>Monochoria</i> spp.) Watergrass (early and late) (<i>Echinochloa</i> spp.)	15 lbs. of product/A (0.036 lb. a.i/A) up to 3-leaf
Weeds Suppressed Common Name (Scientific Name)	
Gregg's Arrowhead (<i>Sagittaria longiloba</i>) Redstem (<i>Ammannia</i> spp.) ³ Rice Mimic (<i>Echinochloa</i> spp.) Ricefield Bulrush (<i>Scirpus mucronatus</i>) ³ Smallflower Umbrellaplant (<i>Cyperus difformis</i>)	
¹ Marble may not reliably control known ALS-resistant weed biotypes. Weeds that emerge after the treatment period will not be controlled. ² Under conditions of late application when weeds are greater than 3-leaf stage, dense weed populations, or average water depth greater than 6 inches deep, 18.5 lbs. (0.044 lb. a.i. penoxsulam) of Marble per acre may be applied. Application of Marble under these conditions may result in reduced weed control. ³ Ricefield bulrush (<i>Scirpus mucronatus</i>) and redstem (<i>Ammannia</i> spp.) are effectively controlled if emerged at the time of treatment with Marble .	

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in the original container. **DO NOT** store in a manner where cross-contamination with other pesticides, fertilizers, food, or feed could occur. In the event of a spill during handling or storage, absorb with sand or other inert material and dispose of absorbent in accordance with the **PESTICIDE DISPOSAL** instructions listed below.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Non-refillable Plastic Containers (Capacity Equal to or Less than 50 Pounds): **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two or more times. Offer for recycling, or puncture and dispose of in a sanitary landfill, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Non-refillable Plastic Containers (Capacity Equal to or more than 50 Pounds): **DO NOT** reuse or refill this container. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by State and local ordinances.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

NOTES

PENOXSULAM GROUP 2 HERBICIDE

Marble

For Selective Weed Control in Water-Seeded Rice in California.

ACTIVE INGREDIENT:	WT. BY %
Penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4] triazolo [1,5c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide	0.24%
OTHER INGREDIENTS:	99.76%
TOTAL:	100.00%

Contains 0.24 lb. of active ingredient per 100 lbs. bag.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

See label booklet for additional
Precautionary Statements and Directions For Use.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For non-emergency resource information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 Monday – Friday 8 am – Noon Pacific Time, (NPIC Web site: www.npic.orst.edu). For Chemical Spill, Leak, Fire or Exposure, call CHEMTREC 800-424-9300 .	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

Except when treating rice fields as specified in this product label, **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Non-Target Organism Advisory Statement

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Groundwater Advisory Statement

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory Statement

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of penoxsulam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Read all Directions for Use carefully before applying.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in the original container. **DO NOT** store in a manner where cross-contamination with other pesticides, fertilizers, food, or feed could occur. In the event of a spill during handling or storage, absorb with sand or other inert material and dispose of absorbent in accordance with the **PESTICIDE DISPOSAL** instructions listed below.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Non-refillable Plastic Containers (Capacity Equal to or Less than 50 Pounds): **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two or more times. Offer for recycling, or puncture and dispose of in a sanitary landfill, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Manufactured For: Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 83529-247 EPA Est. No. **DI** 05905-IA-001; **GH** 70815-GA-002; **MA** 83411-MN-001;
MC 89332-GA-001; **SC** 39578-TX-001; **TX** 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: 45 lbs. (20.41 kg)

OPEN HERE