

AMINOPYRALID	GROUP	4	HERBICIDE
2,4-D	GROUP	4	HERBICIDE

Amrita

SPECIALTY HERBICIDE

For control of susceptible weeds and certain woody plants, including invasive and noxious weeds, on rangeland, permanent grass pastures (including grasses grown for hay), Conservation Reserve Program (CRP) acres, non-cropland areas including industrial sites, rights-of-way (such as roadsides, electric utility and communication transmission lines, pipelines, and railroads) and non-irrigation ditch banks, natural areas (such as wildlife management areas including seasonally dry flood plains, deltas, marshes, prairie potholes, or vernal pools, wildlife openings, wildlife habitats, recreation areas, campgrounds, trailheads and trails), and grazed areas in and around these sites

Not For Sale, Distribution, or Use in New York State.

Not For Sale, Distribution, or Use in the San Luis Valley of Colorado.

Active Ingredients:	% w/w
Aminopyralid, Trisopropanolammonium salt: Trisopropanolammonium salt of 2-pyridine carboxylic acid, 4-amino-3,6-dichloro	8.34%
2,4-D, dimethylamine salt: Dimethyl amine salt of (2,4-dichlorophenoxy) acetic acid	41.96%
Other Ingredients	49.70%
Total	100.00%
Acid Equivalents: aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) – 4.3% - 0.41 lb./gal. (50 g/L) 2,4-D [(2,4-dichlorophenoxy) acetic acid] – 34.85% – 3.27 lbs./gal. (400 g/L)	

KEEP OUT OF REACH OF CHILDREN DANGER

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the poison control center at 1-800-222-1222 for emergency medical treatment information. **Note to Physician:** Probable mucosal damage may contraindicate the use of gastric lavage.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the "Directions for Use" section for information about this standard.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS PELIGRO

PRIMEROS AUXILIOS

Si entra en contacto con los ojos: Mantenga los ojos abiertos y enjuáguelos lenta y cuidadosamente con agua, durante 15 a 20 minutos. Si utiliza lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento. **Si se ingiere:** Llame de inmediato a un centro de control de envenenamientos o a un médico para consejo de tratamiento. Si la persona puede tragar, haga que beba un vaso de agua lentamente. No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. No administre nada por boca a una persona que haya perdido el conocimiento. **NÚMERO DE TELÉFONO DIRECTO:** Cuando llame a un centro de control de envenenamiento, o a un médico, o intentante obtener tratamiento, tenga a la mano el recipiente o la etiqueta del producto. También puede comunicarse con el centro de control de intoxicaciones al 1-800-222-1222 para obtener información sobre tratamiento médico de emergencia. **NOTA AL MÉDICO:** El probable daño de las mucosas puede contraindicar el uso de un lavado gástrico.

Refer to inside of label booklet for additional precautionary information including Directions for Use. Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product call 1-800-262-8200.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Produced For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 83529-253

EPA Est. No. **AG** 72159-GA-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
DANGER

Corrosive. Causes Irreversible Eye Damage. Harmful if Swallowed. Do not get in eyes or on clothing.

Personal Protective Equipment (PPE)

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton \geq 14 mils, when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See **Engineering Controls** for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607 (d-e)] the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607 (f)].

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Take care to minimize the incidental overspray along the shoreline when applying to terrestrial plants at the water's edge or to water in areas where surface water is present. Do not apply directly to intertidal areas below the mean high-water mark, except as permitted on this label. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory

2,4-D and Aminopyralid are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Users are advised not to apply these chemicals where soils have a rapid to very rapid permeability (such as loamy sand to sand) and the water table of an underlying aquifer is shallow or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching both surface water and aquatic sediment via runoff for several months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of aminopyralid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Non-Target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Irrigation Water Statement

Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not apply to snow or frozen ground.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying. This product is not intended for reformulation or repackaging into other end-use products.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not For Sale, Distribution, or Use in New York State.

Not For Sale, Distribution, or Use in the San Luis Valley of Colorado.

Not for use on pastures in Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, and Vermont. All other labeled uses are permitted in these states including grazed areas in and around these sites.



Light Gray = states where use in pastures is not permitted
Dark Gray = NY where the product is not registered

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton \geq 14 mils.
- Protective eyewear
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, feed or fertilizer by storage or disposal.

Pesticide Storage: If this product is exposed to subfreezing temperatures, the active ingredient may crystallize and settle out of solution. Under these conditions the product should be warmed to at least 40°F and agitated well to dissolve any crystallized material prior to use.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Non-refillable containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Nonrefillable containers 5 gallons or larger:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

PRODUCT INFORMATION

Amrita specialty herbicide controls broadleaf weeds and certain woody plants, including invasive and noxious weeds, on rangeland, permanent grass pastures (including grasses grown for hay), Conservation Reserve Program (CRP) acres, non-cropland areas including industrial sites, rights-of-way (such as roadsides, electric utility and communication transmission lines, pipelines, and railroads) and non-irrigation ditch banks, natural areas (such as wildlife management areas including seasonally dry flood plains, deltas, marshes, prairie potholes, or vernal pools, wildlife openings, wildlife habitats, recreation areas, campgrounds, trailheads and trails), and grazed areas in and around these sites.

Consult with a Sharda USA LLC representative if you do not understand the Precautions and/or Restrictions.

WEED RESISTANCE MANAGEMENT

For resistance management, **Amrita** is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to **Amrita** and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of **Amrita** or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.

- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weed), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plant mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

MIXING AND APPLICATION INSTRUCTIONS

Mixing Instructions

Mixing with Water

To prepare the spray, add about half the required amount of water in the spray tank. Then, with agitation, add the specified amount of **Amrita** and other registered tank mix herbicides. Finally, with continued agitation, add the rest of the water and additives such as surfactants or drift control and deposition aids.

Addition of Surfactants or Adjuvants on All Labeled Use Sites: The addition of a high quality non-ionic surfactant (of at least 80% active ingredient) at 0.25 to 0.5% volume per volume (1 to 2 quarts per 100 gallons of spray) is recommended to enhance herbicide activity under adverse environmental conditions (such as, high temperature, low relative humidity, drought conditions, dusty plant surfaces) or when weeds are heavily pubescent or more mature.

Tank Mixing with Other Herbicides

Amrita at rates of up to 2.1 pints (34 fl. oz.) per acre may be mixed with labeled rates of other labeled herbicides to broaden the spectrum of weeds controlled or to improve control of certain weeds. **Amrita** may be applied in tank mix combination with labeled rates of other herbicides provided: (1) the tank mix product is labeled for the timing and method of application for the use site to be treated and (2) mixing is not prohibited by the label of the registered tank mixed products, and (3) that the tank mix combination is physically compatible (see **Tank Mix Compatibility Testing** below). When tank mixing, use only in accordance with the restrictions, precautions and limitations on the respective product labels.

- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Do not exceed specified application rates. If products containing the same active ingredient are mixed, do not exceed the maximum allowable active ingredient use rates.
- For direct injection or other spray equipment where the product formulations will be mixed in undiluted form, special care should be taken to ensure tank mix compatibility.
- Always perform a jar test to ensure the compatibility of products to be used in tank mixture.

Tank Mixing Precautions:

- For products packaged in water soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment has been adequately cleaned. (See **Sprayer Clean-Out Instructions**.)
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

Tank Mix Compatibility Testing: Perform a jar test prior to mixing in a spray tank to ensure compatibility of **Amrita** and other pesticides or carriers. Use a clear glass jar with lid and mix ingredients in the same order and proportions as will be used in the spray tank. The mixture is compatible if the materials mix readily when the jar is inverted several times. The mixture should remain stable after standing for 1/2 hour or, if separation occurs, should readily mix if agitated. An incompatible mixture is indicated by separation into distinct layers that do not readily remix when agitated and/or the presence of flakes, precipitates, gels, or heavy oily film in the jar. Use of an appropriate compatibility agent may resolve mix incompatibility.

Mixing with Sprayable Liquid Fertilizer Solutions: **Amrita** is usually compatible with liquid fertilizer solutions. It is anticipated that **Amrita** will not require a compatibility agent for mixing with fertilizers; however, a compatibility test (jar test) should be made prior to large scale batch mixing. Jar tests are particularly important when a new batch of fertilizer or pesticide is used, when water sources change, or when tank mixture ingredients or concentrations are changed. Compatibility may be determined by mixing the spray components in the desired order and proportions in a clear glass jar before large scale mixing of spray components in the spray tank. When required, a compatibility agent could be used to help obtain and maintain a uniform spray solution during mixing and application. **Note:** The lower the temperature of the liquid fertilizer, the greater the likelihood of mixing problems. **Mixing Amrita in N-P or N-P-K liquid fertilizer solutions is more difficult than mixing with straight nitrogen fertilizer and should not be attempted without first conducting a successful compatibility jar test.** Agitation in the spray tank must be vigorous to be comparable with jar test agitation. Apply the spray mixture the same day it is prepared while maintaining continuous agitation. Rinse the spray tank thoroughly after use.

Suggested Mixing Procedure:

1. With continuous vigorous agitation, dilute herbicide with water (1 part herbicide to 2 parts water) before adding to liquid nitrogen fertilizer solution.
2. Apply as soon as mixing is complete, maintaining continuous, vigorous agitation throughout mixing and application without interruption.
3. Application during very cold (near freezing) weather is not advisable. The likelihood of mixing or compatibility problems with liquid fertilizer increases under cold conditions.
4. Do not store the spray mixture.

Note: Foliar-applied liquid fertilizers themselves can cause injury (such as: yellowing and burning) to the foliage of forage grasses and other vegetation especially in the summer. The addition of a surfactant to fertilizer blends may increase the injury potential.

MANDATORY SPRAY DRIFT MANAGEMENT**Aerial Applications:**

- Do not release spray at a height greater than 10 ft. above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select a nozzle and pressure combination that delivers a medium or coarser droplet size (ASABE S641).
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wing-span for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- If the windspeed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 - 15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
- If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.
- Only apply this product if the wind direction favors on-target deposition and there are no sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.
- Do not apply during temperature inversions.

Ground Boom Applications:

- For applications on pastures and rangeland, do not release spray at a height greater than 4 ft. above the ground. For all other uses, do not release spray at a height greater than 3 ft. above the ground or crop canopy.
- Applicators are required to select a nozzle and pressure combination that delivers a medium or coarser droplet size (ASABE S572).
- Do not apply when wind speeds exceed 15 mph at the application site.
- If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.
- Only apply this product if the wind direction favors on-target deposition and there are no sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.
- Do not apply during temperature inversions.

Boom-less Ground Sprayer Applications:

- Applicators are required to select a nozzle and pressure combination that delivers a medium or coarser droplet size (ASABE S572) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

- An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstance where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

WIND

Drift potential generally increases with wind speed.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.


Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

IMPORTANT USE PRECAUTIONS AND RESTRICTIONS TO PREVENT INJURY TO DESIRABLE PLANTS

- Carefully read the section **Restrictions in Hay or Manure Use**.
- It is mandatory to follow the **Use Precautions** and **Use Restrictions** section of this label.
- Manure and urine from animals consuming grass or hay treated with this product may contain enough aminopyralid to cause injury to sensitive broad-leaf plants.
- Hay can only be used on the farm or ranch where product is applied.
- Consult with a Sharda USA LLC representative if you do not understand the **Use Precautions** and **Use Restrictions**.
- Warning: Do not move treated plant materials or manure from animals who have grazed on treated plant materials to sites where manure may be collected or sensitive crops are grown.

Forage and Manure Management



USE RESTRICTIONS

- This product is persistent and may be present in treated plant materials for months to years after application.
- Do not sell or transport treated plant materials or manure from animals that have grazed on treated plant materials off-site for compost distribution or for use as animal bedding/feed for 18 months after application. Treated plant materials can be recycled on-site or left in the field to decompose.
- Manure from animals that have grazed or eaten forage or hay harvested from treated areas within the previous three days may only be applied to the fields where the following crops will be grown: pasture grasses and grass grown for seed.
- Animals that have been fed **Amrita**-treated forage must be fed forage free of aminopyralid for at least 3 days before movement to an area where manure may be collected, or sensitive crops are grown.
- For more information on how to manage aminopyralid treated materials and to prevent aminopyralid from contaminating compost please visit: <https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides>
- Do not sell or distribute hay treated with **Amrita** in the preceding 18 months off the farm or ranch where harvested.
- Do not move hay and silage made from grass treated with **Amrita** within the preceding 18 months off the farm or ranch.
- Do not use hay or straw from areas treated with **Amrita** within the preceding 18 months or manure from animals feeding on hay treated with **Amrita** in compost.
- Do not use hay from areas treated with **Amrita** in the preceding 18 months for silage, haylage, baleage, or green chop.
- Do not use grasses treated with **Amrita** in the preceding 18 months for seed production.
- Do not use grasses treated with **Amrita** in the preceding 18 months for hay intended for export outside the United States.

• **For applications to pasture:**

- o The applicator must document that they have notified property owners/operators, or customers, in writing, of the compost and animal bedding/feed prohibitions within 14 days of the application. Applicators must keep the records of notification for two years. This record must include date of application, the name of the applicator, the EPA registration number of the product applied, the area(s) treated, and a copy of the written notification provided to the property owner/operator. Notification may be made via email, mail, paper handout, or by any other written communication method. Records must be made available to State Pesticide Regulatory Official(s), and to EPA upon request. If this information is already being retained, duplicate records are not needed.
- o It is recommended that applicators also transmit at the time of notification relevant educational materials for managing treated plant matter, as available. Additional educational materials for aminopyralid will be posted at: <https://www.epa.gov/ingredients-used-pesticide-products/registration-review-pyridine-and-pyrimidine-herbicides>
- o Applications to pasture by property owners/operators on their own property are exempt from this notification and record keeping requirement.
- o Applications to pasture on public land (i.e., lands managed directly by state, tribal, or local authorities) are exempt from this notification requirement.
- Do not apply this product on lawns, turf, ornamental plantings, urban walkways, driveways, tennis courts, golf courses, athletic fields, commercial sod operations, or other high-maintenance, fine turfgrass areas, or similar areas.
- Do not use this product for impregnation on dry bulk fertilizer.
- Do not apply this product through any type of irrigation system.
- Do not contaminate water intended for irrigation or domestic purposes. Do not treat inside banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes.
- Do not treat frozen soil where runoff could damage sensitive plants.
- Untreated trees can occasionally be affected by root uptake of **Amrita** through movement into the soil or by excretion of the product from the roots of nearby treated trees. Do not apply **Amrita** within the root zone of desirable trees.
- **Crop Rotation:** Do not rotate non-cropland to cropland for one year following an application of **Amrita**. Do not plant a broadleaf crop until an adequately sensitive field bioassay shows that the level of **Amrita** present in the soil will not adversely affect that broadleaf crop.
- **Seeding Legumes or Wildflowers:** Do not plant legumes or wildflowers until a soil bioassay has been conducted to determine if residues of **Amrita** remaining in the soil will adversely affect establishment of legumes and wildflowers.
- **Restrictions in Hay or Manure Use:**
 - o Do not use treated plant residues, including hay or straw from areas treated within the preceding 18-months, in compost, mulch or mushroom spawn.
 - o Do not use manure from animals that have grazed forage or eaten hay harvested from treated areas within the previous 3 days, in compost, mulch or mushroom spawn.
 - o Do not spread manure from animals that have grazed or consumed forage or hay from treated areas within the previous 3 days on land used for growing susceptible broadleaf crops.
 - o Manure from animals that have grazed forage or hay harvested from **Amrita**-treated areas within the previous 3 days may only be spread on pasture grasses and grass grown for seed.
 - o Do not plant a broadleaf crop (including soybeans, sunflower, tobacco, vegetables, field beans, peanuts, and potatoes) in fields treated with manure from animals that have grazed forage or eaten hay harvested from **Amrita**-treated areas until an adequately sensitive field bioassay is conducted to determine that the aminopyralid concentration in the soil is at level that is not injurious to the crop to be planted.
 - o Do not plant a broadleaf crop in fields treated in the previous year with manure from animals that have grazed forage or hay harvested from **Amrita**-treated areas until an adequately sensitive field bioassay is conducted to determine that the **Amrita** residues in the soil is at a level that is not injurious to the crop to be planted.
- **Grazing and Haying Restrictions:** There are no restrictions on grazing or hay harvest following application of **Amrita** at labeled rates. Do not transfer grazing animals from areas treated with **Amrita** to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough **Amrita** to cause injury to broadleaf plants.
- **Grazing Poisonous Plants:** Herbicide application may increase palatability of certain poisonous plants. Do not graze treated areas until poisonous plants are dry and no longer palatable to livestock.
- **Maximum Application Rate:** Do not broadcast apply more than 2.1 pints (34 fl. oz.) (0.86 lb. a.i. 2,4-D and 0.11 lb. a.i. aminopyralid) per acre of **Amrita** per year. The total amount of **Amrita** applied broadcast, as a re-treatment, and/or spot treatment per year must not exceed 2.1 pints (34 fl. oz.) (0.86 lb. a.i. 2,4-D and 0.11 lb. ae aminopyralid) per acre. Spot treatments may be applied at an equivalent broadcast rate of up to 4.2 pints (68 fl. oz.) of **Amrita** (1.7 lbs. a.i. 2,4-D and 0.22 lb. a.i. aminopyralid) per acre per year; however, not more than 50% of an acre may be treated at that rate.

USE PRECAUTIONS

- Applications made during periods of intense rainfall, to soils saturated with water, surfaces paved with materials such as asphalt or concrete, or soils through which rainfall will not readily penetrate may result in runoff and movement of **Amrita**. Injury to crops may result if treated soil and/or runoff water containing **Amrita** is washed, or moved onto land used to produce crops. Exposure to **Amrita** may injure or kill susceptible crops and other plants, such as grapes, soybeans, tobacco, sensitive ornamentals.
- **Seeding grasses:**
 - **Preemergence:** Grasses may be reseeded in the fall following an application of **Amrita**—applied in the spring or early summer.
 - **Postemergence:** During the season of establishment, **Amrita** should be applied only after perennial grasses are well established (have developed a good secondary root system and show good vigor). Most perennial grasses are tolerant to **Amrita** at this stage of development. **Amrita** may suppress certain established grasses, such as smooth brome-grass (*Bromus inermis*), especially when plants are stressed by adverse environmental conditions. Plants should recover from this transient suppression with the onset of environmental conditions favorable to grass growth and upon release from weed competition.
- **Field Bioassay Instructions:** In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample variability in field conditions such as soil texture, soil organic matter, soil pH, rainfall pattern or drainage. The field bioassay can be initiated starting a minimum of one year after herbicide application and following harvest of the treated crop. Observe the test crop for symptoms of herbicidal activity, such as poor stand (effect on seed germination), chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the intended rotational crop; plant only to wheat, forage grasses, native grasses or grasses grown for hay.

NON-CROPLAND AND NATURAL AREAS

Amrita may be applied alone or in tank mix combination to non-cropland areas, such as non-irrigation ditch banks, industrial and storage areas, airports, roadsides, railroad and utility rights-of-way, including grazed areas on these sites as an aerial or ground broadcast treatment, as a spot application, or as a high volume foliar application (see **Application Methods** section). Refer to the **Broadleaf Weeds Controlled** section for application rates listed for specific broadleaf weeds.

Restrictions:

Postemergence (annual and perennial weeds):

- Limited to 2 applications per year.
- Maximum of 2.1 pints (34 fl. oz.) (0.86 lb. a.i./acre 2,4-D and 0.11 lb. a.i./acre aminopyralid)/acre per application.
- Do not apply more than 2.1 pints per acre per year.
- Minimum of 30 days between applications.

Postemergence (woody plants)

- Limited to 1 application per year.
- Maximum of 2.1 pints (34 fl. oz.) (0.86 lb. ae/acre 2,4-D and 0.11 lb. ae/acre aminopyralid)/acre per year.

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Restrictions for Non-Irrigation Canal Ditchbank Application and Terrestrial Weeds near and up to the Water's Edge

Use Rate Restriction:

- Limited to 2 application per year.
- Minimum of 30 days between application.
- Maximum of 2.1 pints (34 fl. oz.) (0.86 lb. a.i./acre 2,4-D and 0.11 lb. a.i./acre aminopyralid)/acre per broadcast application.
- Do not apply more than 2.1 pints per acre per year.

Spot treatments may be applied at an equivalent broadcast rate of up to 4.2 pints (64 fl. oz.) of **Amrita** per acre per year; however, not more than 50 % of an acre may be treated at that rate.

Do not use on small canals with a flow rate less than 10 cubic feet per second (CFS) where water will be used for drinking purposes. CFS may be estimated by using the formula below. The approximate velocity needed for the calculation can be determined by observing the length of time that it takes a floating object to travel a defined distance.

Divide the distance (ft.) by the time (sec.) to estimate velocity (ft. per sec.). Repeat 3 times and use the average to calculate CFS.

Average Width (ft.) x Average Depth (ft.) x Average Velocity (ft. per sec.) = CFS

For ditchbank weeds:

Do not allow boom spray to be directed onto water surface. Do not spray across stream to opposite bank.

For shoreline weeds:

Allow no more than 2 foot overspray onto water.

Sprayer Clean-Out Instructions

It is recommended that separate spray equipment be used on highly sensitive crops such as tobacco, soybeans, peanuts, and tomatoes.

Do not use spray equipment used to apply **Amrita** for other applications to land planted to, or to be planted to, crops or desirable sensitive plants, unless it has been determined that all residues of this herbicide have been removed by thorough cleaning of equipment.

Equipment used to apply **Amrita** should be thoroughly cleaned before reusing to apply any other chemicals as follows.

1. Rinse and flush application equipment thoroughly after use. Dispose of rinse water away from water supplies.
2. Rinse a second time, adding 1 quart of household ammonia or tank cleaning agent for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 minutes). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Spray nozzles and screens should be removed and cleaned separately.

APPLICATION METHODS

Apply the specified rate of **Amrita** as a coarse low-pressure spray. **Do not apply this product with mist blower systems that deliver very fine spray droplets. Use of mist blower equipment can reduce weed control and increase spray drift potential.**

Spray volume should be sufficient to uniformly cover foliage. Increase spray volume to ensure thorough and uniform coverage when target vegetation is tall and/or dense. To enhance foliage wetting and coverage, an approved non-ionic agricultural surfactant may be added to the spray mixture as specified by the surfactant label.

Ground Broadcast Application: Higher spray volumes (greater than 10 gallons per acre) generally provide better coverage and better control, particularly in dense and/or tall foliage.

Aerial Broadcast Application: Do not apply less than 2 gallons per acre total spray volume. Five gallons per acre or greater will generally provide better coverage and better control, particularly in dense and/or tall foliage.

High-Volume Foliar Application: High volume foliar treatments may be applied at rates equivalent to a maximum of 2.1 pints (34 fl. oz.) per acre per annual growing season. Use sufficient spray volume to thoroughly and uniformly wet foliage and stems.

Spot Application: Spot treatments may be applied at rates equivalent to broadcast-applied rate of up to a maximum of 4.2 pints (68 fl. oz.) per acre on 50% of the treated field. Spray volume should be sufficient to thoroughly and uniformly wet weed foliage. Repeat treatments may be made, but the total amount of **Amrita** applied must not exceed 2.1 pints (34 fl. oz.) per acre per year (see comments in the **Use Precautions and Restrictions** section above on **Maximum Application Rate**).

Table 1: Amount of **Amrita** herbicide (in fl. oz.) to mix in 3 gallons of water.

Amrita Amount (In fl. oz.) To Mix In 3 Gals. Of Water With Various Application Rates

GPA	19 fl. oz./A	24 fl. oz./A	34 fl. oz./A
20	2.9	3.6	5.1
30	1.9	2.4	3.4
40	1.4	1.8	2.6
50	1.1	1.4	2.0
60	1.0	1.2	1.7
70	0.8	1.0	1.4
80	0.7	0.9	1.3
90	0.6	0.8	1.1
100	0.6	0.7	1.0

Table 2: Application rates in the table below are based on treating an area of 1000 sq. ft. An area of 1000 sq. ft. is about 10.5 by 10.5 yards in size. Mix the amount of **Amrita** (fl. oz. or milliliters) corresponding to the desired broadcast rate in 0.5 to 2.5 gallons of water, depending upon the spray volume required to treat 1000 sq. ft. A delivery volume of 0.5 gallons per 1000 sq. ft. is equivalent to 22 gallons per acre and 2.5 gallons per 1000 sq. ft. is equivalent to 109 gallons per acre.

Amount of Amrita per 1000 sq. ft. to Equal Broadcast Rate			
Broadcast Rate		Amount of Amrita per 1000 sq. ft.	
(fl. oz./acre)	(pints/acre)	(fl. oz.)	(mL)
19	1.2	0.44	13
24	1.5	0.55	16
34	2.1	0.78	23

Note: 1 mL = 1cc and 1 fluid ounce (fl. oz.) = 29.6 milliliters (mL) = 2 tablespoons = 6 teaspoons

To calculate the amount of Amrita for areas larger than 1000 sq. ft.: Multiply the table value (fl. oz. or milliliters) by the area to be treated in “thousands” of square feet. For example, if the area to be treated is 3500 sq. ft., multiply the table value by 3.5 (3500 sq. ft. divided by 1000 sq. ft. = 3.5).

USE RATES AND TIMING

Do not use Amrita if loss of legumes species or other broadleaf species cannot be tolerated.

Amrita may be applied postemergence as a broadcast spray or as a spot application to control weeds listed on this label; weeds other than those listed may also be controlled by this herbicide. When a rate range is given, use a higher rate in the range to control weeds at advanced growth stages or under less-than-favorable growing conditions (e.g., drought stress) or for longer residual control. Best weed control results are obtained when spray volume is sufficient to provide uniform coverage of treated plants. For optimum uptake and translocation of the herbicide, avoid mowing, haying, shredding, burning or soil disturbance in treated areas for at least 7 days following application.

For most species, 2 hours between application and rainfall provides a sufficient amount of time to avoid loss in weed control due to herbicide wash-off of foliage.

Amrita also provides preemergence control of germinating seeds or emerging seedlings of susceptible weeds and re-growth of certain perennial weeds following application. Weed establishment following **Amrita** application will depend upon application rate, season of application, and growing condition.

Amrita can provide long-term control of weeds. The length of control is dependent upon the application rate, condition and growth stage of target weeds, environmental conditions at and following application, and the density and vigor of competing desirable vegetation. Long-term broadleaf weed control is most effective where forage grasses are allowed to recover from overgrazing, drought, etc., and compete with weeds.

Amrita can be an important component of integrated vegetation management programs designed to renovate or restore desired non-cropland plant communities. To maximize and extend the benefits of weed control provided by **Amrita**, it is important that vegetation management practices, including grazing management, biological control agents, replanting, fertilization, prescribed fire, reseeding with desirable plants, etc., be used to increase the competitiveness of desired forages. Used as part of an integrated management program, **Amrita** can serve as a catalyst for rapid improvement of rangeland, permanent grass pasture, and CRP, and non-cropland sites by alleviating the adverse competitive effect of weeds on the yield and quality of forages and other desirable plant species.

Agricultural and natural resources specialists with federal and state government agencies can provide guidance on best management practices and development of integrated vegetation management systems.

Broadleaf Weeds Controlled

The following weeds will be controlled at 1.2 to 2.1 pints (19 to 34 fl. oz.) per acre. For best results, apply when weeds are actively growing and conditions favorable for plant growth. Use a higher rate in the rate range when growing conditions are less than favorable, when weeds are mature, or when weed foliage is tall and dense or when residual control is important. **Amrita** also provides preemergence control of germinating seeds or emerged seedlings of susceptible weeds following application.

Table 3: Broadleaf Weeds Controlled

Weed Species			
Common Name	Scientific Name	Life Cycle***	Plant Family
Rate Range: 1.2 to 1.5 pints (19 to 24 fl. oz.) per acre			
bedstraw	<i>Galium</i> spp.	perennial	Rubiaceae
bedstraw, smooth	<i>Galium mollugo</i>	perennial	Rubiaceae
beggerticks	<i>Bidens</i> spp.	annual	Asteraceae
carrot, wild*	<i>Daucus carota</i>	biennial	Apiaceae
chamomile, scentless	<i>Matricaria inodora</i>	annual	Asteraceae
Cinquefoil, hoary	<i>Potentilla argentea</i>	perennial	Rosaceae
cinquefoil, sulfur**,**	<i>Potentilla recta</i>	perennial	Rosaceae
clover, sweet	<i>Melilotus officinalis</i>	biennial	Fabaceae
clover, white	<i>Trifolium repens</i>	perennial	Fabaceae
crownvetch*	<i>Securigera varia</i>	perennial	Fabaceae
daisy, oxeye**,**	<i>Leucanthemum vulgare</i>	perennial	Asteraceae
false dandelion, Carolina*	<i>Pyrrophappus carolinianus</i>	annual/biennial	Asteraceae
gumweed, curlycup	<i>Grindelia squarrosa</i>	biennial	Asteraceae
horsenettle, Carolina**,**	<i>Solanum carolinense</i>	perennial	Solanaceae
jimsonweed	<i>Datura stramonium</i>	annual	Solanaceae
pokeweed, common	<i>Phytolacca americana</i>	perennial	Phytolaccaceae
ragweed, common**,**	<i>Ambrosia artemisiifolia</i>	annual	Asteraceae
ragweed, western	<i>Ambrosia psilostachya</i>	perennial	Asteraceae
ragwort, tansy**,**	<i>Senecio jacobaea</i>	perennial	Asteraceae
starthistle, yellow**,**	<i>Centaurea solstitialis</i>	annual	Asteraceae
thistle, bull**,**	<i>Cirsium vulgare</i>	biennial	Asteraceae
thistle, musk**,**	<i>Carduus nutans</i>	biennial	Asteraceae
thistle, plumeless**,**	<i>Carduus acanthoides</i>	biennial	Asteraceae
thistle, scotch	<i>Onopordum acanthium</i>	biennial	Asteraceae
tickclover	<i>Desmodium</i> sp.	perennial	Fabaceae
vetch, common*	<i>Vicia sativa</i>	annual	Fabaceae
woodsorrel, yellow*	<i>Oxalis stricta</i>	perennial	Oxalidaceae
wormwood, absinth**,**	<i>Artemisia absinthium</i>	perennial	Asteraceae
Rate Range: 1.5 to 2.1 pints (24 to 34 fl. oz.) per acre			
actinomeris, wingstem	<i>Verbesina alternifolia</i>	perennial	Asteraceae
amaranth, spiny*	<i>Amaranthus spinosus</i>	annual	Amaranthaceae
broomweed, annual*	<i>Amphichyris dracunculoides</i>	annual	Asteraceae

(continued)

Weed Species			
Common Name	Scientific Name	Life Cycle***	Plant Family
Rate Range: 1.5 to 2.1 pints (24 to 34 fl. oz.) per acre (continued)			
burdock, common*,**	<i>Arctium minus</i>	biennial	Asteraceae
buttercup, hairy*	<i>Ranunculus sardous</i>	perennial	Ranunculaceae
buttercup, tall*,**	<i>Ranunculus acris</i>	perennial	Ranunculaceae
camphorweed*	<i>Heterotheca subaxillaris</i>	annual	Asteraceae
cat's ear, common	<i>Hypochaeris radicata</i>	perennial	Asteraceae
chickweed, common*	<i>Stellaria media</i>	annual	Caryophyllaceae
chicory*	<i>Cichorium intybus</i>	perennial	Asteraceae
cocklebur*	<i>Xanthium strumarium</i>	annual	Asteraceae
croton, woolly*,**	<i>Croton capitatus</i>	annual	Euphorbiaceae
cudweed, purple	<i>Gnaphalium purpureum</i>	annual	Asteraceae
dandelion, common*	<i>Taraxacum officinale</i>	perennial	Asteraceae
deadnettle, purple	<i>Lamium purpureum</i>	annual/biennial	Lamiaceae
dock, broadleaf*	<i>Rumex obtusifolius</i>	perennial	Polygonaceae
dock, curly*	<i>Rumex crispus</i>	perennial	Polygonaceae
dogfennel***	<i>Eupatorium capillifolium</i>	perennial	Asteraceae
evening primrose, cutleaf*	<i>Oenothera lacinata</i>	annual	Asteraceae
false dandelion, Carolina*	<i>Tragopogon dubius</i>	biennial	Asteraceae
fiddleneck, common	<i>Amsinckia intermedia</i>	annual	Boraginaceae
fireweed	<i>Epilobium angustifolium</i>	perennial	Onagraceae
fleabane, annual*	<i>Erigeron annuus</i>	annual	Asteraceae
goldenrod, Canada*	<i>Solidago canadensis</i>	perennial	Asteraceae
goldenrod, Missouri*	<i>Solidago missouriensis</i>	perennial	Asteraceae
goldenrod, rigid	<i>Solidago rigida</i>	perennial	Asteraceae
hawkweed, orange*,**	<i>Hieracium aurantiacum</i>	perennial	Asteraceae
hawkweed, yellow*,**	<i>Hieracium pratense</i>	perennial	Asteraceae
henbit*	<i>Lamium amplexicaule</i>	annual/biennial	Lamiaceae
horseweed*	<i>Conyza canadensis</i>	annual	Asteraceae
ironweed, tall	<i>Vernonia gigantea</i>	perennial	Asteraceae
ironweed, western	<i>Vernonia baldwinii</i>	perennial	Asteraceae
knapweed	<i>Centaurea</i> sp.	biennial	Asteraceae
knapweed, brown	<i>Centaurea jacea</i>	perennial	Asteraceae
knapweed, diffuse*,**	<i>Centaurea diffusa</i>	biennial	Asteraceae
knapweed, Russian*,**	<i>Acrotilon repens</i>	perennial	Asteraceae
knapweed, spotted*,**	<i>Centaurea stoebe</i>	biennial	Asteraceae
kudzu*,**	<i>Pueraria montana</i>	perennial	Fabaceae

(continued)

Weed Species			
Common Name	Scientific Name	Life Cycle***	Plant Family
Rate Range: 1.5 to 2.1 pints (24 to 34 fl. oz.) per acre (continued)			
lambquarters, common*	<i>Chenopodium album</i>	annual	Chenopodiaceae
lespedeza, annual	<i>Lespedeza striata</i>	annual	Fabaceae
lettuce, prickly*	<i>Lactuca serriola</i>	annual	Asteraceae
locust	<i>Robinia pseudoacacia</i>	perennial	Fabaceae
marshelder, annual*	<i>Iva annua</i>	annual	Asteraceae
mayweed, scentless*	<i>Tripleurospermum perforata</i>	annual	Asteraceae
mayweed, stinking*,**	<i>Anthemis cotula</i>	annual	Asteraceae
medic, black*	<i>Medicago lupulina</i>	perennial	Fabaceae
mexican tea	<i>Dysphania ambrosioides</i>	annual/perennial	Chenopodiaceae
mullein****	<i>Verbascum</i> spp.	biennial	Scrophulariaceae
mugwort	<i>Artemisia vulgaris</i>	Perennial	Asteraceae
partridgepea*	<i>Chamaecrista fasciculata</i>	annual	Fabaceae
plantain, broadleaf*	<i>Plantago major</i>	perennial	Plantaginaceae
plantain, buckhorn*	<i>Plantago lanceolata</i>	perennial	Plantaginaceae
sicklepod*	<i>Senna obtusifolia</i>	annual	Fabaceae
smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>	annual	Polygonaceae
sneezeweed, bitter*	<i>Helenium amarum</i>	annual	Asteraceae
soda apple, tropical*,**	<i>Solanum viarum</i>	perennial	Solanaceae
sowthistle, perennial*,**	<i>Sonchus arvensis</i>	perennial	Asteraceae
sowthistle, prickly*	<i>Sonchus asper</i>	annual	Asteraceae
Spanish needles	<i>Bidens bipinnata</i>	annual	Asteraceae
starthistle, yellow*,**	<i>Centaurea solstitialis</i>	annual	Asteraceae
St. Johnswort, common	<i>Hypericum perforatum</i>	perennial	Ciusiaceae
sunflower, common*	<i>Helianthus annua</i>	annual	Asteraceae
teasel*	<i>Dipsacus</i> spp.	biennial	Dipsacaceae
thistle, Canada*,**	<i>Cirsium arvense</i>	perennial	Asteraceae
vervain, blue*	<i>Verbena hastata</i>	perennial	Asteraceae
vervain, hoary*	<i>Verbena stricta</i>	perennial	Asteraceae
yarrow, common*	<i>Achillea millefolium</i>	perennial	Asteraceae

*These plants are indicated to be invasive in the USDA-NRCS, PLANTS Database (<http://plants.usda.gov/index.html>).

**Plants designated as noxious weeds in at least one state (PLANTS Database, USDA-NRCS, <http://plants.usda.gov/index.html>).

***Spot treatment at rates up to 4.2 pints (68 fl. oz.) per acre of **Amrita** may be particularly effective against dense patches of perennial broadleaf plants.

****Apply during rosette stage.

Susceptible Brush such as Multiflora Rose:

Individual Plant Treatment - Use 24 fl. oz. (0.25% v/v) of **Amrita** tank-mixed with Remedy® Ultra (triclopyr, butoxyethyl ester EPA Reg. # 62719-552) at the appropriately labeled use rate per 100 gallons of water. Apply from full leaf through flowering. For best results, delay treatment for 9 - 12 months after mowing. Spot treatments may be applied at an equivalent broadcast rate of up to 4.2 pints (68 fl. oz.) of **Amrita** per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate.

Broadcast - Use 1.2 to 2.1 pints (19 to 34 fl. oz.) of **Amrita** tank mixed with Remedy Ultra at the appropriately labeled use rate per acre. Apply from full leaf through flowering. For best results, delay treatment for 9 - 12 months after mowing.

Control of Terrestrial Weeds near and up to the Water's Edge

Amrita can be used to treat terrestrial weeds that extend up to the water's edge. **Do not apply directly to water.** This product must not be used to treat vegetation standing in the water. When controlling terrestrial weed species near and up to the water's edge, take precautions to minimize incidental overspray to the adjacent water. Consult local public water control authorities before applying this product near public waters. Permits may be required to treat such areas. Apply the specified rate, listed in **Table 2**, of **Amrita** as a coarse low-pressure spray as ground broadcast or spot applications. Do not apply aerially for control of weeds growing at or near the water's edge. Spray volume should be sufficient to uniformly cover foliage. Increase the spray volume to ensure thorough and uniform coverage when target vegetation is tall and/or dense. It is also permissible to treat target weeds within dry non-irrigation ditches and seasonally dry transitional areas between upland and lowland sites (such as flood plains, deltas, marshes, prairie potholes or vernal pools), but only at times when those sites are dry and are forecasted or managed by water control systems to remain dry for at least 2 weeks following application.

Use Rate Restrictions:

Limited to 2 application per year.

Minimum of 30 days between applications.

Maximum of 2.1 pints (34 fl. oz.) per acre per broadcast application.

Do not apply more than 2.1 pints per acre per year.

Spot treatments may be applied at an equivalent broadcast rate of up to 4.2 pints (64 fl. oz.) of **Amrita** per acre per year; however, not more than 50% of an acre may be treated at that rate.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

WARRANTY DISCLAIMER

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Sharda USA LLC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Sharda USA LLC or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Sharda USA LLC election's, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Sharda USA LLC shall not be liable for losses or damages resulting from handling or use of this product unless Sharda USA LLC is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Sharda USA LLC be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Sharda USA LLC or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

DECLARACIONES PREVENTIVAS RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS PELIGRO

Corrosivo. Causa daño ocular irreversible. Letal si se ingiere. No permita que caiga en los ojos o la ropa.

Equipo de Protección Personal (PPE, por sus siglas en inglés)

Todos los mezcladores, cargadores, bandereros y otros manipuladores deben usar:

- Camisa de manga larga y pantalones largos
 - Zapatos y calcetines
 - Lentes o gafas protectoras
 - Guantes resistentes a productos químicos fabricados con laminados impermeables, goma de butilo ≥ 14 mils; goma de nitrilo ≥ 14 mils; goma de neopreno ≥ 14 mils; goma natural ≥ 14 mils; polietileno, cloruro de polivinilo ≥ 14 mils, o Viton ≥ 14 mils, al aplicar con cualquier boquilla o equipo portátil, mezclar o cargar, limpiar derrames o equipos, o de cualquier otra forma estar expuesto al concentrado
 - Delantal resistente a productos químicos cuando se mezcle o se cargue, se limpien derrames o el equipo o cuando se pueda estar expuesto al producto concentrado
- Consulte **Controles de ingeniería** para conocer los requisitos adicionales.

Deseche la ropa y otros materiales absorbentes que hayan sido empapados o muy contaminados con el concentrado de este producto. No vuelva a reutilizarlos. Siga las instrucciones del fabricante para la limpieza/mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE separadamente de otra ropa para lavar.

DECLARACIÓN DE CONTROL TÉCNICO

Cuando los manipuladores de este producto utilizan sistemas cerrados o cabinas cerradas de conformidad con los requisitos estipulados en el Estándar para la Protección del Trabajador (WPS, según siglas en inglés) relativos a los pesticidas agrícolas [40 CFR 170.607 (d-e)], es posible reducir o modificar los requisitos relativos al equipo protector personal del manipulador, según lo especificado en el WPS.

Los pilotos deben utilizar una cabina que cumpla con los requisitos establecidos en la Ley de Protección al Trabajador (WPS, por sus siglas en inglés) para pesticidas agrícolas [40 CFR 170.607(d-f)].

RIESGOS AMBIENTALES

Este producto es tóxico hacia peces e invertebrados acuáticos. La deriva o la escorrentía pueden afectar negativamente a invertebrados acuáticos y las plantas no objetivo. No aplique directamente a cuerpos de agua. Tenga cuidado de minimizar el rociado accidental a lo largo de la costa cuando aplique a plantas terrestres en el borde del agua o al agua en áreas donde hay agua superficial. No aplicar directamente en áreas intermareales debajo de la marca media de pleamar, excepto según lo permitido en esta etiqueta. La deriva o el rociado y las escorrentías pueden ser peligrosas para los organismos acuáticos en aguas adyacentes a las áreas tratadas. No contamine el agua al desechar el agua de lavado o enjuague del equipo.

Advertencia Sobre Aguas Subterráneas

Se sabe que [2,4-D y Aminopirralid] se filtra a través del suelo hacia las aguas subterráneas en determinadas condiciones como resultado del uso agrícola. Estos químicos se pueden filtrar a aguas subterráneas si es utilizado en áreas donde el suelo es permeable, especialmente en donde la tabla de agua está cercana a la superficie. Los usuarios deben evitar la aplicación de estos químicos en suelos con una permeabilidad de rápida a muy rápida en todo el perfil (como de arena limosa a arena) y donde la capa freática de un acuífero subyacente es poco profunda, o en suelos que contienen sumideros sobre roca caliza, superficies severamente fracturadas y sustratos que permitirían la introducción directa en un acuífero. Las agencias agrícolas locales pueden proporcionar más información sobre el tipo de suelo en su área y la ubicación del agua subterránea.

Advertencia Referente a Las Aguas Superficiales

Este producto puede impactar la calidad de cuerpos de agua superficiales debido a escorrentías de aguas pluviales. Esto es especialmente notable en suelos con poco drenaje y suelos con una tabla de agua que está cercana a la superficie. Este producto está clasificado como de alto potencial para alcanzar agua superficial a través del escurrimiento por varios meses después su aplicación. Una franja de protección o amortiguamiento vegetativo, nivelada y bien mantenida entre las áreas donde este producto es utilizado y cuerpos de agua superficiales tales como lagos (lagunas), ríos y manantiales, reducirá el posible movimiento de Aminopirralid hacia aguas de escorrentía y sedimentos. Escorrentías de este producto pueden ser reducidas al evitar aplicaciones cuando estén pronosticados lluvia o irrigación dentro de 48 horas.

Declaración Consultiva Sobre Organismos No Objetivo

Este producto es tóxico para las plantas y puede afectar negativamente al forraje y al hábitat de organismos no objetivo, incluidos los polinizadores, en zonas adyacentes al lugar tratado. Proteja el forraje y el hábitat de los organismos no objetivo siguiendo las instrucciones de la etiqueta destinadas a minimizar la deriva de la pulverización.

Declaración de Agua de Riego

No contamine el agua destinada a riego o fines domésticos. Para evitar daños a los cultivos u otras plantas deseables, no trate ni permita que la deriva de la pulverización o la escorrentía del agua de lluvia caigan en las orillas o fondos de zanjas de riego, ya sea secas o con agua, u otros canales que transporten agua que pueda ser utilizada para riego o fines domésticos. No aplicar sobre nieve o suelo congelado.

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal.

ALMACENAMIENTO Y DESECHO

No contamine agua, comida ni forrajes mediante el almacenamiento y desecho.

Almacenamiento de pesticidas: Si este producto se expone a temperaturas bajo cero grados centígrados, el ingrediente activo puede cristalizarse y sedimentar. En estas condiciones, el producto debe calentarse a por lo menos 40°F y agitarse bien para disolver cualquier material cristalizado antes de su uso.

Desecho de pesticidas: Los residuos de pesticidas son tóxicos. La disposición inadecuada del pesticida sobrante, mezcla de aerosol o aguas de enjuague de contenedores, constituye una violación de la Ley Federal. Si estos residuos no pueden ser eliminados de acuerdo a las instrucciones de la etiqueta, contacte la agencia para el manejo de pesticidas o control ambiental de su estado, o el representante para el Manejo de Residuos Peligrosos de la Oficina Regional de la EPA más cercana para obtener indicaciones del proceso a seguir para su disposición.

Manipulación de envases:

Recipiente no rellenable. No reutilice ni rellene este recipiente. Ofrezca para el reciclaje si está disponible o perfore y deseche en un área adecuada para desechos sanitarios, o por otros procedimientos aprobados por las autoridades estatales y locales o la incineración.

Enjuague tres veces (o equivalente) el recipiente inmediatamente después de vaciarlo. Enjuague tres veces siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezclar, y drene por 10 segundos después de que el flujo empiece a gotear. Llene el recipiente 1/4 lleno con agua y tape de nuevo. Agite por 10 segundos. Eche el enjuague en el equipo de aplicación o tanque de mezclar, o almacene el enjuague para uso más tarde o eliminación. Drene por 10 segundos después de que el flujo empiece a gotear. Repita este procedimiento dos veces más. Enjuague a presión siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezclar, y drene por 10 segundos después de que el flujo empiece a gotear. Mantenga el recipiente boca abajo sobre el equipo de aplicación o un tanque de mezclar, o recolecte el enjuague para uso más tarde o la eliminación. Inserte la boquilla de enjuague a presión en el lado del recipiente y enjuague a aproximadamente 40 PSI por al menos 30 segundos. Drene por 10 segundos después de que el flujo empiece a gotear.

NOTES

NOTES

AMINOPYRALID	GROUP	4	HERBICIDE
2,4-D	GROUP	4	HERBICIDE

Amrita

SPECIALTY HERBICIDE

For control of susceptible weeds and certain woody plants, including invasive and noxious weeds, on rangeland, permanent grass pastures (including grasses grown for hay), Conservation Reserve Program (CRP) acres, non-cropland areas including industrial sites, rights-of-way (such as roadsides, electric utility and communication transmission lines, pipelines, and railroads) and non-irrigation ditch banks, natural areas (such as wildlife management areas including seasonally dry flood plains, deltas, marshes, prairie potholes, or vernal pools, wildlife openings, wildlife habitats, recreation areas, campgrounds, trailheads and trails), and grazed areas in and around these sites

Not For Sale, Distribution, or Use in New York State.

Not For Sale, Distribution, or Use in the San Luis Valley of Colorado.

Active Ingredients:	% w/w
Aminopyralid, Triisopropanolammonium salt:	
Triisopropanolammonium salt of 2-pyridine carboxylic acid,	
4-amino-3,6-dichloro	8.34%
2,4-D, dimethylamine salt:	
Dimethyl amine salt of (2,4-dichlorophenoxy) acetic acid.	41.96%
Other Ingredients	49.70%
Total	100.00%

Acid Equivalents:
 Aminopyralid (2-pyridine carboxylic acid, 4-amino-3,6-dichloro-) –
 4.3% - 0.41 lb./gal. (50 g/L)
 2,4-D [(2,4-dichlorophenoxy) acetic acid] –
 34.85% - 3.27 lbs./gal. (400 g/L)

KEEP OUT OF REACH OF CHILDREN DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the poison control center at 1-800-222-1222 for emergency medical treatment information.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS - HAZARD TO HUMANS AND DOMESTIC ANIMALS - DANGER - Corrosive. Causes Irreversible Eye Damage. Harmful if Swallowed. Do not get in eyes or on clothing.

Produced For: Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 83529-253 EPA Est. No. [AG] 72159-GA-001; [MC] 89332-GA-001; [SC] 39578-TX-001

The EPA Establishment Number is identified by the letters above that match the first two letters in the batch number.

Net Contents: **2.5 Gals.*** **265 Gals.**

* Unless alternate checked

Environmental Hazards - This product is toxic to fish and aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Take care to minimize the incidental overspray along the shoreline when applying to terrestrial plants at the water's edge or to water in areas where surface water is present. Do not apply directly to intertidal areas below the mean high-water mark, except as permitted on this label. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food, feed or fertilizer by storage or disposal.

Pesticide Storage: If this product is exposed to subfreezing temperatures, the active ingredient may crystallize and settle out of solution. Under these conditions the product should be warmed to at least 40°F and agitated well to dissolve any crystallized material prior to use.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Non-refillable containers 5 gallons or less: Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons: Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Nonrefillable containers 5 gallons or larger: Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product call 1-800-262-8200.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

OPEN HERE