

<b>TEBUCONAZOLE</b>	<b>GROUP</b>	<b>3</b>	<b>FUNGICIDE</b>
<b>TRIFLOXYSTROBIN</b>	<b>GROUP</b>	<b>11</b>	<b>FUNGICIDE</b>

# Entire

## FUNGICIDE

For control of certain diseases in the following crops: barley, corn, peanut, pecan, soybeans, wheat, and grasses grown for seed.

ACTIVE INGREDIENTS:	WT. BY %
Trifloxystrobin: Benzeneacetic acid, alpha-(methoxyimino)-2-[[[[[3-trifluoromethyl]phenyl]ethylidene]amino]methyl]-, methyl ester	22.63%
Tebuconazole: ( $\alpha$ -[2-(4-chlorophenyl)-ethyl]- $\alpha$ -(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol)	22.63%
<b>OTHER INGREDIENTS:</b>	<b>54.74%</b>
<b>TOTAL:</b>	<b>100.00%</b>


Contains 2.18 lbs. of trifloxystrobin and 2.18 lbs. of tebuconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN /  
MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS  
CAUTION / PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you **DO NOT** understand this label, find someone to explain it to you in detail.)

**See label booklet for complete First Aid, Precautionary Statements,  
Directions For Use, and Storage and Disposal.**

**Manufactured For:**

**Sharda USA LLC** 

7217 Lancaster Pike, Suite A  
Hockessin, Delaware 19707

EPA Reg. No. 83529-271

EPA Est. No. **AG** 72159-GA-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first five minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor immediately for treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at <b>1-800-222-1222</b> .	
<b>NOTE TO PHYSICIAN</b>	
No specific antidote. Treat symptomatically.	

## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

#### **CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

#### **Personal Protective Equipment (PPE)**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants,
- Shoes plus socks, and
- Chemical-resistant gloves such as: Barrier Laminate, Butyl Rubber  $\geq$  14 mils, Nitrile Rubber  $\geq$  14 mils, Neoprene Rubber  $\geq$  14 mils, Polyvinyl Chloride (PVC)  $\geq$  14 mils, Viton  $\geq$  14 mils.

#### **ENGINEERING CONTROL STATEMENTS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### **USER SAFETY RECOMMENDATIONS**

##### **Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

#### **Groundwater Advisory**

Several trifloxystrobin degradates have properties and characteristics associated with chemicals detected in groundwater. The use of trifloxystrobin and tebuconazole in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use.

### Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** use this product until you have read the entire label. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

**PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:**

- Long-sleeved shirt and long pants,
- Shoes plus socks, and
- Chemical-resistant gloves such as: Barrier Laminate, Butyl Rubber  $\geq$  14 mils, Nitrile Rubber  $\geq$  14 mils, Neoprene Rubber  $\geq$  14 mils, Polyvinyl Chloride (PVC)  $\geq$  14 mils, Viton  $\geq$  14 mils.

### PRODUCT INFORMATION

**Entire** is a broad spectrum fungicide for the control of certain diseases of barley, corn, peanut, pecan, soybean, wheat, and grasses grown for seed. **Entire** works by interfering with both energy and cell membrane production by plant pathogenic fungi.

UNDER CERTAIN CONDITIONS CONDUCIVE TO EXTENDED INFECTION PERIODS, ADDITIONAL FUNGICIDE APPLICATIONS BEYOND THE NUMBER ALLOWED BY THIS LABEL MAY BE NEEDED. UNDER THESE CONDITIONS, USE ANOTHER FUNGICIDE REGISTERED FOR THE CROP/DISEASE.

### RESISTANCE MANAGEMENT

The active ingredients in **Entire** belong to two different chemistry classes. Tebuconazole belongs to the DMI (Group 3) class of chemistry which exhibits no known cross resistance to other chemical classes. Tebuconazole may exhibit cross resistance to other Group 3 fungicides, such as propiconazole and myclobutanil. Trifloxystrobin belongs to the QoI (Group 11) class of chemistry which exhibits no known cross-resistance to other chemical classes. Trifloxystrobin does exhibit cross-resistance to other Group 11 fungicides, such as azoxystrobin, pyraclostrobin, and kresoxim-methyl. The NA-QoI Working Group has established the following general guidelines for the maximum number of applications of a Group 11-containing fungicide. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Entire** or other Group 3/Group 11 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your local Sharda USA representative at <https://shardausa.com/>.

### SPRAY EQUIPMENT

Thorough coverage is necessary to provide good disease control. Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control. For ground application equipment, a minimum of 10 gallons per acre is specified. For aerial application equipment, a minimum of 2 gallons per acre is specified.

## Broadcast Ground Sprayers

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate the sprayer before use.

Use a pump with the capacity to: (1) maintain a minimum of 35 psi at nozzles, and (2) provide sufficient agitation in the tank to keep the mixture in suspension - this requires recirculation of 10% of the tank volume per minute. Use jet agitators or a liquid sparge tube for vigorous agitation.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump must be 16-mesh or coarser. **DO NOT** place a screen in the recirculation line. Use 50-mesh screens at the nozzles. Check nozzle manufacturer's recommendations.

For information on spray equipment and calibration, consult sprayer manufacturer's and/or state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.

## Aerial Application

Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. **DO NOT** apply directly to humans or animals. Not registered for aerial application in New York State.

## Chemigation

Apply **Entire** through irrigation equipment only to crops for which chemigation is specified on this label.

**Entire** alone or in combination with other pesticides which are registered for application through irrigation systems, may be applied through irrigation systems. Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. **DO NOT** apply this product through any other type of irrigation system. Illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

## Operating Instructions

The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. **DO NOT** apply when wind speed favors drift beyond the area intended.

## Center Pivot Irrigation Equipment

### Restrictions:

- Use only with drive systems which provide uniform water distribution.
- **DO NOT** use end guns when chemigating **Entire** through center pivot systems because of non-uniform application.

Determine the size of the area to be treated. Determine the time required to apply 1/8 - 1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. When applying **Entire** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80 - 95% of the manufacturer's rated capacity. Using water, determine the injection pump output when operated at normal line pressure. Determine the amount of **Entire** required to treat the area covered by the irrigation system. Add the required amount of **Entire** and sufficient water to meet the injection time requirements to the solution tank. Make sure the system is fully charged with water before starting injection of the **Entire** solution. Time the injection to last at least as long as it takes to bring the system to full pressure. Maintain constant solution tank agitation during the injection period. Continue to operate the system until the **Entire** solution has cleared the sprinkler head.

## Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

When applying **Entire** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Determine the amount of **Entire** required to treat the area covered by the irrigation system. Add the required amount of **Entire** into the same quantity of water used to calibrate the injection period. Operate the system at the same pressure and time interval established during the calibration. Stop injection equipment after treatment is completed. Continue to operate the system until the **Entire** solution has cleared the last sprinkler head.

## MIXING PROCEDURES

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. **DO NOT** let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

**Entire Alone:** Add approximately 1/2 of the required amount of water to the mix tank. With the agitator running, add the **Entire** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the **Entire** has completely and uniformly dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

**Entire + Tank Mix Partners:** Add approximately 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank-mix partners. In general, tank-mix partners must be added in this order: products packaged in water soluble packaging\*, wettable powders, wettable granules (dry flowables), liquid flowables such as **Entire**, liquids, and emulsifiable concentrates. Always allow each tank-mix partner to become fully and uniformly dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

When using **Entire** in tank mixtures, all products in water-soluble packaging must be added to the tank before any other tank-mix partner, including **Entire**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix partner to the tank. If using **Entire** in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix partner product label. No label dosage rate must be exceeded, and the most restrictive label precautions and limitations must be followed. This product must not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

**Entire** is compatible with most insecticide, fungicide, and foliar nutrient products. However, the physical compatibility of **Entire** with tank-mix partners must be tested before use. To determine the physical compatibility of **Entire** with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

The crop safety of all potential tank mixes including additives and other pesticides on all crops has not been tested. Before applying any tank mixture not specifically recommended on this label, the safety to the target crop should be confirmed. To test for crop safety, apply **Entire** to the target crop in a small area and in accordance with label instructions for the target crop.

#### MANDATORY SPRAY DRIFT MANAGEMENT

##### Aerial Applications:

- For aerial applications, **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** release spray at a height greater than 10 feet above the ground or vegetative canopy unless a greater application height is necessary for pilot safety.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

##### Airblast Applications:

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

##### Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

#### SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

##### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

##### **Controlling Droplet Size - Ground Boom**

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

##### **Controlling Droplet Size - Aircraft**

**Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

**BOOM HEIGHT - Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

**RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift.

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

**TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

**WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

**Handheld Technology Applications**

- Take precautions to minimize spray drift.

---

**CROP SPECIFIC USE DIRECTIONS**


---

**BARLEY**

Diseases Controlled	Rate per Acre	Use Restrictions
Glume Blotch <i>(Stagonospora nodorum)</i> Leaf Blotch <i>(Stagonospora avenae)</i> Net Blotch <i>(Pyrenophora teres)</i> Powdery Mildew <i>(Blumeria graminis)</i> Rusts <i>(Puccinia spp.)</i> Scald <i>(Rhynchosporium secalis)</i> Spot Blotch <i>(Cochliobolus sativus)</i>	3.3 fl. oz.	<p><b>DO NOT</b> apply more than 3.3 fl. oz. of per season.</p> <p><b>DO NOT</b> apply within 40 days of harvest. For optimum disease control, the lowest labeled rate of a spray non-ionic surfactant (NIS) may be tank-mixed.</p> <p>For resistance management, <b>DO NOT</b> apply more than 2 consecutive applications of a Group 11 or Group 11-containing fungicide per acre per year without alternation with at least 2 applications of fungicide from a different (not Group 11) mode of action.</p> <p><b>DO NOT</b> allow livestock to graze within the treated area within 30 days after application, and <b>DO NOT</b> harvest the treated crop for forage within 30 days after application or for hay within 45 days after application.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>
<p><b>Use Directions:</b></p> <ul style="list-style-type: none"> <li>• May be applied by ground, aerial or chemigation.</li> <li>• Begin applications preventatively when conditions are favorable for disease development.</li> </ul>		

**SWEET CORN (AND SEED PRODUCTION)**

Diseases Controlled	Rate per Acre	Use Restrictions
Anthracnose Leaf Blight <i>(Colletotrichum graminicola)</i> Common Rust <i>(Puccinia sorghi)</i> Eye Spot <i>(Aureobasidium zeae)</i> Gray Leaf Spot <i>(Cercospora zeae-maydis)</i> Northern Corn Leaf Blight <i>(Setopshaeria turcica)</i> Northern Corn Leaf Spot <i>(Cochliobolus carbonum)</i> Southern Corn Leaf Blight <i>(Cochliobolus heterostrophus)</i> Southern Rust <i>(Puccinia polysora)</i>	5.0 - 6.0 fl. oz.	<p><b>DO NOT</b> apply more than 24 fl. oz. per acre per use season.</p> <p><b>DO NOT</b> apply more than 4 applications per use season.</p> <p>May be applied up to 7 days before the harvest of ears and forage.</p> <p><b>DO NOT</b> apply within 49 days of harvest for fodder.</p> <p>In programs with Group 11 tank mixes, or other pre-mixes containing a Group 11 fungicide, the number of Group 11 fungicide must be no more than one-half of the total number of fungicide applications per season. Alternate every application of with at least one application of a non-Group 11 fungicide.</p> <p>Restricted-entry interval (REI) = 19 days.</p>
<p><b>Use Directions:</b></p> <ul style="list-style-type: none"> <li>• May be applied by ground, air or chemigation.</li> <li>• Should be applied in a minimum of 10 gal./A of spray solution by ground sprayer or in a minimum of 2 gal./A by aircraft spray equipment. For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed.</li> <li>• Apply when disease first appears and continue on a 10 - 14-day interval if favorable conditions for disease development persist. Use of shorter spray intervals and higher rates are specified when disease pressure is severe.</li> </ul>		

**CORN (FIELD CORN, FIELD CORN GROWN FOR SEED, AND POPCORN)**

Diseases Controlled	Rate per Acre	Use Restrictions
Anthracnose Leaf Blight <i>(Colletotrichum graminicola)</i> Common Rust <i>(Puccinia sorghi)</i> Eye Spot <i>(Aureobasidium zeae)</i> Gray Leaf Spot <i>(Cercospora zeae-maydis)</i> Northern Corn Leaf Blight <i>(Setopshaeria turcica)</i> Northern Corn Leaf Spot <i>(Cochliobolus carbonum)</i> Southern Corn Leaf Blight <i>(Cochliobolus heterostrophus)</i> Southern Rust <i>(Puccinia polysora)</i>	5.0 - 6.0 fl. oz.	<p><b>DO NOT</b> apply more than 12 fl. oz. per acre per use season.</p> <p><b>DO NOT</b> apply more than 2 applications per use season.</p> <p>May be applied up to 36 days before the harvest of grain and fodder.</p> <p><b>DO NOT</b> apply within 21 days of harvest for forage.</p> <p><b>DO NOT</b> apply more than two sequential applications.</p> <p>Limit the number of Group 11-containing fungicide applications to no more than two per acre per crop.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>
<p><b>Use Directions:</b></p> <ul style="list-style-type: none"> <li>• May be applied by ground, air or chemigation.</li> <li>• Must be applied in a minimum of 10 gal./A of spray solution by ground sprayer or in a minimum of 2 gal./A by aircraft spray equipment. For optimum disease control, the lowest labeled rate of a spray surfactant may be tank-mixed.</li> <li>• Apply when disease first appears and continue on a 10 - 14-day interval if favorable conditions for disease development persist. Use of shorter spray intervals and higher rates are specified when disease pressure is severe.</li> </ul>		

**GRASSES GROWN FOR SEED (NORTHWEST U.S. ONLY)**

Diseases Controlled	Rate per Acre	Use Restrictions
Rust ( <i>Puccinia</i> spp.) Powdery Mildew ( <i>Erysiphe graminis</i> )	5 - 7.7 fl. oz.	<b>DO NOT</b> apply more than 32 fl. oz. of per acre per year. <b>DO NOT</b> apply more than 2 sequential applications, or other Group 11 containing fungicide without alternation to at least 2 applications of a fungicide from a different (not Group 11) mode of action. For optimum performance, the lowest specified rate of a spray surfactant containing methylated seed oil, or other equivalent oil-based product, should be tank mixed. <b>DO NOT</b> apply within 4 days of harvest. <b>DO NOT</b> forage or cut green crop for feed purposes. Chaff, screenings, and straw from treated areas may be used for feed purposes, but <b>DO NOT</b> use seed for feed purposes. Regrowth may be grazed starting 17 days after the last application. Restricted-entry interval (REI) = 12 hours.
<b>Use Directions:</b> <ul style="list-style-type: none"> <li>Begin applications when rust and powdery mildew infections are noticeable and beginning to increase in number. Continue applications on a 21-day application interval.</li> <li>Continue applications if favorable conditions for disease development persist. Use higher rates when disease pressure is severe.</li> <li>Most bluegrass has little resistance to rust or powdery mildew. It is important to begin applications early in the growing season for bluegrass and other more susceptible species.</li> <li>Apply in a minimum of 20 gal./A for ground application, or in a minimum of 10 gal./A for aerial application.</li> </ul>		

**PEANUTS**

Diseases Controlled	Rate per Acre	Use Restrictions
Early Leaf Spot ( <i>Cercospora arachidicola</i> ) Late Leaf Spot ( <i>Cercosporidium personatum</i> ) Rust ( <i>Puccinia arachidis</i> ) Web Blotch ( <i>Phoma arachidicola</i> )	3.5 fl. oz.	<b>DO NOT</b> exceed more than 4 total applications per season. <b>DO NOT</b> apply within 14 days of harvest. If 4 or less total fungicide sprays are planned, alternate each application with a non-Group 11 containing fungicide. If 5 or more fungicide sprays are planned, use a maximum of 2 consecutive applications alternated with at least 2 applications of a non-Group 11 containing fungicide before returning to another Group 11 fungicide. To limit development of disease resistance, <b>DO NOT</b> apply a Group 11 containing fungicide for more than 1/2 of the seasonal sprays.
White Mold ( <i>Sclerotium rolfsii</i> )	3.5 fl. oz. + 5.2 fl. oz. Folicur® 3.6 F	<b>DO NOT</b> feed hay or threshings or allow livestock to graze in treated areas. Restricted-entry interval (REI) = 12 hours.
Limb Rot ( <i>Rhizoctonia solani</i> )	7.0 fl. oz.	
<b>Use Directions:</b> <ul style="list-style-type: none"> <li>Begin applications when conditions are favorable for diseases but before infection. Apply on a 10 - 14-day spray schedule. Use shorter intervals when disease pressure is severe.</li> <li>May be applied by ground, aerial, or chemigation.</li> <li>For White Mold: Folicur® tank-mix - Begin applications when conditions are favorable for diseases, typically within timings 3 - 6 in a seven-spray program. This Folicur® tank-mix, when part of a Folicur® four block program, will also provide protection against Rhizoctonia limb rot. May be applied by ground, aerial, or chemigation.</li> <li>For Limb Rot: Begin applications when conditions are favorable for diseases, typically within timings 3 - 6 in a seven-spray program. In the southeast, applications at approximately 90 and 104 days after planting may be the most effective timings for control of limb rot.</li> <li>May be applied by ground, aerial, or chemigation.</li> </ul>		

## PECAN

Diseases Controlled	Rate per Acre	Use Restrictions
Pecan Scab ( <i>Cladosporium caryigenum</i> ) Anthracnose ( <i>Glomerella cingulata</i> )	5 - 7.67 fl. oz.	<b>DO NOT</b> make more than 6 applications per season. <b>DO NOT</b> apply more than 46 oz. per acre per season. <b>DO NOT</b> cut cover crops in treated areas for feed or allow livestock to graze treated areas. To limit the potential for resistance to develop apply up to 2 consecutive applications, then make at least 2 applications with an effective fungicide with a different mode of action (a non-Group 11) before returning to <b>Entire</b> . To limit development of disease resistance, <b>DO NOT</b> apply a Group 11 containing fungicide for more than 1/2 of the seasonal sprays. Restricted-entry interval (REI) = 12 hours.
<b>Use Directions:</b> <ul style="list-style-type: none"> <li>• Begin applications when conditions are favorable for disease development and continue throughout the season using a 14 - 21-day interval.</li> <li>• Will control scab on both the leaf and shuck. <b>DO NOT</b> apply after shuck split or within 30 days of harvest.</li> <li>• A surfactant may be added to the spray solution for optimum control of the indicated diseases.</li> <li>• May be applied by ground, aerial, or chemigation.</li> </ul>		

## SOYBEAN

Diseases Controlled	Rate per Acre	Use Restrictions
Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum truncatum</i> ) Asian Soybean Rust ( <i>Phakopsora pachyrhizi</i> ) Brown spot ( <i>Septoria glycinis</i> ) Cercospora Blight ( <i>Cercospora kikuchii</i> ) Frogeye Leaf Spot ( <i>Cercospora sojina</i> ) Pod and Stem Blight ( <i>Diaporthe phaseolorum</i> ) Powdery Mildew ( <i>Microspheera diffusa</i> ) Rhizoctonia Aerial Blight ( <i>Rhizoctonia solani</i> )	5.0 - 6.0 fl. oz.	Applications may not be made within 21 days of harvest. <b>DO NOT</b> apply more than 3 applications per season (i.e. 18 fl. oz. of product which is equivalent to 0.3 lb. of active ingredient). <b>DO NOT</b> apply more than two sequential applications. For optimum disease control, the lowest labeled rate of a spray non-ionic surfactant (NIS) may be tank-mixed. Limit the number of Group 11-containing fungicide applications to no more than two per acre per crop. <b>DO NOT</b> graze or feed soybean forage or hay. Restricted-entry interval (REI) = 12 hours.
<b>Use Directions:</b> <ul style="list-style-type: none"> <li>• For diseases other than Asian soybean rust, for best disease control, apply at early flowering or prior to disease development.</li> <li>• For Asian soybean rust control, apply as a preventative spray or at first visible symptoms of disease.</li> <li>• Repeat applications on a 10 - 21-day spray interval if environmental conditions are favorable for continued disease development. Use of shorter spray intervals and higher rates are recommended when disease pressure is severe.</li> </ul>		

## WHEAT

Diseases Controlled	Rate per Acre	Use Restrictions
Glume Blotch <i>(Stagonospora nodorum)</i> Leaf Blight <i>(Septoria tritici)</i> Powdery Mildew <i>(Blumeria graminis f. sp. tritici)</i> Rusts <i>(Puccinia spp.)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	5.0 fl. oz.	<p><b>DO NOT</b> apply more than 5 fl. oz. of per season.</p> <p><b>DO NOT</b> apply within 35 days of harvest. For optimum disease control, the lowest labeled rate of a spray non-ionic surfactant (NIS) may be tank-mixed.</p> <p>For resistance management, <b>DO NOT</b> apply more than 2 consecutive applications of <b>Entire</b> or other Group 11 or Group 11-containing fungicide per acre per year without alternation with at least 2 applications of fungicide from a different (non-Group 11) mode of action.</p> <p><b>DO NOT</b> allow livestock to graze within the treated area within 30 days after application, and <b>DO NOT</b> harvest the treated crop for forage within 30 days after application or for hay and wheat straw within 45 days after application.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>
<p><b>Use Directions:</b></p> <ul style="list-style-type: none"> <li>Begin applications preventatively when conditions are favorable for disease development.</li> <li>Early season leaf disease suppression: apply 3 - 4 fl. oz. per acre for suppression of Tan Spot, Leaf Blight, and Powdery Mildew.</li> <li>May be applied by ground, aerial or chemigation.</li> </ul>		

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store this product in a cool, dry place in its original container only. **DO NOT** store this product near fertilizers, seeds, or other pesticides. If this product is spilled, sweep up the spillage and dispose pursuant to the below **PESTICIDE DISPOSAL** instructions.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

**Less Than or Equal to 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

**Greater Than 5 Gallons:** Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**For Bulk and Mini-Bulk Containers:** Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

<b>PRIMEROS AUXILIOS</b>	
<b>SI SE INGIERE:</b>	<ul style="list-style-type: none"> <li>• Llame inmediatamente a un centro de control de envenenamientos o a un médico para consejo de tratamiento.</li> <li>• Si la persona puede tragar, haga que tome pequeños sorbos de agua.</li> <li>• <b>NO</b> induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico.</li> <li>• <b>NO</b> administre nada por boca a una persona que haya perdido el conocimiento.</li> </ul>
<b>SI CAE EN LA PIEL O LA ROPA:</b>	<ul style="list-style-type: none"> <li>• Qúitese la ropa contaminada.</li> <li>• Enjuague la piel inmediatamente con bastante agua por 15 - 20 minutos.</li> <li>• Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.</li> </ul>
<b>SI ES INHALADO:</b>	<ul style="list-style-type: none"> <li>• Traslade a la persona al aire fresco.</li> <li>• Si la persona no está respirando llame al 911 o a una ambulancia, luego dé respiración artificial, preferiblemente de boca a boca, si es posible.</li> <li>• Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.</li> </ul>
<b>SI ENTRA EN CONTACTO CON LOS OJOS:</b>	<ul style="list-style-type: none"> <li>• Mantenga los ojos abiertos y enjuáguelos lenta y cuidadosamente con agua, durante 15 a 20 minutos.</li> <li>• Si utiliza lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos.</li> <li>• Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.</li> </ul>
<b>NÚMERO DE TELÉFONO DIRECTO</b>	
<p>Cuando llame a un centro de control de envenenamiento, o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. Para información de emergencia sobre este producto, llame al centro de control de envenenamientos a <b>1-800-222-1222</b>.</p>	
<b>NOTA AL MÉDICO</b>	
<p>No hay antídoto específico disponible. Tratar sintomáticamente.</p>	

## DECLARACIONES PREVENTIVAS

### RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS

#### PRECAUCIÓN

Dañino si se ingiere o inhala o es absorbido por la piel. Causa irritación moderada en los ojos. Evite el contacto con la piel, los ojos o la ropa. Evite respirar el vapor o la neblina de aspersión. Lávese bien con agua y jabón después de manipular y antes de comer, beber, masticar chicle, o usar tabaco. Qúitese y lave la ropa contaminada antes de reutilizarla.

#### **Equipo de protección personal (PPE, por sus siglas en inglés)**

Siga las instrucciones del fabricante para la limpieza y el mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE por separado del resto de la ropa.

#### **Los aplicadores y otros manipuladores deberán usar:**

- Camisa de manga larga y pantalones largos,
- Zapatos y calcetines, y
- Guantes resistentes a productos químicos, por ejemplo: laminados impermeables, goma de butilo  $\geq$  14 mils, goma de nitrilo  $\geq$  14 mils, goma de neopreno  $\geq$  14 mils, cloruro de polivinilo (PVC)  $\geq$  14 mils, Viton  $\geq$  14 mils.

#### **DECLARACIÓN DE CONTROL TÉCNICO**

Quando los manipuladores de este producto utilizan sistemas cerrados de manera que cumplan con los requisitos estipulados en el Estándar para la Protección de los Trabajadores (WPS, según siglas en inglés) relativos a los pesticidas agrícolas [40 CFR 170.240 (d)(4-6)], es posible reducir o modificar los requisitos relativos al equipo protector personal del manipulador, según lo especificado en el WPS.

#### **RIESGOS AMBIENTALES**

Este pesticida es tóxico para los mamíferos, los peces y los invertebrados acuáticos. **NO** aplique el producto directamente al agua, a zonas donde haya presencia de agua superficial ni a las zonas intermareales por debajo de la marca de pleamar media. La deriva y la escorrentía representan un peligro para los organismos acuáticos en las aguas adyacentes a las zonas tratadas. **NO** contamine el agua cuando disponga de las aguas de lavado de equipo o las aguas.

#### **Advertencia sobre aguas subterráneas**

Varios productos de degradación de trifloxystrobin presentan propiedades y características asociadas a sustancias químicas detectadas en las aguas subterráneas. El uso de trifloxystrobin y tebuconazole en zonas con suelos permeables, especialmente donde el nivel freático es poco profundo, puede provocar la contaminación de las aguas subterráneas. Se sabe que Tebuconazole se filtran a través del suelo hacia las aguas subterráneas en determinadas condiciones como resultado de su uso indicado en la etiqueta.

#### **Advertencia referente a las aguas superficiales**

Este producto puede contaminar el agua debido a la deriva de la aspersión por la acción del viento. Este producto tiene potencial de escorrentía durante varios meses o más después de la aplicación. Los suelos con drenaje deficiente y los suelos con niveles freáticos poco profundos son más propensos a generar escorrentía que contiene

este producto. Una franja de protección o amortiguamiento vegetativo, nivelada y bien mantenida entre las áreas donde este producto es utilizado y cuerpos de agua superficiales tales como estanques, arroyos y manantiales, reducirá el potencial de contaminación del agua por escorrentía de lluvia. La escorrentía de este producto puede reducirse al evitar aplicaciones cuando se pronostique lluvia dentro de 48 horas.

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal.

## ALMACENAMIENTO Y DESECHO

**NO** contamine el agua, los alimentos ni los forrajes durante el almacenamiento o el desecho.

**ALMACENAMIENTO DE PESTICIDAS:** Almacene este producto en un lugar fresco y seco, únicamente en su envase original. **NO** lo guarde cerca de fertilizantes, semillas ni otros pesticidas. En caso de derrame, recoja el derrame y deséchelo siguiendo las instrucciones de eliminación de pesticidas que se indican a continuación.

**DESECHO DE PESTICIDAS:** Los residuos de pesticidas son tóxicos. La disposición inadecuada del pesticida sobrante, mezcla de pulverización, o aguas de enjuague, constituye una violación de la Ley Federal. Si estos residuos no pueden ser eliminados de acuerdo con las instrucciones de la etiqueta, contacte la agencia estatal de pesticidas o de control ambiental de su estado, o el representante para el Manejo de Residuos Peligrosos de la Oficina Regional de la EPA más cercana para obtener instrucciones sobre métodos adecuados de eliminación.

### MANIPULACIÓN DE ENVASES:

**Menos o igual a 5 galones:** Envase no rellenable. **NO** reutilice ni rellene este envase. Ofrézcalo para reciclaje si está disponible. Enjuague tres veces el envase (o equivalente) inmediatamente después de vaciarlo. Enjuague tres veces siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezcla, y drene por 10 segundos después de que el flujo empiece a gotear. Llene el envase hasta 1/4 de su capacidad con agua y tape de nuevo. Agite por 10 segundos. Vierta las aguas de enjuague en el equipo de aplicación o tanque de mezcla, o almacene el enjuague para uso posterior o eliminación. Drene por 10 segundos después de que el flujo empiece a gotear. Repita este procedimiento dos veces más. Luego ofrézcalo para reciclaje si está disponible o reacondiciónelo si es apropiado, o perfore y deseche en un relleno sanitario o mediante incineración.

**Más de cinco galones:** Envase rellenable. Rellene este envase sólo con pesticida. **NO** reutilice este envase para ningún otro propósito. El triple enjuague de este envase es responsabilidad de la persona que lo desecha. La limpieza antes de rellenarlo es responsabilidad de la persona que lo rellena. Para limpiar el envase antes de su eliminación final, vacíe el contenido restante de este envase en el equipo de aplicación o en el tanque de mezcla. Llene el envase con un 10% de agua. Agite vigorosamente o haga recircular el agua con la bomba durante 2 minutos. Vierta o bombee las aguas de enjuague en el equipo de aplicación o en el sistema de recolección de aguas de enjuague. Repita este procedimiento dos veces más.

**Para contenedores y mini-contenedores a granel:** Envase rellenable. Rellene este envase sólo con pesticida. **NO** reutilice este envase para ningún otro propósito. La limpieza del envase antes de su eliminación final es la responsabilidad de la persona que se deshace del envase. La limpieza antes de rellenarlo es responsabilidad de la persona que lo rellena. Para limpiar el envase antes de su eliminación final, vacíe el contenido restante de este envase en el equipo de aplicación o en el tanque de mezcla. Llene el envase con un 10% de agua. Agite vigorosamente o haga recircular el agua con la bomba durante 2 minutos. Vierta o bombee las aguas de enjuague en el equipo de aplicación o en el sistema de recolección de aguas de enjuague. Repita este procedimiento dos veces más. Luego ofrézcalo para reciclaje si está disponible o perfore y deseche en un relleno sanitario, o por otros procedimientos aprobados por las autoridades estatales y locales.

**¡EL ENVASE NO ES SEGURO PARA ALIMENTOS, FORRAJES, NI AGUA POTABLE!**

**NOTES**

**NOTES**

**NOTES**

TEBUCONAZOLE	GROUP	3	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE

# Entire

For control of certain diseases in the following crops: barley, corn, peanut, pecan, soybeans, wheat, and grasses grown for seed.

ACTIVE INGREDIENTS:	WT. BY %
Trifloxystrobin: Benzenecetic acid, alpha-(methoxyimino)-2-[[[3-(trifluoromethyl)phenyl]ethylidene]amino]methyl]-, methyl ester . . . . .	22.63%
Tebuconazole: (α-[2-(4-chlorophenyl)-ethyl]-cc-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol) . . . . .	22.63%
<b>OTHER INGREDIENTS:</b> . . . . .	54.74%
<b>TOTAL:</b> . . . . .	100.00%

Contains 2.18 lbs. of trifloxystrobin and 2.18 lbs. of tebuconazole per gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

**FIRST AID - IF SWALLOWED:** • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. • **DO NOT** give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice. **IF INHALED:** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice. **IF IN EYES:** • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first five minutes, then continue rinsing. • Call a poison control center or doctor immediately for treatment advice.

**HOTLINE NUMBER** - Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222.

**NOTE TO PHYSICIAN** - No specific antidote. Treat symptomatically.

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

**PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION** - Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**Manufactured For:** Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 83529-271 EPA Est. No. **AG** 72159-GA-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

**ENVIRONMENTAL HAZARDS** - This pesticide is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. **Groundwater Advisory:** Several trifloxystrobin degradates have properties and characteristics associated with chemicals detected in groundwater. The use of trifloxystrobin and tebuconazole in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. **Surface Water Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

**DIRECTIONS FOR USE** - It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** use this product until you have read the entire label. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage and disposal. **PESTICIDE STORAGE:** Store this product in a cool, dry place in its original container only. **DO NOT** store this product near fertilizers, seeds, or other pesticides. If this product is spilled, sweep up the spillage and dispose pursuant to the below **PESTICIDE DISPOSAL** instructions. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER HANDLING: Less Than or Equal to 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration. **Greater Than 5 Gallons:** Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times. **For Bulk and Mini-Bulk Containers:** Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities. **CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

Net Contents: **2.5 Gals.\***  **265 Gals.**

\* Unless alternate checked