

AXYL SHIELD ST

FOR USE AS A SEED TREATMENT FOR CONTROL OF SYSTEMIC DOWNY MILDEW, PYTHIUM SEED ROT, PYTHIUM DAMPING-OFF, AND EARLY SEASON PHYTOPHTHORA DISEASES OF CERTAIN CROPS.

ACTIVE INGREDIENT:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester 30.00%

OTHER INGREDIENTS: 70.00%

TOTAL: **100.00%**

Contains 2.7 pounds metalaxyl per gallon (323 grams per liter).

EPA REG. NO. 83529-36

EPA EST. NO. 70815-GA-001

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for complete First Aid, Precautionary Statements, Directions for Use, and Limitation of Warranty and Liability.

Manufactured for:

Sharda USA LLC 

7217 Lancaster Pike Suite A
Hockessin, DE 19707

NET CONTENTS 2.5 GAL. (9.46 L)

FIRST AID

If Swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by to an unconscious person.
If in Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-hour medical emergency assistance (human or animal) call **1-800-222-1222**. For chemical emergency assistance (spill, leak, fire, or accident) call ChemTrec at **1-800-424-9300**.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks
- Chemical-resistant gloves such as barrier laminate or nitrile rubber or neoprene rubber or viton

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory Statement: This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves such as barrier laminate or nitrile rubber or neoprene rubber or viton

USE DIRECTIONS

This product is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora* spp. For control of other soil borne diseases, combination of captan, thiram and carboxin registered formulations are compatible. **DO NOT** use with other seed treatment products unless previous experience assures compatibility.

Note: This product is a systemic fungicide having a specific mode of action. This product could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible metalaxyl resistant strains of fungi which may occur. FAILURE TO FOLLOW DIRECTIONS OF THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

In California: Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

On-Farm Seed Treatment: **DO NOT** apply this product through any type of irrigation system.

DO NOT use treated seed for food or feed purposes or process for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. **DO NOT** store excess treated seeds beyond planting time.

Seed treatment on agricultural establishments at or immediately before planting is within the scope of WPS, while commercial treatment of seeds is not within the scope of WPS.

Planter Box Treatment: Apply at the specified rate and premix with the seed directly in the planter box at planting. Place half the amount of seed into the planter box and apply half the required amount of pesticide to the seed. Stir the seed thoroughly with a paddle until the seed is uniformly coated. Place the remaining half of the seed into the planter box and add the remaining half of the pesticide. Again stir the seed with the paddle until all seed is thoroughly coated.

For Commercial Seed Treatment: Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before sale.

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with metalaxyl fungicide.
- **DO NOT** use for feed, food, or oil purposes.
- User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.
- **DO NOT** use rape greens grown and harvested from Axl-Shield ST treated seed for human and feed consumption.
- Rapeseed grown and harvested from Axl-Shield ST treated seed is only for industrial uses and cannot be used for edible oil or any other human/ feed consumption.
- **DO NOT** use metalaxyl or mefenoxam for control of *Pythium* in water-seeded rice.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seed treated with Axl-Shield ST:

- Store away from food and feedstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, and waterproof gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwater.
- Dispose of seed packaging in accordance with local requirements.
- Excess treated seed may be used for ethanol production only if (1) byproducts are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used for agronomic practice.

Tank Mixing: When tank mixing follow the most restrictive label instructions for any tank mix partner. Do not tank mix with any product that contains a prohibition against tank mixing.

SEED TREATMENT

Apply this product as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

DOMESTIC USE

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Brassica Leafy Vegetables Broccoli (including Chinese and raab), Brussels sprout, Cabbage (including Chinese bok choy, Chinese napa, and mustard), Cauliflower, Cavalo broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, and Rape greens	<i>Pythium</i> damping-off control	0.75
Canola* Mustard Seed* Rapeseed*	<i>Pythium</i> damping-off control NOTE: Do not harvest mustard greens. Do not use rape greens grown and harvested from Axyl-Shield ST treated seed for human and feed consumption. Rapeseed grown and harvested from Axyl-Shield ST treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed.	0.25 to 0.50
Cotton	<i>Pythium</i> seed rot and damping-off control	0.75 to 1.50
Cucurbit Vegetables Chayote (fruit), Chinese wax gourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd (edible)(includes Hyotan, Cucuzza, Hechima, and Chinese okra), <i>Momordica</i> spp. (includes Balsam apple, Balsam pear, Bitter melon, and Chinese cucumber), Muskmelon (including hybrids and/or varieties of <i>Cucumis melo</i> including True cantaloupe, Cantaloupe, Casaba, Santa Claus melon, Crenshaw melon, Honeydew melon, Honey balls, Persian melon, Golden pershaw melon, Mango melon, Pineapple melon, and Snake melon), Watermelon (including hybrids and/or varieties of <i>Citrullus</i> spp.), Pumpkin, and Squash (summer, winter including Butternut squash, Calabaza, Hubbard squash, Acorn squash, and Spaghetti squash)	<i>Pythium</i> damping-off control	0.75
Forage Grasses Grasses grown for hay, grazing or silage, Corn (fodder or silage), Sorghum (hay or silage), and Small grain grown for hay, grazing or silage	<i>Pythium</i> damping-off control	0.75
Forage Legumes Alfalfa, Beans (for forage), Clover, Cowpeas (for forage), Cowpea hay, Lespedeza, Peanuts (for forage), Peanut hay, Peas (for forage), Pea vine hay, Soybeans (for forage), Soybean hay, Trefoil, Velvet beans (for forage), and Vetch	<i>Pythium</i> damping-off and early season <i>Phytophthora</i> control	0.75 to 1.50
Fruiting Vegetables Eggplant, Ground cherry (<i>Physalis</i> spp.), Pepino, Pepper (includes Bell pepper, Chili pepper, Cooking pepper, Pimento, and Sweet pepper), Tomatillo, and Tomato	<i>Pythium</i> damping-off control	0.75

* Not registered for use in California

(continued)

Domestic Use (continued)

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Grain Crops Barley, Buckwheat, Corn (Field corn, Sweet corn, and Popcorn), Millet, Milo, Oats, Rye, Triticale, and Wheat	<i>Pythium</i> damping-off control	0.75
	Systemic downy mildew control of sweet corn	1.50
Use Rate Conversions for Field Corn		
PRODUCT FLUID OUNCES /		
100 lbs. seed*	1,000 seeds	80,000 seed count
0.75	0.0042	0.334
1.50	0.0084	0.672

This product contains 9,404.5 milligrams metalaxyl (active ingredient) per fluid ounce.

*Based on standard number of 1,786 seeds per pound (100,000 seeds per 56.0 pound bushel).

NOTE: To maintain milligrams active ingredient per seed, adjust fluid ounces per 100 pounds in proportion to the actual seeds per pound compared to the standard number of seeds per pound for that crop.

For example, if there are 1,600 corn seeds per pound and the targeted rate is 0.040 milligram active ingredient per seed at a non-adjusted rate of 0.75 fluid ounce per 100 pounds of seed, use: $(1,600 \div 1,786) \times 0.75 = 0.67$ fluid ounce per 100 pounds of seed.

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Leafy Vegetables Amaranth (Chinese spinach), Cardoon, Celery, Celery (Chinese), Celeriac, Celtuce, Chervil, Chrysanthemum (edible-leaved, garland), Corn salad, Cress (garden, upland), Dandelion, Dock (sorrel), Endive (escarole), Fennel (Florence), Lettuce (head and leaf), Orach, Parsley, Purslane (garden, winter), Radicchio (Red chicory), Rhubarb, Spinach (including New Zealand and vine), and Swiss chard	<i>Pythium</i> damping-off control	0.75
	NOTE: Do not apply this product to spinach which is destined to be grown in the greenhouse. Containers or bags containing Axyl-Shield ST treated spinach seed must be labeled with this restriction.	
Onions (dry bulb and green)	<i>Pythium</i> damping-off control	0.75
Peanuts*	<i>Pythium</i> seed rot and damping-off control	0.75
Rice*	<i>Pythium</i> damping-off control	0.75 to 1.50
	Note: If using Axyl-Shield ST treated rice do not use metalaxyl or mefenoxam for control of <i>Pythium</i> in water-seeded rice. Containers or bags containing Axyl-Shield ST treated rice seed must be labeled with this restriction.	
Root and Tuber Vegetables (except potato) Arracacha, Arrowroot, Artichoke (Chinese, Jerusalem), Beet (garden, sugar), Burdock (edible), Canna (edible), Carrot, Cassava (bitter and sweet), Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Parsley (turnip-rooted), Parsnip, Radish (including Oriental), Rutabaga, Salsify (including black and Spanish), Skirret, Sweet potato, Tanier, Turmeric, Turnip, Yam bean, and Yam (true)	<i>Pythium</i> damping-off control	0.75
Seed and Pod Vegetables Black-eyed peas, Chickpea (Garbanzos), Cowpeas, Edible soybeans, Field beans, Field peas, Garden peas, Green beans, Kidney beans, Lentils, Lima beans, Lupines, Navy beans, Okra, Peas, Pole beans, Snap beans, String beans, and Wax beans	<i>Pythium</i> damping-off control	0.75
	Early season <i>Phytophthora</i> control	0.75
	Control of <i>Pythium</i> damping-off and systemic downy mildew of peas	1.50
Sorghum	<i>Pythium</i> damping-off control	0.375 to 0.75
	Systemic downy mildew (races susceptible to chemical control)	0.75 to 3.00
	Note: Apply 0.75 to 1.50 fl. oz. of this product on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.50 to 3.00 fl. oz. of this product. Only high quality sorghum seed (90% germination and above) should be treated with this product.	

*Not registered for use in California

Domestic Use (continued)

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)	
Soybeans	<i>Pythium</i> damping-off control and early season <i>Phytophthora</i> control	0.75 to 1.50	
Use Rate Conversions for Soybeans			
PRODUCT FLUID OUNCES /			MG A.I. / SEED
100 lbs. seed*	1,000 seeds	80,000 seed count	
0.75	0.0025	0.20	0.0235
1.50	0.005	0.40	0.047

This product contains 9,404.5 milligrams metalaxyl (active ingredient) per fluid ounce.

*Based on standard number of 3,000 seeds per pound (150,000 seeds per 50.0 pound bushel).

NOTE: To maintain milligram active ingredient per seed, adjust fluid ounces per 100 pounds in proportion to the actual seeds per pound compared to the standard number of seeds per pound for that crop. For example, if there are 2,800 soybean seeds per pound and the targeted rate is 0.0235 milligram active ingredient per seed at a non-adjusted rate of 0.75 fluid ounces per 100 pounds of seed, use: $(2,800 \div 3,000) \times 0.75 = 0.70$ fluid ounce per 100 pounds of seed.

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Sunflowers	Systemic downy mildew control	1.50 to 3.00
Turf Grasses Golf courses, Home lawns, Reclamation and Erosion control projects	<i>Pythium</i> damping-off control	1.50

DIRECTIONS FOR USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES

This product is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. This product may be applied as a water-based slurry through standard slurry of mist-type commercial seed treatment equipment.

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Barley, Corn (field), Oats, Rye, and Wheat	Aid in control of seed decay and damping-off caused by <i>Pythium</i>	0.10 to 0.375
Beans (field, kidney, navy and pinto), Lentils, Peas, and Soybeans		
Peanuts*		
Rice*		
Sunflowers		

* Not registered for use in California

Note: Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.

FOR EXPORT USE ONLY

For treatment of the following seed for export to countries requiring treatment with metalaxyl before importation.

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Brassica (Cole) leafy vegetables	<i>Pythium</i> damping-off control	0.75 to 1.50
Canola	<i>Pythium</i> damping-off control	0.50 to 1.70 (31 to 100 ml per 100 kg of seed)
Corn	<i>Pythium</i> damping-off control	0.75 to 1.50
	Systemic downy mildew control	3.00
Cucurbit vegetables	<i>Pythium</i> damping-off control	0.75
Fruiting vegetables	<i>Pythium</i> damping-off control	0.75 to 1.50
Leafy vegetables	<i>Pythium</i> damping-off control	0.75 to 1.50
Millet	<i>Pythium</i> damping-off control	0.75 to 1.50
	Systemic downy mildew control	2.50
Onions	<i>Pythium</i> damping-off control	0.75 to 1.50
Peas	<i>Pythium</i> damping-off control and early season <i>Phytophthora</i> control	3.50
Root and Tuber vegetables	<i>Pythium</i> damping-off control	0.75 to 1.50
	Note: Do not apply this product to potatoes.	
Sorghum	<i>Pythium</i> damping-off control	0.75 to 1.50
	Systemic downy mildew control	5.25
Sunflower	Systemic downy mildew control	9.00

Seed treated in accordance with the export use pattern must be labeled as follows:

**FOR EXPORT ONLY
NOT FOR DOMESTIC SALES OR USE**

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store in a cool, dry location, out of direct sunlight, and away from exposure to air and moisture. Do not store in temperatures over 90°F for prolonged period of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 56 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC: 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, Sharda USA LLC, MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

AXYL SHIELD ST

***FOR USE AS A SEED TREATMENT FOR CONTROL OF SYSTEMIC
DOWNY MILDEW, PYTHIUM SEED ROT, PYTHIUM DAMPING-OFF, AND
EARLY SEASON PHYTOPHTHORA DISEASES OF CERTAIN CROPS.***

ACTIVE INGREDIENT:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester..... 30.00%

OTHER INGREDIENTS: 70.00%

TOTAL: 100.00%

Contains 2.7 pounds metalaxyl per gallon (323 grams per liter).

EPA REG. NO. 83529-36

EPA EST. NO. 70815-GA-001

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for complete First Aid, Precautionary Statements, Directions for Use, and Limitation of Warranty and Liability.

Manufactured for:

Sharda USA LLC



7217 Lancaster Pike Suite A
Hockessin, DE 19707

NET CONTENTS 2.5 GAL. (9.46 L)

PEEL BACK BOOK HERE ▲