

Sharda Imidacloprid 5SC

INSECTICIDE

Active Ingredient:	
Imidacloprid 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	48.7%
Other Ingredients:	51.3%
Total:	100.0%
Contains 5 lbs. active ingredient per gallon (600 grams per liter).	

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA Reg. No. 83529-14

EPA Est. No. **AG** 72159-GA-001; **MA** 83411-MN-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Manufactured For:

Sharda USA LLC  7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

Net Contents: **1 Gal.**

FIRST AID	
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
For 24-hour medical emergency assistance (human or animal) call 1-800-222-1222 . For chemical emergency assistance (spill, leak, fire, or accident) call ChemTrec at 1-800-424-9300 .	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made out of any waterproof material (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride, or viton, selection Category A).

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride or Viton)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately, from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds.

Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treatment facilities. Use is also permitted as an end-use seed treatment on agricultural establishments at planting, or immediately before planting, as specified in the **Specific Crop Use Directions**. This product is to be used in liquid or slurry treaters only.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Mix thoroughly before use or use entire container at one time. All tank mixes should be pre-tested to determine physical compatibility between formulations. Follow the most restrictive precautions and limitations on the labeling of all products used in mixtures.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception. If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride, or Viton)
- Shoes plus socks
- Protective eyewear

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

Pesticide Storage: Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): Do not reuse container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): Do not reuse container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

STORED SEED PROTECTION

To protect stored seed of crops listed on this label, apply a minimum of 0.8 fluid ounces of **Sharda Imidacloprid 5SC** per 100 pounds of seed. Pests controlled by this application include Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryzae*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*). If seed being treated is known to have existing pest infestation, fumigation may be necessary prior to bagging.

EARLY SEASON PROTECTION AGAINST SUCKING INSECTS

Crop	Pest	Rate (fl. oz./ hundredweight seed)	Comments
Canola, Rapeseed and Mustard Seed	Aphids Flea Beetles Wireworms	10.24 - 25.6	Use higher rate within the specified rate range when pest populations are above historical levels.
	Lygus Bugs (including <i>Lygus</i> spp.) Seedpod Weevil Larvae (including <i>Ceutorhynchus assimilis</i>)	15.36 - 25.6	Suppression only. Use higher rate within the specified rate range when pest populations are above historical levels.

Canola, Rapeseed and Mustard Seed (For end use application at agricultural establishments):

Shake product well before making slurry and apply using a HCBT or Batch Treater. Apply 5.1 - 12.8 fl. oz. of **Sharda Imidacloprid 5SC** per 50 pound bag. May be diluted with an approved fungicide for disease control. Treat 1/2 of seed with 1/2 of slurry then add remainder of seed and slurry and continue mixing until all seed are covered.

Crop	Pest	Rate (fl. oz. per 80,000 units of seed)	Comments
Field Corn	Corn Root Worm (including Northern, Western, Southern, and Mexican) ⁽¹⁾ Flea Beetle Chinch Bug Southern Green Stinkbug White Grub Seed Corn Maggot Thrips Wireworm Corn Leaf Aphid Imported Fire Ant Southern Corn Leaf Beetle Billbug ⁽²⁾ Grape Colaspis ⁽²⁾ Black Cutworm ⁽³⁾	6.0 (Equivalent to 1.34 milligrams a.i. per kernel)	⁽¹⁾ In areas of heavy to severe corn rootworm populations, protection will not be adequate. Use only in areas of light to moderate corn root-worm populations. Consult your State Agricultural Extension Service on levels of corn rootworm populations. ⁽²⁾ Reduces early season feeding damage. ⁽³⁾ Will reduce feeding damage caused by leaf feeding black cutworms that are 1/2 inch or less in length. ⁽⁴⁾ Reduces feeding damage during emergence and seedling stages.
	Flea Beetle Chinch Bug Seed Corn Maggot Thrips Wireworm Corn Leaf Aphid Imported Fire Ant Grape Colaspis ⁽²⁾ White Grub	2.70 (Equivalent to 0.60 milligrams a.i. per kernel)	
	Seed Corn Maggot (Seed Protection Only) Wireworm (Seed Protection Only) Flea Beetle (Through 1-leaf stage) Imported Fire Ant White Grub ⁽⁴⁾	0.72 (Equivalent to 0.16 milligrams a.i. per kernel)	

Field Corn (For end use application at agricultural establishments):

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8 - 10 fluid ounces per 50 pound bag of seed. Treat 1/2 of seed with 1/2 of slurry then add remainder of seed and slurry and continue mixing until all seed are covered. After slurry treatment, the addition of dry talc at the rate of 3/4 - 1 ounce per 50 pounds of seed is recommended to minimize risk moisture buildup in planter.

Crop	Pest	Rate (fl. oz./ hundredweight seed)	Comments
Wheat, Barley, Oats, Rye, Triticale	Aphids (including Bird Cherry-Oat, English Grain, Greenbug, and Russian Wheat Aphid) Hessian Fly	0.8 - 2.4	Reduces possible yellow dwarf virus disease from aphid vectoring. Use the higher rate within the specified rate range to increase time of protection and/or when insect pressure is heavy.
	Wireworm	0.13 - 0.26	Suppression only. Use the higher rate within the specified rate range to increase time of protection and/or when insect pressure is heavy.
	Grasshopper	1.2 - 2.4	Treated seed may be planted as a 50- to 60-foot border around the perimeter of the field to help min- imize early season damage from grasshoppers. Consult your local agricultural extension agent for grasshopper control information specific to your area. Use the higher rate within the specified rate range to increase time of protection and/or when insect pressure is heavy.

Wheat, Barley, Oats, Rye, Triticale (For End-Use Application at Agricultural Establishments):

Make application using a Total Slurry Treater (TST), Farmer Applied Seed Treater (F.A.S.T.) or other on-farm seed treating equipment capable of accurately applying low rates of **Sharda Imidacloprid 5SC**. Apply 0.8 - 2.4 fl. oz. per hundredweight of seed. **Sharda Imidacloprid 5SC** may be mixed with a fungicide product for protection against diseases. **Sharda Imidacloprid 5SC** may be applied to seed previously treated with a fungicide. In this case, dilution is necessary.

Do not graze or feed livestock on treated areas for 45 days after planting.

Crop	Pest	Rate (fl. oz./ hundredweight seed)	Comments
Sorghum	Aphids (including Corn Leaf, English Grain, Greenbug, and Yellow Sugar Cane Aphid) Chinch Bugs Imported Fire Ants Wireworms	6.4	Thorough coverage is necessary for optimal control.

Sorghum (For End-Use Application at Agricultural Establishments):

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8 - 10 fl. oz. per 50 pound bag of seed. Treat 1/2 of seed with 1/2 of slurry then add remainder of seed and slurry and continue mixing until all seed are covered. After slurry treatment, the addition of dry talc at the rate of 3/4 - 1 ounce per 50 pounds of seed is recommended to minimize risk moisture buildup in planter.

Do not graze or feed livestock on treated areas for 45 days after planting.

Crop	Pest	Rate (fl. oz./ hundredweight seed)	Comments
Cotton (Delinted Seed Only)	Thrips & Aphids	12.8 (Equivalent to 0.375 milligram a.i./seed)	Regardless of the type of application (seed treatment, soil, or foliar) do not apply more than a total of 0.5 lb. of imidacloprid/acre/cropping cycle.

Cotton (Delinted Seed Only) (For End-Use Application at Agricultural Establishments):

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8 - 10 fluid ounces per 50 pound bag of seed. Treat 1/2 of seed with 1/2 of slurry then add remainder of seed and slurry and continue mixing until all seed are covered.

Crop	Pest	Rate (fl. oz. per hundredweight seed)⁽¹⁾	Comments
Sugar Beets (Commercial Seed Treatment)	Whitefly Aphids Leafhoppers ⁽²⁾ Root Aphid Thrips Wireworms	2.4 - 5.0	⁽¹⁾ On a unit of pelleted sugar beet seed with a weight ratio of 2:1 pelleting mixture to raw seed (seed count 100,000 seed – approximately 1 kilogram by weight). ⁽²⁾ Including to reduce risk of spread of Curly top and Yellow mosaic virus. If applying to non-pelleted raw seed, do not exceed a rate of 2.4 fluid ounces per 100,000 seed (approximately 1 kilogram by weight).

Crop	Pest	Rate (fl. oz. per hundredweight seed)	Comments
Sweet Corn (Commercial Seed Treatment)	Flea Beetle Early Season Corn Leaf Aphid Seed Corn Maggot Wireworm	6.4	Mix slurry so that it is applied at a rate of 16 - 20 fl. oz. of diluted solution per hundredweight of seed.
	Imported Fire Ant Early Season Corn Leaf Aphid Seed Corn Maggot Wireworm	3.2 - 6.4	
	Imported Fire Ant Seed Corn Maggot (Seed Protection) Wireworm (Seed Protection)	1.6 - 3.2	
	Wireworm (Seed Protection)	0.8 - 1.6	
Popcorn (Commercial Seed Treatment)	Flea Beetles	6.4	

Cotton (Delinted Seed Only) (For End-Use Application at Agricultural Establishments): *(continued)*

Crop	Pest	Rate (fl. oz. per hundredweight seed)	Comments
Soybean	Seed Corn Maggot Soybean Aphids Over-Wintering Bean Leaf Beetles	1.6 - 3.2	Use higher rate within the specified rate range to provide increased length of protection and for heavy insect pressure. Can be used as an over-treatment. Do not graze or feed livestock on soybean forage or hay.
Adzuki Bean, Asparagus Bean, Broad Bean (Succulent or Dry), Catjang Bean, Chinese Longbean, Field Bean, Guar Bean, Jackbean, Kidney Bean, Lablab Bean, Lima Bean (Succulent or Dry), Moth Bean (Succulent or Dry), Mung Bean, Navy Bean, Pinto Bean, Rice Bean, Runner Bean, Snap Bean, Sword Bean, Tepary Bean, Urd Bean, Wax Bean, Yardlong Bean, Blackeyed Pea (Succulent or Dry), Chickpea, Cowpea (Succulent or Dry), Crowder Pea, Dwarf Pea, Edible-Pod Pea, English Pea, Field Pea, Garden Pea, Green Pea, Pigeon Pea (Succulent or Dry), Snow Pea, Southern Pea (Succulent or Dry), Sugar Snap Pea, Grain Lupin, Sweet Lupin, White Lupin, White Sweet Lupin, Lentil*	Wireworm Bean Leaf Beetle Imported Fire Ant Aphid	1.6 - 3.2	* Seed-and-pod vegetable seed treated in California must be destined for planting in states other than California and is not to be planted in California.
Carrot	Seed Corn Maggot Wireworm	6.4	

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals must be observed for listed crops:

Rotational Plant-Back Intervals*				
Immediate Plant-back				30-Day Plant-back
Artichoke Barley Brassica (cole) Leafy Vegetable Canola Cilantro Corn, Field Corn, Sweet Cotton Cranberry	Cucurbits Eggplant Ground Cherry Leaf Petiole Vegetable Leafy Vegetables Legume Vegetable (Succulent or Dried, Except Soybean) Millet Mustard Seed	Oats Okra Pepinos Pepper Popcorn Potato Rapeseed Rye Safflower Sorghum	Soybean Strawberry Sugar Beet Tomatillo Tomato Triticale Vegetable Root and Tuber Watercress Wheat	Cereals, including: Buckwheat Rice
* Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed. * For all other crops not listed on an Imidacloprid label, or for crops for which no tolerances for the active ingredient has been established, a 12-month plant-back interval must be observed.				

Notification of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

NOTE: The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40 CFR, Part 180.1001. Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with **Sharda Imidacloprid 5SC** must be labeled in accordance with all applicable requirements of the Federal Seed Act.

USE RESTRICTIONS

- **Rape greens grown and harvested from Sharda Imidacloprid 5SC treated seed** must not be used for human and feed consumption.
- **Rapeseed grown and harvested from treated seed** is only for industrial uses and must not be used for edible oil or any other human/feed consumption.
- **Wheat, Barley, Oats, Rye, Triticale, and Sorghum:** Do not graze or feed livestock on treated areas for 45 days after planting.
- **Cotton (delinted seed only):** Regardless of the type of application (seed treatment, soil, or foliar) do not apply more than a total of 0.5 lb. imidacloprid per acre per cropping cycle.
- Do not use as a planter (hopper) box treatment.
- Do not use treated seed for feed, food, or oil purposes.
- Store treated seed away from feeds and foodstuffs.

- Do not allow children, pets, or livestock to have access to treated seed.
- The maximum application rate for imidacloprid (including seed treatments, foliar applications, and soil applications) is 0.5 lb. per acre per year. Do not apply more than 0.5 lb. a.i. imidacloprid per acre per year.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed may be hazardous to birds. Cover or incorporate spilled treated seed. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.

SEED BAG LABELING REQUIREMENTS

Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEEDS BE LABELED WITH THE FOLLOWING STATEMENTS:

- This seed has been treated with **Sharda Imidacloprid 5SC** (imidacloprid).
- Do not use for feed, food, or oil purpose.

THE US ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON BAGS CONTAINING SEEDS TREATED WITH **SHARDA IMIDACLOPRID 5SC** (IMIDACLOPRID):

- **Pollinator Precautions:** Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants, and chemical-resistant gloves when handling treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed may be hazardous to birds. Cover or incorporate spilled treated seeds. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash water.
- Do not allow children, pets, or livestock to have access to treated seed.
- In the event of a crop failure or harvest of a crop grown from **Sharda Imidacloprid 5SC** treated seed, the field may be replanted immediately to artichoke, barley, Brassica (cole) leafy vegetables, canola, cilantro, corn (field or sweet), cotton, cranberry, cucurbits, eggplant, ground cherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried including soybean), millet, mustard seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeets, tomatillo, tomato, triticale, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat, and rice, the minimum plant-back interval is 30 days from the date **Sharda Imidacloprid 5SC** treated seed was planted. Cover crops for soil building or erosion control may be planted at any time; but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid label, or for crops for which no imidacloprid, tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.
- The maximum application rate (including seed treatment, foliar application, and soil application) per acre per year for imidacloprid is 0.5 lb.
- This seed has been treated with X lbs. imidacloprid per lbs. of seed [or X mg imidacloprid per seed].
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

Conditions of Sale and Limited Warranty Statement

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Sharda USA LLC, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sharda USA LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Sharda USA LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SHARDA USA LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

Sharda Imidacloprid 5SC

Active Ingredient:

Imidacloprid 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine

48.7%

Other Ingredients:

51.3%

Total:

100.0%

Contains 5 lbs. active ingredient per gallon (600 grams per liter).

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID - IF SWALLOWED: • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice. **IF INHALED:** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. **IF IN EYES:** • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For 24-hour medical emergency assistance (human or animal) call **1-800-222-1222**. For chemical emergency assistance (spill, leak, fire, or accident) call ChemTrec at **1-800-424-9300**.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long sleeved shirt, long pants, shoes, socks, and chemical-resistant

gloves made out of any waterproof material (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride, or viton, selection Category A).

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds. Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited. **Pesticide Storage:** Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal: NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS):** Do not reuse container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Manufactured For: Sharda USA LLC,

7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 83529-14

EPA Est. No. (AG) 72159-GA-001; (MA) 83411-MN-001;

(MC) 89332-GA-001; (SC) 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: **1 Gal.**

OPEN HERE