

Cyril WDG Fungicide

CYPRODINIL GROUP 9 FUNGICIDE

FUNGICIDE



ACTIVE INGREDIENT:	WT. BY %
Cyprodinil: 4-cyclopropyl-6-methyl-N-phenylpyrimidin-2-amine	75.0%
OTHER INGREDIENTS:	25.0%
TOTAL:	100.0%
Cyril WDG Fungicide is a water-dispersible granule containing 75% cyprodinil.	

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

See label booklet for complete First Aid, Precautionary Statements, Directions For Use, and Storage and Disposal.

Manufactured For:

Sharda USA LLC

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 82633-132-83529

EPA Est. No. **(AG)** 72159-GA-001; **(MA)** 83411-MN-001; **(SC)** 39578-TX-001; **(TX)** 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: **50 oz.**

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378 , Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	
NOTE TO PHYSICIAN	
If ingested, induce emesis or lavage stomach. Treat symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Handlers applying this product as a pre-plant dip to strawberry roots and crowns and workers packaging or preparing treated roots and crowns for shipment must wear:

- Chemical-resistant apron made of any waterproof material
- Elbow-length chemical-resistant gloves made of any waterproof material
- Chemical-resistant boots made of any waterproof material

All other applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or Viton[®] \geq 14 mils
- Shoes plus socks

In addition, mixers and loaders for aerial and ground boom applications must wear:

- Filtering facepiece respirator (N95, R95, or P95) (e.g., a dust mask)

Mixer/loaders supporting ground boom applications to corn, canola, soybean, cotton, must wear:

- Long-sleeved shirt and long pants
- Shoes and socks
- Chemical-resistant gloves including barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton[®] \geq 14 mils

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS**Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Surface and Ground Water Advisory

This chemical may contaminate water through runoff. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This chemical has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this chemical. A level, well maintained vegetative buffer strip between areas to which this chemical is applied and surface water features including ponds, streams, and springs will reduce the potential loading of cyprodinil from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

Physical or Chemical Hazards

DO NOT use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

PRODUCT INFORMATION

Cyrl WDG Fungicide is a broad spectrum fungicide which controls certain diseases in fruits, herbs, nuts and vegetables.

PRODUCT USE PRECAUTIONS

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

APPLICATION INFORMATION

Cyrl WDG Fungicide has preventative and systemic properties and is labeled for the control of many important plant diseases. **Cyrl WDG Fungicide** provides excellent disease control of Botrytis, several leaf spots and powdery mildews. **Cyrl WDG Fungicide** is applied as a foliar spray and can be used in block, alternating spray, or tank-mix programs with other crop protection products. Make all applications according to the use directions that follow.

PRODUCT USE INSTRUCTIONS

Application: Thorough coverage is necessary to provide good disease control. Make no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Adjuvants: When an adjuvant is to be used with this product, Sharda advises the use of a Chemical Producers and Distributors Association certified adjuvant.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if the maximum amount of **Cyrl WDG Fungicide** has been used. If resistant isolates to Group 9 fungicides are present, efficacy can be reduced

for certain diseases. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, with highly susceptible varieties, or when environmental conditions are conducive to disease.

Integrated Pest Management (IPM): Integrate **Cyril WDG Fungicide** into an overall disease and pest management strategy whenever the use of a fungicide is required. Follow cultural practices known to reduce disease. Consult your local agricultural authorities for additional IPM strategies established for your area. **Cyril WDG Fungicide** may be used in state agricultural extension advisory (disease forecasting) programs which direct application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

CYPRODINIL	GROUP	9	FUNGICIDE
------------	-------	---	-----------

For resistance management, **Cyril WDG Fungicide** contains a Group 9 fungicide/bactericide. Any fungal/bacterial population may contain individuals naturally resistant to **Cyril WDG Fungicide** and other Group 9 fungicides/bactericides. A gradual or total loss of pest control may occur over time if these fungicides/bactericides are used repeatedly in the same fields. Appropriate resistance management strategies must be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of **Cyril WDG Fungicide** or other Group 9 fungicides/bactericides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM guidelines for specific crops and pathogens.
- For information or to report suspected resistance, contact your local Sharda Cropchem, Ltd. representative.

Crop Tolerance: Plant tolerance has been found to be acceptable for all crops on the label, however, not all possible tank-mix combinations have been tested under all conditions. When possible, test the combinations on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application.

Rotational Crop Restrictions

DO NOT plant any crop which is not registered for use with cyprodinil for a period of 30 days, unless a shorter interval is specified on the following list.

Rotational Crop	Planting Time from Last Cyril WDG Fungicide Application
Artichoke, Globe Beans (dried and succulent except cowpeas) Berries (bushberries 13-07B, caneberries 13-07A) Brassica, Leafy Greens (Crop Subgroup 4-16B) Brassica, Head and Stem (Crop Group 5-16) Celtuce Citrus Fruit (Crop Subgroup 10-10B) Cucurbits (Crop Group 9) Fennel, Florence, fresh leaves and stalk Herbs (fresh and dried) Kohlrabi Leafy Greens (Crop Subgroup 4-16A) Leaf Petiole Vegetables (Crop Subgroup 22B) Leaves of Root and Tuber Vegetables Onions (dry bulb, garlic, and green) Peppers Tuberous and Corm Vegetables (Crop Subgroup 1C) Root and Tuber Vegetables, except Sugar beet (Crop Subgroup 1B) Strawberries Tomatoes and Tomatillos Watercress Crops Not Intended for Food or Feed	0 days
All Other Crops Intended for Food or Feed	30 days

Restriction: In annual crops, where multiple crops can be grown per year (double/triple cropping), **DO NOT** apply more than 1.3 lb. a.i. per acre per year to an individual plot of land.

APPLICATION AND MIXING PROCEDURES

Mixing

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. **DO NOT** let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area. Buffer the spray solution to a pH of 5.0 - 7.0 when tank mixed with a fungicide containing Iprodione.

Cyril WDG Fungicide Alone: Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the **Cyril WDG Fungicide** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the **Cyril WDG Fungicide** has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Cyril WDG Fungicide + Tank Mixtures: Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank-mix partners. Add tank-mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) including **Cyril WDG Fungicide**, liquid flowables, liquids, and emulsifiable concentrates. Always allow each tank-mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

When using **Cyril WDG Fungicide** in tank mixtures, add all products in water-soluble packaging to the tank before any other tank-mix partner, including **Cyril WDG Fungicide**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix partner to the tank.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If using **Cyril WDG Fungicide** in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. **DO NOT** exceed label dosage rates and follow the most restrictive label precautions and limitations. **Cyril WDG Fungicide** must not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Additives: **Cyril WDG Fungicide** is compatible with most crop protection additives. **DO NOT** use X-77[®] with **Cyril WDG Fungicide** for bloom sprays applied to stone fruits, almonds and pistachios.

Application

For best disease control, apply **Cyril WDG Fungicide** in sufficient water to provide thorough and uniform coverage. Use minimum ground spray volumes of 50 gal./A for tree crops, 30 gal./A for vine crops, and 15 gal./A for field and vegetable crops. For aerial application, see spray volume requirements in the specific crop directions for use.

To prevent spray drift, **DO NOT** apply when conditions favor drift beyond the target area. Spray overlap may cause crop injury.

For air assisted or air blast sprayers, move spray droplets into the canopy using a forced air stream. Set up the fan to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area.

Equip sprayers with nozzles that provide accurate and uniform application. Check whirl plates and nozzle discs for wear and replace as necessary. Calibrate sprayer before use.

Use a pump with capacity to maintain the correct rated pressure for the nozzles selected. Maintain sufficient agitation to keep the spray mixture in suspension. Use a jet agitator, liquid sparge tube, or mechanical paddle for agitation. **DO NOT** air sparge.

Use screens to prevent nozzles from clogging. Use 50-mesh or coarser screens placed after the tank and before the nozzles. Check nozzle manufacturer's specifications.

For more information on spray equipment and calibration, consult sprayer manufacturers and state directions. For specific local directions and spray schedules, consult the current state agricultural experiment station directions.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH PONDS.

- **DO NOT** apply within 75 ft. of bodies of water including lakes, reservoirs, rivers, permanent streams, natural ponds, marshes or estuaries.
- Shut off the sprayer when row ends.
- **DO NOT** cultivate within 10 ft. of aquatic areas in order to allow a vegetative filter strip.
- **DO NOT** apply when weather conditions favor drift to aquatic areas. **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during a temperature inversion. Mist or fog may indicate the presence of an inversion in humid areas.
- For perennial crops including tree crops and grapes:
 - o For all plantings within 150 ft. of bodies of water as described above, spray crops from outside the planting away from the bodies of water.
 - o Spray last three rows windward of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas. Adjust or turn off top nozzles to prevent spray going over the tops of trees. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row or passing tree gaps in the rows.

Ground Spray Drift Restrictions

DO NOT apply when wind speeds exceed 15 miles per hour at the application site.

Aerial Spray Directions

Avoid applications under conditions when uniform coverage cannot be obtained or when excessive drift may occur.

Aerial Spray Drift Restrictions

Observe the following restrictions when spraying in the vicinity of aquatic areas including lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.

- Use only on crops where aerial applications are indicated.
- **DO NOT** apply by air within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Release spray at the lowest height consistent with pest control and flight safety.
- **DO NOT** make applications more than 10 ft. above the crop canopy.
- **DO NOT** apply when weather conditions favor drift to aquatic areas.
- **DO NOT** apply during a temperature inversion. Mist or fog may indicate the presence of an inversion in humid areas.

Aerial Spray Precautions

Observe the following precautions when spraying in the vicinity of aquatic areas including lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.

- Use the largest droplet size consistent with good pest control.
- Formation of very small droplets may be minimized by appropriate nozzle selection, by orientating nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Reduce risk of exposure to aquatic areas by avoiding applications when wind direction is toward the aquatic area.
- Low humidity and high temperatures increase the evaporation rate of spray droplets, and therefore the likelihood of increased spray drift to aquatic area. Avoid spraying during conditions of low humidity and/or high temperatures.

SPRAY DRIFT MANAGEMENT

Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure directed for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

Adjust Nozzles - Follow nozzle manufacturer's directions for setting up nozzles. To reduce fine droplets, orient nozzles parallel with the air flow in flight.

Boom Height – Ground Boom

- Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage.
- For ground equipment, the boom must remain level with the crop and have minimal bounce.

Release Height – Aircraft

- Higher release heights increase the potential for spray drift.
- When applying aurally to crops, **DO NOT** release spray at a height greater than 10 ft. above the crop canopy, unless a greater application height is necessary for pilot safety.

Shielded Sprayers

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.125 - 0.25 inches/A of water. Excessive water may reduce efficacy.
- If you have questions about calibration, contact state extension service specialists, equipment manufacturers, or other experts.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, for example a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended.

Center Pivot Irrigation Equipment

Restrictions:

- (1) **DO NOT** inject **Cyril WDG Fungicide** at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part **Cyril WDG Fungicide**. **Cyril WDG Fungicide** is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used but must be replaced once a year. **DO NOT** use Viton®, Buna-N, Neoprene, or PVC seals.
- (2) Use only with drive systems which provide uniform water distribution.
- (3) **DO NOT** use end guns when chemigating **Cyril WDG Fungicide** through center pivot systems because of non-uniform application.
 - Determine the size of the area to be treated.
 - Determine the time required to apply 1/8 - 1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as directed by the equipment manufacturer. When applying **Cyril WDG Fungicide** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80 - 95% of the manufacturer's rated capacity.
 - Using water, determine the injection pump output when operated at normal line pressure.
 - Determine the amount of **Cyril WDG Fungicide** required to treat the area covered by the irrigation system.
 - Add the required amount of **Cyril WDG Fungicide** and sufficient water to meet the injection time requirements to the solution tank.
 - Make sure the system is fully charged with water before starting injection of the **Cyril WDG Fungicide** solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
 - Maintain constant solution tank agitation during the injection period.
 - Continue to operate the system until the **Cyril WDG Fungicide** solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20- to 30-minute interval. When applying **Cyril WDG Fungicide** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of **Cyril WDG Fungicide** required to treat the area covered by the irrigation system.
- Add the required amount of **Cyril WDG Fungicide** into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the **Cyril WDG Fungicide** solution has cleared the last sprinkler head.

SPECIFIC INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, for example a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

CROP USE DIRECTIONS – TREES, NUTS AND VINES

Crop	Disease	Product Rate oz./Acre	Use Directions
<p>Almonds</p>	<p>Brown rot blossom blight (<i>Monilinia</i> spp.) Suppression: Green fruit rot (Jacket rot) (<i>Botrytis cinerea</i>) Shot hole (<i>Wilsonomyces carpophilus</i>)</p>	<p>5 - 10*</p>	<p>Apply Cyril WDG Fungicide at 5 - 10% bloom. Additional applications at 50 - 100% bloom and petal fall may be necessary. When used for control of brown rot blossom blight, Cyril WDG Fungicide will provide suppression of shot hole. For broad spectrum disease control in tank mixture, apply Cyril WDG Fungicide at a minimum rate of 5 oz. in tank mixtures with other fungicides registered for use on almonds. For suppression of green fruit rot, apply Cyril WDG Fungicide at full bloom. Disease suppression for almond diseases refers to erratic control from fair to good, or consistent control at a level below that obtained with products registered for control.</p>
<p>*5 oz. product contains 0.24 lb. a.i. cyprodinil/A *10 oz. product contains 0.47 lb. a.i. cyprodinil/A</p>			
<p align="center">Application Instructions</p>			
<p>Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air.</p>			
<p align="center">Specific Use Restrictions</p>			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for almonds. • DO NOT apply more than 3 applications per year at the highest rate. • Minimum Application Interval: 14 days • DO NOT apply more two applications by air. • DO NOT apply more than 30 oz./A of Cyril WDG Fungicide (1.4 lb. a.i./A of cyprodinil) per year. • DO NOT apply within 60 days of harvest (60-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Citrus, Crop Subgroup 10-10B** Lemon Lime Not for Use in California	Alternaria stem end rot (<i>A. citri</i>) Anthracnose (<i>Colletotrichum gloeosporioides</i>) Blue mold (<i>Penicillium italicum</i>) Green mold (<i>Penicillium digitatum</i>)	5.5 - 7*	Make one application near harvest to prevent post-harvest fruit rot. The application may be made up to and including the day of harvest.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
**Complete List of Citrus Crop Group 10-10B			
Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; kumquat; lemon; lime; mount white lime; New Guinea wild lime; Russell River lime; sweet lime; Tahiti lime; cultivars, varieties, and/or hybrids of these.			
Application Instructions			
Good coverage is essential for good disease control.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Citrus, Crop Subgroup 10-10B. • DO NOT apply more than 1 application per year at the highest rate. • Application must be made by ground only. • DO NOT apply more than 7 oz./A of Cyrol WDG Fungicide (0.33 lb. a.i./A of cyprodinil) per year. • DO NOT apply more than 0.33 lb. a.i./A of cyprodinil-containing products per year. • May be applied on the day of harvest (0-day PHI). • DO NOT exceed one application per year. 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Small Fruit Vine Climbing Subgroup 13-07F (except Fuzzy kiwifruit) Grapes Amur river grape Hardy kiwifruit Maypop Schisandra berry And cultivars and/or hybrids of these	Botrytis bunch rot <i>(Botrytis spp.)</i> Suppression: Powdery mildew <i>(Uncinula necator)</i>	Cyril WDG Fungicide alone 10*	Begin applications of Cyril WDG Fungicide at early bloom. Make an additional application at berry touch, veraison, or preharvest using at least a 7-day spray interval. Botrytis bunch rot is most effectively controlled by ground application, using sufficient water volume to provide thorough coverage. Thorough coverage of bunches is essential. When used at 10 oz./A, Cyril WDG Fungicide will provide significant suppression (approximately 60% control) of powdery mildew.
		Cyril WDG Fungicide tank mixtures 5 - 10*	Apply Cyril WDG Fungicide in tank mixture with the label rate of another fungicide registered on grapes for control of Botrytis bunch rot.
*5 oz. product contains 0.24 lb. a.i. cyprodinil/A			
*10 oz. product contains 0.47 lb. a.i. cyprodinil/A			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Small Fruit Vine Climbing Subgroup 13-07F. • DO NOT apply more than 3 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT apply more than two applications by air. • DO NOT apply more than 30 oz./A of Cyril WDG Fungicide (1.4 lb. a.i./A of cyprodinil) per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Kiwi	Botrytis fruit rot (<i>Botrytis</i> spp.)	10*	Make 1 - 2 applications on a 7- to 10-day interval prior to harvest to prevent post-harvest fruit rot.
*10 oz. product contains 0.47 lb. a.i. cyprodinil/A			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Kiwi. • DO NOT apply more than 2 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT apply more than 20 oz. of Cyril WDG Fungicide (0.94 lb. a.i./A of cyprodinil) per year. • May be applied on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Pistachios	Botrytis (<i>Botrytis</i> spp.) Alternaria (<i>Alternaria alternata</i>)	5.5 - 7*	Make the first application during early bloom and repeat applications at 14-day intervals if conditions remain favorable for disease development. After 2 applications, alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Pistachios. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 14 days • DO NOT apply more than two applications by air. • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Pome Fruits** Apples Crabapples Loquat Mayhaw Pears (see tank mixture instructions) Quince And cultivars and/or hybrids of these	Prebloom Scab <i>(Venturia spp.)</i>	Cyril WDG Fungicide alone 5*	For pome fruits except pear, begin application at green tip and continue on a 7- to 10-day interval. Under severe disease pressure, use the shorter interval.
		Cyril WDG Fungicide tank mixtures 3 - 5*	For scab control utilizing multiple modes of action, apply Cyril WDG Fungicide in tank mixture with the label rate of a protectant or systemic fungicide registered on pome fruit. Make applications on a 7- to 10-day interval.
	Pink, bloom, post-bloom Scab <i>(Venturia spp.)</i>	Cyril WDG Fungicide tank mixtures 3 - 5*	Apply Cyril WDG Fungicide in tank mix combination with the label rate of a protectant fungicide. Use of the label rate of an EBDC fungicide will broaden the disease control spectrum to include cedar apple rust. The addition of the label rate of sulfur or a sterol inhibitor (SI) fungicide to this tank mix will broaden the disease control spectrum to include powdery mildew.
*3 oz. product contains 0.14 lb. a.i. cyprodinil/A *5 oz. product contains 0.24 lb. a.i. cyprodinil/A			
**Complete List of Pome Fruit Crops			
Apple; Azarole; Crabapple; Loquat; Mayhaw; Medlar; Pear; Pear, Asian; Quince; Quince, Chinese; Quince, Japanese; Tejocote and cultivars, varieties and/or hybrids of these.			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Pome Fruits. • DO NOT apply more than 6 applications per year at the highest rate. • Minimum Application Interval: 7 days • When applying to pears, apply Cyril WDG Fungicide in tank mix only. • DO NOT apply more than two applications by air. • DO NOT apply more than 30 oz./A of Cyril WDG Fungicide (1.4 lb. a.i./A of cyprodinil) per year. • Apply up to and on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Stone Fruits Crop Group 12-12** Apricots Tart cherries Nectarines Peaches Plums Prunes And cultivars and/or hybrids of these	Brown rot blossom blight <i>(Monilinia spp.)</i>	5*	Begin applications at bloom stage (apricots at red bud, cherries at popcorn, peaches and nectarines at pink bud, plums and prunes at green tip). Make a second application at full bloom. If disease pressure persists, make subsequent applications on a 7- to 10-day interval. Use Cyril WDG Fungicide alone or in tank mixture with the label rate of another fungicide registered for stone fruit.
	California only Fruit brown rot <i>(Monilinia spp.)</i>	Cyril WDG Fungicide alone 10*	Apply a maximum of 2 applications of Cyril WDG Fungicide during the preharvest period on a 7-day interval up through 2 days prior to harvest as needed.
		Cyril WDG Fungicide tank mixtures 5*	Apply Cyril WDG Fungicide in tank mixture with the label rate of another fungicide registered on stone fruit for control of fruit brown rot.
*5 oz. product contains 0.24 lb. a.i. cyprodinil/A			
*10 oz. product contains 0.47 lb. a.i. cyprodinil/A			
**Complete List of Stone Fruit			
Apricot; Apricot, Japanese; Cherry, tart; Jujube, Chinese; Nectarine; Peach; Plum; Plum, American; Plum, beach; Plum, Canada; Plum, cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plum, prune; Plumcot; Prune; Sloe; as well as other cultivars and hybrids of these.			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Stone Fruits. • Crop Group 12-12. • DO NOT apply Cyril WDG Fungicide to sweet cherries. • Minimum Application Interval: 7 days • DO NOT apply more than two applications by air. • DO NOT apply more than 30 oz./A of Cyril WDG Fungicide (1.4 lb. a.i./A of cyprodinil) per year. • DO NOT apply more than a maximum total of 4 applications (air plus ground) per year when applied at reduced rates. • DO NOT apply within 2 days of harvest (2-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Tropical and Subtropical, Small fruit, inedible peel Subgroup 24A** Lychee Longan Spanish lime And cultivars and/or hybrids of these	Botrytis fruit rot <i>(Botrytis spp.)</i> Alternaria fruit rot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i>	5.5 - 7*	Make the first application during early bloom and repeat on 7- to 10-day intervals if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyrol WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A *7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
**Complete List of Subgroup 24A Crops			
Aisen; bael fruit; Burmese grape; cat's-eyes; inga; longan; Lychee; madras-thorn; manduro; matisia; mesquite; mongongo, fruit; pawpaw, small-flower; satinleaf; Sierra Leone-tamarind; Spanish lime; velvet tamarind; wampi; white star apple; cultivars, varieties, and hybrids of these commodities.			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Tropical and Subtropical, Small fruit, inedible peel Subgroup 24A. • Make no more than two applications by air. • Minimum Application Interval: 7 days • DO NOT apply more than 28 oz./A of Cyrol WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply more than a maximum total of 4 applications (air plus ground) per year. • May be applied on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Specific Tropical Fruits Acerola Avocado Black sapote Canistel Dragon fruit Feijoa Guava Jaboticaba Mamey sapote Mango Papaya Passionfruit Pulasan Rambutan Sapodilla Star apple Starfruit Wax jambu	Botrytis fruit rot (<i>Botrytis</i> spp.) Alternaria fruit rot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.)	5.5 - 7*	Make the first application during early bloom and repeat on 7- to 10-day intervals if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Specific Tropical Fruits. • DO NOT apply more than two applications by air. • Minimum Application Interval: 7 days • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply more than a maximum total of 4 applications (air plus ground) per year. • Apply up to and on the day of harvest (0-day PHI). 			

CROP USE DIRECTIONS – BERRIES, FRUITS, HERBS AND VEGETABLES

Crop	Disease	Product Rate oz./Acre	Use Directions
Artichoke, Globe Not for Use in California	Ramularia leaf spot Ramularia bud spot <i>(R. cynarae)</i>	5.5 - 7.0*	Begin applications prior to disease onset when conditions are conducive for disease. Apply Cyrial WDG Fungicide on a 14-day schedule, making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A *7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
Application Instructions			
For best results, sufficient water volume must be used to provide thorough coverage. Cyrial WDG Fungicide can be applied by ground, chemigation, or aerial application. For ground applications, apply in 50 - 200 gallons of water per acre to obtain coverage without excessive runoff. For aerial applications, use a minimum of 10 gal./A of water. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Artichoke, Globe. • DO NOT apply more than two applications by air. • Minimum Application Interval: 14 days • DO NOT apply more than 28 oz./A of Cyrial WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply more than 1.3 lb. a.i./A/year of a cyprodinil-containing product. • DO NOT apply more than a maximum total of 4 applications (air plus ground plus chemigation) per year. • DO NOT apply Cyrial WDG Fungicide within 3 days of harvest (3-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Beans (Dried and Succulent except cowpeas) Chickpea (garbanzo bean) Bean (<i>Lupinus</i> spp.) (grain lupin, sweet lupin, white lupin, white sweet lupin) Bean (<i>Phaseolus</i> spp.) (kidney, lima, mung, navy, pinto, snap, wax) Broad Bean (fava bean) Bean (<i>Vigna</i> spp.) (asparagus, blackeyed pea) Not for Use in California	Gray mold (<i>Botrytis cinerea</i>)	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Beans (Dried and Succulent except cowpeas). • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT apply more than two applications by air. • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Berries Bushberry Subgroup 13-07B Aronia berry Black currant Blueberry, high and low bush Buffalo currant Chilean guava Edible honeysuckle Elderberry European barberry Gooseberry Highbush cranberry Huckleberry Jostaberry Juneberry (Saskatoon berry) Lingonberry Native currant Red currant Salal Sea buckthorn Not for Use in California Caneberry Subgroup 13-07A Blackberry Loganberry Red and Black raspberry Wild raspberry And cultivars and/or hybrids of these Not for Use in California	Mummy berry (<i>Monilinia vaccinicorymbosi</i>) Anthracnose (<i>Colletotrichum</i> spp.) Alternaria fruit rot (<i>Alternaria tenuissima</i>) Botrytis fruit rot (<i>Botrytis cinerea</i>)	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A *7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air.			

(continued)

Specific Use Restrictions

- **Maximum Single Application Rate: DO NOT** exceed the maximum rate listed above for Berries Bushberry Subgroup 13-07B.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** apply more than two applications by air.
- **DO NOT** apply more than 28 oz./A of **Cyril WDG Fungicide** (1.3 lb. a.i./A of cyprodinil) per year.
- Apply up to and on the day of harvest (0-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Brassica Head and Stem Vegetable Crop Group 5-16** Broccoli Brussels sprouts Cabbage Cabbage, Chinese Cauliflower And cultivars and/or hybrids of these Not for Use in California	Powdery mildew <i>(Erysiphe polygoni)</i>	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
**Complete List of Brassica Head and Stem Vegetables Crop Group 5-16			
Broccoli; Brussels sprouts; Cabbage; Cabbage, Chinese (napa); Cauliflower cultivars, varieties, and hybrids of these commodities.			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Brassica Head and Stem Vegetable Crop Group 5-16. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT apply more than two applications by air. • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Brassica Leafy Greens Subgroup 4-16B** Arugula Chinese cabbage Bok Choy Collards Kale Mustard greens Turnip greens And cultivars and/or hybrids of these. See separate instructions for Watercress. Not for Use in California	Powdery mildew <i>(Erysiphe polygoni)</i>	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A *7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
**Complete list of Brassica Leafy Greens Vegetable Subgroup 4-16B			
Arugula; broccoli, Chinese; broccoli raab; cabbage, abyssinian; cabbage, seakale; cabbage, Chinese, bok choy; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; cultivars, varieties, and hybrids of these commodities.			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Brassica Leafy Greens Subgroup 4-16B. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • Make no more than two applications by air. • DO NOT use roots of treated turnips for food or feed. Only turnip varieties harvested for their leaves may be treated. • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Cucurbits Crop Group 9** Cantaloupe Chayote Chinese waxgourd Cucumber Gourds Honeydew Momordica spp. (Bitter melon, Balsam apple) Muskmelon Watermelon Pumpkin Squash Zucchini And cultivars and/or hybrids of these Not for Use in California	Alternaria leaf blight (<i>A. cucumerina</i>) Alternaria leaf spot (<i>A. alternate</i>) Gummy stem blight (<i>Didymella bryoniae</i>) Powdery mildew (<i>Sphaerotheca fuliginea</i> , <i>Erysiphe cichoracearum</i>)	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			

(continued)

****Complete List of Cucurbit vegetable Crop Group 9**

Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon); pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon.

Application Instructions

Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed above for Cucurbits Crop Group 9.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** apply more than two applications by air.
- **DO NOT** apply more than 28 oz./A of **Cyril WDG Fungicide** (1.3 lb. a.i./A of cyprodinil) per year.
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products per year.
- Apply up to 1 day before harvest (1-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Herbs* (Dried and Fresh) Basil Chive Coriander, leaves (cilantro) Dillweed Lemongrass Parsley Rosemary Sage Tarragon Thyme Not for Use in California	Alternaria leaf spot <i>(Alternaria spp.)</i> Botrytis leaf blight <i>(Botrytis spp.)</i>	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
**Additional Herbs (dried and fresh)			
Angelica, Balm, Borage, Burnet, Camomile, Catnip, Chervil (dried leaves), Clary, Costmary, Culantro (leaves), Curry (leaves), Horehound, Hyssop, Lavender, Lovage (leaves), Marigold, Marjoram, Nasturtium, Pennyroyal, Rue, Savory (summer and winter), Sweet bay, Tansy, Wintergreen, Woodruff, and Wormwood.			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Herbs. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT apply more than two applications by air. • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Kohlrabi Not for Use in California	Powdery mildew <i>(Erysiphe polygoni)</i>	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Kohlrabi. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • Make no more than two applications by air. • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
<p>Leafy Greens, Crop Subgroup 4-16A**, except Parsley Leaf Petiole Vegetables Crop Subgroup 22B*** Amaranth Cardoon Celery Celery, Chinese Celtuce Chervil Chrysanthemum, edible Corn salad Dandelion Dock Endive (escarole) Fennel, Florence Lettuce, head and leaf New Zealand spinach Orach Purslane Radicchio Rhubarb Spinach Swiss chard And cultivars and/or hybrids of these Not for Use in California</p>	<p>Alternaria leaf spot (<i>Alternaria</i> spp.) Gray mold (<i>Botrytis cinerea</i>) Suppression: Powdery mildew (<i>Erysiphe cichoracearum</i>)</p>	<p>5.5 - 7*</p>	<p>Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide, alternate with another fungicide with a different mode of action for 2 applications.</p>
<p>*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A *7 oz. product contains 0.33 lb. a.i. cyprodinil/A</p>			

(continued)

****Complete List of Leafy Greens Crop Subgroup 4-16A includes**

Amaranth, Chinese; amaranth, leafy; aster, Indian; blackjack; cat's whiskers; cham-chwi; cham-na-mul; chervil, fresh leaves; chipilin; chrysanthemum, garland; cilantro, fresh leaves; corn salad; cosmos; dandelion, leaves; dang-gwi, leaves; dillweed; dock; dol-nam-mul; ebolo; endive; escarole; fameflower; feather cockscomb; good king henry; huauzontle; jute, leaves; lettuce, bitter; lettuce, head; lettuce, leaf; orach; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; spinach; spinach, Malabar; spinach, New Zealand; spinach, tanier; Swiss chard; violet, Chinese, leaves; cultivars, varieties, and hybrids of these commodities.

*****Leaf Petiole Vegetables Crop Subgroup 22B includes**

Cardoon; celery; celery, Chinese; fuki; rhubarb; udo; zuiki; cultivars, varieties, and hybrids of these commodities.

Application Instructions

Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed above for Leafy Greens, Crop Subgroup 4-16A (except Parsley) and Leaf Petiole Vegetables Crop Subgroup 22B.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** apply more than two applications by air.
- **DO NOT** apply more than 28 oz./A of **Cyril WDG Fungicide** (1.3 lb. a.i./A of cyprodinil) per year.
- May be applied on the day of harvest (0-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Leaves of Root and Tuber Vegetables Crop Group 2** Carrot Beet, garden Beet, sugar Parsnip Radish Radish (Oriental) Rutabaga Sweet potato Turnip Yam (true) Not for Use in California	Alternaria leaf blight <i>(Alternaria dauci)</i> Powdery mildew <i>(Erysiphe spp.)</i>	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A *7 oz. product contains 0.33 lb. a.i. cyprodinil/A			

(continued)

****Additional Leaves of Root and Tuber Vegetables**

Burdock (edible), Cassava, Celeriac, Chicory, Dasheen, Salsify (black and Spanish), Tanier, and Turnip rooted chervil.

Application Instructions

Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions

- **Maximum Single Application Rate: DO NOT** exceed the maximum rate listed above for Leaves of Root and Tuber Vegetables Crop Group 2.
- **DO NOT** apply more than 4 applications per year at the highest rate except for radish.
- **Minimum Application Interval:** 7 days
- **DO NOT** apply more than two applications by air.
- Radish ONLY – **DO NOT** apply more than two applications per year.
- Radish ONLY – **DO NOT** apply more than 14 oz. of **Cyril WDG Fungicide** (0.66 lb. a.i./A of cyprodinil) per crop per year.
- Radish ONLY – **DO NOT** apply more than 0.66 lb. a.i./A of cyprodinil-containing products per year.
- **DO NOT** apply more than 28 oz./A of **Cyril WDG Fungicide** (1.3 lb. a.i./A of cyprodinil) per year for specified leaves of root and tuber vegetables except radish.
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products per year for specified leaves of root and tuber vegetables except radish.
- **DO NOT** apply within 7 days of harvest (7-day PHI).
- **DO NOT** allow cattle or other livestock to feed upon the leaves of root vegetables.

Crop	Disease	Product Rate oz./Acre	Use Directions
<p>Onions Bulb Vegetables Crop Group 3-07A and 3-07B Bulb Onion Chinese onion Dry bulb onion Daylily bulb Fritillaria bulb Garlic Great-headed garlic Lily bulb Pearl onion Potato onion Serpent garlic Shallot Green Onion Beltsville bunching onion Chinese chive fresh leaves Fresh chive leaves Fritillaria leaves Fresh onion Green onion Hosta elegans Kurrat Lady's leek Leek Macrostem onion Shallot fresh leaves Tree tops onion Welsh onion tops Wild leek Wild onion Onions Grown for Seed And cultivars and/or hybrids of these</p>	<p>Botrytis leaf blight or blast (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) Suppression: Neck rot (<i>Botrytis</i> spp.)</p>	<p>5.5 - 10*</p>	<p>Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. For optimal effect on neck rot, apply on a 7-day schedule at the 10 oz. rate. Resistance Management: After 2 applications of Cyril WDG Fungicide, alternate with another fungicide with a different mode of action for 2 applications.</p>
<p>*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A *7 oz. product contains 0.33 lb. a.i. cyprodinil/A</p>			

Application Instructions

Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed above for Onions, Bulb Vegetables Crop Group 3-07A and 3-07B, Bulb Onion, Green Onion, and Onions Grown for Seed.
- **DO NOT** apply more than 2 applications (air and ground) per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** apply more than two applications by air.
- **DO NOT** apply more than 28 oz./A of **Cyril WDG Fungicide** (1.3 lb. a.i./A of cyprodinil) per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Potatoes Tuberous and Corm Vegetables Crop Subgroup 1C** Sweet Potatoes Not for Use in California	Brown spot <i>(Alternaria alternata)</i> Early blight <i>(A. solani)</i> Powdery mildew <i>(Erysiphe cichoracearum)</i> Septoria leaf spot <i>(Septoria lycopersici)</i> Tan spot <i>(Botrytis cinerea)</i>	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
**Additional Vegetables, Tuberous and Corm Subgroup 1C			
Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna, Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen (taro), Ginger, Leren, Sweet Potato, Tanier, Turmeric, Yam (bean and true), and cultivars and/or hybrids of these.			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Potatoes, Tuberous and Corm Vegetables Crop Subgroup 1C, and Sweet Potatoes. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT apply more than two applications by air. • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products per year. • DO NOT apply within 14 days of harvest (14-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Root and Tuber Vegetables except Sugar beet, Crop Subgroup 1B** Carrot Beet, garden Ginseng Horseradish Parsnip Radish Radish (Oriental) Rutabaga Turnip Not for Use in California	Alternaria leaf blight <i>(Alternaria dauci)</i> Powdery mildew <i>(Erysiphe spp.)</i>	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
**Additional Root and Tuber Vegetables			
Burdock, edible, Celeriac, Chicory, Salsify (including black and Spanish), Skirret, Turnip-root parsley, and Turnip rooted chervil.			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Root and Tuber Vegetables except Sugar beet, Crop Subgroup 1B. • DO NOT apply more than 4 applications per year at the highest rate except for radish. • Minimum Application Interval: 7 days • DO NOT apply more than two applications by air. • Radish ONLY – DO NOT apply more than two applications per year. • Radish ONLY – DO NOT apply more than 14 oz. of Cyril WDG Fungicide (0.66 lb. a.i./A of cyprodinil) per crop per year. • Radish ONLY – DO NOT apply more than 0.66 lb. a.i./A of cyprodinil-containing products per year. • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year for specified leaves of root and tuber vegetables, except radish. • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products per year for specified leaves of root and tuber vegetables, except radish. • DO NOT apply within 7 days of harvest (7-day PHI). • DO NOT allow cattle or other livestock to feed upon the leaves of root vegetables. 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Berry, Low Growing Subgroup 13-07G (except Cranberry)** Strawberry Not for Use in California	Gray mold <i>(Botrytis cinerea)</i> Powdery mildew <i>(Sphaerotheca macularis)</i> Anthracnose <i>(Colletotrichum spp.)</i>	5.5 - 10*	Begin application at or before bloom and continue on a 7- to 10-day interval. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
	Root and crown anthracnose at planting <i>(Colletotrichum spp.)</i>	2.5 - 4 oz. per 100 gal. water	Apply as a preplant dip to strawberry roots and crowns at the rate of 2.5 to 4 oz. per 100 gal. of water for suppression of root and crown rot caused by anthracnose. Wash transplants to remove excess soil prior to dipping. This helps to remove adhering spores from the external plant parts. Completely immerse planting stock in dip solution. Dip or expose plants for a minimum of 2 to 5 minutes. DO NOT reuse solution. Dispose of dip solution according to local regulations. Plant treated plants as quickly as possible. For continued anthracnose control, follow with foliar applications of beginning 2 - 3 weeks after transplant.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A *10 oz. product contains 0.47 lb. a.i. cyprodinil/A			

(continued)

****Additional Low Growing Berries**

Bearberry; bilberry; cloudberry; muntries; partridgeberry and cultivars and/or hybrids of these.

Application Instructions

Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions

- **Maximum Single Application Rate: DO NOT** exceed the maximum rate listed above for Berry, Low Growing Subgroup 13-07G (except Cranberry) and Strawberry.
- **DO NOT** apply more than 2 applications (air and plant dip applications) per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** apply more than two applications by air.
- **DO NOT** apply more than one pre-plant dip application per crop.
- **DO NOT** apply more than 20 oz./A of **Cyril WDG Fungicide** (0.94 lb. a.i./A of cyprodinil) per year.
- Apply up to and on the day of harvest (0-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Fruiting Vegetable Crop Group 8-10** Eggplant Groundcherry Pepino Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) Tomatillo Tomatoes Not for Use in California	Early blight <i>(Alternaria solani)</i> Grey mold <i>(Botrytis cinerea)</i> Powdery mildew <i>(Leveillula taurica)</i>	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
**Additional Fruiting Vegetables			
African eggplant; bush tomato; cocona; currant tomato; garden huckleberry; goji berry; martynia; naranjilla; okra; pea eggplant; pepino; pepper, nonbell; roselle; scarlet eggplant; sunberry; tree tomato; cultivars, varieties, and/or hybrids of these.			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Fruiting Vegetable Crop Group 8-10. • Minimum Application Interval: 7 days • DO NOT apply more than two applications by air. • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide per year. • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products per year. • DO NOT apply more than a maximum total of 4 applications (air plus ground plus chemigation) per year. • Apply up to and on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Watercress Not for Use in California	Cercospora leaf spot <i>(Cercospora spp.)</i>	5.5 - 7*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Cyril WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*5.5 oz. product contains 0.26 lb. a.i. cyprodinil/A			
*7 oz. product contains 0.33 lb. a.i. cyprodinil/A			
Application Instructions			
Application may be made by ground or chemigation. Good coverage is essential for good disease control. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed above for Watercress. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT apply more than 28 oz./A of Cyril WDG Fungicide (1.3 lb. a.i./A of cyprodinil) per year. • Applications can be made to a dry bed only. DO NOT apply directly to water. • Apply up to and on the day of harvest (0-day PHI). 			

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container, when not in use. Store in a cool, dry area that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Pesticide wastes may be toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

CONTAINER HANDLING:

Less than or equal to 50 pounds:

Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

WARRANTY AND DISCLAIMER STATEMENT

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury is not expected when applying end-use products, according to label directions and restrictions, that are formulated using this manufacturing use only product. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA CROP CHEM LTD. MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Sharda Cropchem Ltd. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA CROP CHEM LTD. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SHARDA CROP CHEM LTD.'S ELECTION, THE REPLACEMENT OF PRODUCT.

NOTES

CYPRDINIL GROUP 9 FUNGICIDE

Cyril WDG Fungicide

ACTIVE INGREDIENT:	WT. BY %
Cyprodinil: 4-cyclopropyl-6-methyl-N-phenylpyrimidin-2-amine	75.0%
OTHER INGREDIENTS:	25.0%
TOTAL:	100.0%

Cyril WDG Fungicide is a water-dispersible granule containing 75% cyprodinil.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID - IF ON SKIN OR CLOTHING: • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice. **IF IN EYES:** • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER - Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at **1-800-222-1222**. For information about this product, contact the National Pesticides Information Center (NPIC) at **1-800-858-7378**, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>. **NOTE TO PHYSICIAN** - If ingested, induce emesis or lavage stomach. Treat symptomatically.

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION - Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. **ENVIRONMENTAL HAZARDS** - This pesticide is toxic to fish and aquatic invertebrates. For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment wash-water or rinsate. **Surface and Ground Water Advisory** - This chemical may contaminate water through runoff. This product may impact surface water quality due to runoff of rain

water. This is especially true for poorly draining soils and soils with shallow ground water. This chemical has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this chemical. A level, well maintained vegetative buffer strip between areas to which this chemical is applied and surface water features including ponds, streams, and springs will reduce the potential loading of cyprodinil from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff. **Physical or Chemical Hazards - DO NOT** use, pour, spill or store near heat or open flame. **DIRECTIONS FOR USE** - It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL - DO NOT contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE:** Keep this product in its tightly closed original container, when not in use. Store in a cool, dry area that is inaccessible to children and animals. **PESTICIDE DISPOSAL:** Pesticide wastes may be toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance. **CONTAINER HANDLING:** Less than or equal to 50 pounds: Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

See label booklet for additional Precautionary Statements and Directions For Use.

Manufactured For: Sharda US LLC,
7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707
EPA Reg. No. 82633-132-83529 EPA Est. No. **AG** 72159-GA-001;
(MA) 83411-MN-001; **(SC)** 39578-TX-001; **(TX)** 07401-TX-001
The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: **50 oz.**

OPEN HERE