

CYPRODINIL	GROUP	9	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

Button WDG Fungicide

ACTIVE INGREDIENTS:

Cyprodinil: 4-cyclopropyl-6-methyl-N-phenylpyrimidin-2-amine.	WT. BY %
Fludioxonil: 4-(2,2-difluoro-1,3-benzodioxol-4-yl)-1H-pyrrole-3-carbonitrile	37.5%
	25.0%

OTHER INGREDIENTS: 37.5%

TOTAL: **100.0%**

Button WDG Fungicide is a water-dispersible granule containing 37.5% cyprodinil and 25% fludioxonil.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

**See label booklet for First Aid, Precautionary Statements,
Directions For Use, and Storage and Disposal.**

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 82633-133-83529

EPA Est. No. **AG** 72159-GA-001; **GH** 70815-GA-002;

MA 83411-MN-001; **TX** 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: 0.75 lb.

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222. For information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Handlers applying this product as a pre-plant dip to strawberry roots and crowns and workers packaging or preparing treated roots and crowns for shipment must wear:

- Chemical-resistant apron made of any waterproof material
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or Viton \geq 14 mils
- Chemical-resistant boots made of any waterproof material

All other applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or Viton \geq 14 mils
- Shoes plus socks

In addition, mixers and loaders for aerial, ground boom, and chemigation applications must wear:

- A minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter (e.g., N95, R95 or P95); OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air-purifying respirator with an HE filter.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. Aerial applicators must be in enclosed cockpits.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This chemical may contaminate water through drift of spray in wind. This chemical has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this chemical. A level, well maintained vegetative buffer strip between areas to which this chemical is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this chemical will be reduced by avoiding applications when conditions favor runoff (for example when soils are saturated and/or significant rainfall is forecast in the next 48 hours). Sound erosion control practices will reduce this chemical's contribution to surface water contamination.

Physical or Chemical Hazards

DO NOT use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE); notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

PRODUCT INFORMATION

Buttun WDG Fungicide is a broad-spectrum fungicide for the control of certain diseases.

PRODUCT USE RESTRICTIONS

DO NOT plant any crop which is not registered for use with cyprodinil or fludioxonil for a period of 30 days unless a shorter interval is specified on the following list.

Rotational Crop	Planting Time from Last Buton WDG Fungicide Application
Beans (dried and succulent except cowpeas)* Berries (bushberries 13-07B, caneberries 13-07A)* Brassica, Head and Stem (Crop Group 5-16)* Brassica, Leafy greens (Crop Subgroup 4-16B)* Celtnce Citrus Fruit (Crop Group 10-10B) Cucurbits Crop Group 9* Fennel, Florence, fresh leaves, and stalk Herbs (fresh and dried)* Kohlrabi Leafy Greens (Crop Subgroup 4-16A) Leaf petiole vegetables (Crop Subgroup 22B) Leaves of Root and Tuber Vegetables* Onion, Bulb, Crop Subgroup 3-07A Onion, Green, Crop Subgroup 3-07B Fruiting vegetables crop group 8-10 Tuberous and Corm Vegetables (crop subgroup 1C)* Root and Tuber Vegetables except Sugar beet (Crop Subgroup 1B)* Strawberries Tomatoes Watercress Crops Not Intended for Food or Feed	0 days
All Other Crops Intended for Food or Feed	30 days

*See crop lists in **CROP USE DIRECTIONS** section.

In annual crops where multiple crops can be grown per year (double/triple cropping), **DO NOT** apply more than 1.3 lb. a.i. cyprodinil and 0.9 lb. a.i. fludioxonil per acre per year to an individual plot of land.

For the crops to which aerial applications are allowed, refer to the specific crop directions for use. Aerial applicators must be in enclosed cockpits.

Nassau and Suffolk counties of New York: use limited to strawberries and onions.

RESISTANCE MANAGEMENT

CYPRODINIL	GROUP	9	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

For resistance management, please note that **Button WDG Fungicide** contains both a Group 9/cyprodinil and Group 12/fludioxonil fungicide. Cyprodinil is classified in the Group 9 chemical class as an anilinopyrimidine. Fludioxonil is classified in the Group 12 chemical class as phenylpyrrole class of chemistry and has a unique mode of action which prevents fungal respiration. Any fungal population may contain individuals naturally resistant to **Button WDG Fungicide** and other Group 9 and Group 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Button WDG Fungicide** or other Group 9 and Group 12 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For information or to report suspected resistance, contact your local Sharda USA LLC representative.

APPLICATION INSTRUCTIONS

Thorough coverage is necessary to provide good disease control. Applications using sufficient water volume to provide thorough and uniform coverage provide the most effective disease control. Use minimum ground spray volumes of 10 gal./A for field and vegetable crops and 50 gal./A for tree crops. For aerial application, see directions in the specific crop directions for use.

To avoid spray drift, **DO NOT** apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Calibrate sprayer before use.

Use a pump with capacity to maintain the correct rated pressure for the nozzles selected. Maintain sufficient agitation to keep the mixture in suspension. Use a jet agitator, liquid sparge tube, or mechanical paddle for agitation. **DO NOT** air sparge.

Use screens to prevent nozzles from clogging. Use 50-mesh or coarser screens placed after the tank and before the nozzles. Check nozzle manufacturers' specifications.

For more information on spray equipment and calibration, consult sprayer manufacturers' and state directions. For specific local directions and spray schedules, consult the current state agricultural experiment station.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH FARM PONDS.

- **DO NOT** apply within 75 ft. of bodies of water including lakes, reservoirs, rivers, permanent streams, natural ponds, marshes, or estuaries.
- Shut off the sprayer when at row ends.
- **DO NOT** cultivate within 10 ft. of aquatic areas as to allow a vegetative filter strip.
- **DO NOT** apply when weather conditions favor drift to aquatic areas. **DO NOT** apply when gusts or sustained winds exceed 15 mph.
- **DO NOT** apply during a temperature inversion. Mist or fog may indicate the presence of an inversion in humid areas.
- For perennial crops including tree crops and grapes:
- For all plantings within 150 ft. of bodies of water as described above, spray crops from outside the planting away from the bodies of water.
- Spray last three rows windward of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas. Adjust or turn off top nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row or passing tree gaps in the rows.

Ground Application

- Apply in a minimum of 10 gallons of water per acre, unless specified otherwise.

Ground Spray Drift Restrictions

- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.

Aerial Spray Directions

Avoid applications under conditions when uniform coverage cannot be obtained or when excessive drift may occur.

Aerial Spray Drift Restrictions

Observe the following restrictions when spraying in the vicinity of aquatic area including lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.

- Use only on crops where aerial applications are indicated.
- **DO NOT** apply by air within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Release spray at the lowest height consistent with pest control and flight safety. **DO NOT** make applications more than 10 feet above the crop canopy.
- **DO NOT** apply when weather conditions favor drift to aquatic areas.
- **DO NOT** apply during a temperature inversion. Mist or fog may indicate the presence of an inversion in humid areas.

Aerial Spray Precautions

Observe the following precautions when spraying in the vicinity of aquatic area including lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.

- Use the largest droplet size consistent with good pest control.
- Formation of very small droplets may be minimized by appropriate nozzle selection, by orientating nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Reduce risk of exposure to aquatic areas by avoiding applications when wind direction is toward the aquatic area.
- Low humidity and high temperatures increase the evaporation rate of spray droplets, and therefore the likelihood of increased spray drift to aquatic area. Avoid spraying during conditions of low humidity and/or high temperatures.
- For the crops to which aerial applications are allowed, refer to the specific crop directions for use.
- Apply in a minimum of 5 gallons of water per acre, unless specified otherwise.

SPRAY DRIFT MANAGEMENT

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications

- **DO NOT** release spray at a height greater than 10 ft. above the vegetative canopy unless a greater application height is necessary for pilot safety.
- For all applicators, applicators are required to use a medium or coarse spray droplets size (ASABE S572.1).
- For aerial applications: **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind-speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply during temperature inversions.

Ground boom Applications

- Apply with the nozzle height directed by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

- **Pressure** - Use the lowest spray pressure directed for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Boom Height - Ground Boom

- For ground equipment, the boom must remain level with the crop and have minimal bounce.

Controlling Droplet Size - Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturer's directions for setting up nozzles. To reduce fine droplets, orient nozzles parallel with the airflow in flight.

Release Height - Aircraft

- Higher release heights increase the potential for spray drift.

Shielded Sprayers

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature And Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential increases with speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

- Take precautions to minimize spray drift.

Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through drip, microjet, center pivot, solid set, hand move, and moving wheel irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.125 - 0.25 inches/A of water. Excessive water may reduce efficacy.
- If you have questions about calibration, contact state extension service specialists, equipment manufacturers, or other experts.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, for example a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Center Pivot Irrigation Equipment

Restrictions: (1) Use only with drive systems which provide uniform water distribution.

(2) **DO NOT** use end guns when chemigating **Button WDG Fungicide** through center pivot systems because of non-uniform application.

- Determine the size of the area to be treated.
- Determine the time required to apply 1/8 - 1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as directed by the equipment manufacturer. When applying **Button WDG Fungicide** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80 - 95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of **Button WDG Fungicide** required to treat the area covered by the irrigation system.
- Add the required amount of **Button WDG Fungicide** and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the **Button WDG Fungicide** solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the **Button WDG Fungicide** solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20- to 30-minute interval. When applying **Button WDG Fungicide** through irrigation equipment, use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of **Button WDG Fungicide** required to treat the area covered by the irrigation system.
- Add the required amount of **Button WDG Fungicide** into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the **Button WDG Fungicide** solution has cleared the last sprinkler head.

Drip or Microjet Chemigation Systems

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

Spray Preparation: Chemical tank and injector system must be thoroughly cleaned. Flush system with clean water.

Use Directions for Drip or Microjet Irrigation Applications

Drip or Microjet Irrigation: Butten WDG Fungicide may be applied through drip irrigation systems for soil-borne disease control. The soil must have adequate moisture capacity prior to drip application.

- Terminate drip irrigation at fungicide depletion from the main feed supply tank or after 6 hours from start, whichever is shorter. For maximum efficacy, delay subsequent irrigation (water only) for at least for 24 hours following drip application.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, for example a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.
- If you have questions about calibration, contact state extension service specialists, equipment manufacturers or other experts.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

SPECIFIC INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, for example a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

MIXING PROCEDURES

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. **DO NOT** let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

Button WDG Fungicide Alone: Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the **Button WDG Fungicide** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the Switch 62.5WG has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Button WDG Fungicide + Tank Mixtures: **Button WDG Fungicide** is compatible in tank mixtures with many commonly used fungicides, liquid fertilizers, herbicides, insecticides, and biological control products. If tank mixes are desired, observe all directions, precautions, and limitations on labeling of all products used. Consult compatibility charts or your local or state agricultural authorities for compatibility information.

To prepare spray solution, add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) including **Button WDG Fungicide**, liquid flowables, liquids, and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

When using **Button WDG Fungicide** in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including **Button WDG Fungicide**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using **Button WDG Fungicide** in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. **DO NOT** exceed label dosage rates, and follow the most restrictive label precautions and limitations. This product must not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CROP USE DIRECTIONS

When a range of rates is provided, use the higher specified or labeled rates if weather conditions are conducive for higher disease pressure.

Crop	Disease	Product Rate oz./Acre	Use Directions
Beans (Dried and Succulent except cowpeas) Chickpea (garbanzo bean) Bean (<i>Lupinus</i> spp.) (grain lupin, sweet lupin, white lupin, white sweet lupin) Bean (<i>Phaseolus</i> spp.) (kidney, lima, mung, navy, pinto, snap, wax) Broad Bean (fava bean) Bean (<i>Vigna</i> spp.) (asparagus, blackeyed pea)	White mold (<i>Sclerotinia sclerotiorum</i>) Gray mold (<i>Botrytis cinerea</i>)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7-day interval if conditions remain favorable for disease development. For White Mold control, make the first application at 10 - 20% bloom. In some locations a single application at this timing will pro- vide adequate disease control. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

(continued)

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed in the table above for beans.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** make more than two applications by air.
- **DO NOT** apply more than 56 oz./A of **Button WDG Fungicide** per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil).
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Berries Bushberry Subgroup 13-07B* Blueberry Currant Caneberry Subgroup 13-07A** Blackberry Red and Black Raspberry And cultivars and/or hybrids of these.	Mummy berry <i>(Monilinia</i> <i>vaccinilicorymbosi)</i> Anthracnose <i>(Colletotrichum spp.)</i> Alternaria fruit rot <i>(Alternaria tenuissima)</i> Phomopsis <i>(Phomopsis vaccinii)</i> Botrytis fruit rot <i>(Botrytis cinerea)</i>	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day in- terval if conditions remain favor- able for disease development. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
Complete List of Bushberries and Caneberries			
*Bushberries: Aronia berry, Black currant, Blueberry high and low bush, Buffalo currant, Chilean guava, Edible honeysuckle, Elderberry, European barberry, Gooseberry, Highbush cranberry, Huckleberry, Jostaberry, Juneberry (Saskatoon berry), Lingonberry, Native currant, Red currant, Salal, Sea buckthorn **Caneberries: Blackberry, Loganberry, Red and Black Raspberry, Wild raspberry			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for bushberries and caneberries. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT make more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • May be applied on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Brassica Head and Stem Vegetable Crop group 5-16* Broccoli Brussels sprouts Cabbage Cauliflower And cultivars and/or hybrids of these. [Not for use in California]	Powdery mildew <i>(Erysiphe polygoni)</i>	10 - 12 (0.23 - 0.28 lb. cyprodinil/Acre and 0.16 - 0.19 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day in- terval if conditions remain favor- able for disease development.
	Alternaria leaf blight <i>(Alternaria spp.)</i> Suppression: Cercospora leaf spot <i>(Cercospora brassicicola)</i>	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*Complete List of Brassica Head and Stem Vegetables Crop group 5-16			
Broccoli; Brussels sprouts; Cabbage; Cabbage, Chinese (napa); Cauliflower; cultivars, varieties, and hybrids of these commodities			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for Brassica Head and Stem Vegetables Crop group 5-16. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT make more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Brassica Leafy Greens Subgroup 4-16B* Arugula Chinese cabbage Bok Choy Collards Kale Mustard greens Turnip greens And cultivars and/or hybrids of these. See separate instructions for watercress. [Not for use in California]	Powdery mildew <i>(Erysiphe polygoni)</i>	10 - 12 (0.23 - 0.28 lb. cyprodinil/Acre and 0.16 - 0.19 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development.
	Alternaria leaf blight <i>(Alternaria spp.)</i> Suppression: Cercospora leaf spot <i>(Cercospora brassicicola)</i>	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*Complete list of Brassica Leafy Greens Vegetable subgroup 4-16B			
Arugula; broccoli, Chinese; broccoli raab; cabbage, abyssinian; cabbage, seakale; cabbage, Chinese, bok choy; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; cultivars, varieties, and hybrids of these commodities.			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

(continued)

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed in the table above for Brassica Leafy Greens Vegetable subgroup 4-16B.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- Make no more than two applications by air.
- **DO NOT** use roots of treated turnips for food or feed. Only turnip varieties harvested for their leaves may be treated.
- **DO NOT** apply more than 56 oz./A of **Button WDG Fungicide** per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil).
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Citrus, Crop Subgroup 10-10B* Lemon Lime	Alternaria stem end rot (<i>A. citri</i>) Anthracnose (<i>Colletotrichum gloeosporioides</i>) Blue mold (<i>Penicillium italicum</i>) Green mold (<i>Penicillium digitatum</i>)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Make one application near harvest to prevent post-harvest fruit rot. The application may be made up to and including the day of harvest.
*Complete List of Citrus Crop Group 10-10B			
Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; kumquat; lemon; lime; mount white lime; New Guinea wild lime; Russell River lime; sweet lime; Tahiti lime; cultivars, varieties, and/or hybrids of these.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for Citrus Crop Group 10-10B. • DO NOT apply more than 1 application per year at the highest rate. • Application may be made by ground only. • DO NOT apply more than 14 oz./A of Buttton WDG Fungicide per year (0.33 lb. cyprodinil and 0.22 lb. fludioxonil). • DO NOT apply more than 0.33 lb. a.i./A of cyprodinil-containing products and 0.22 lb. a.i./A of fludioxonil-containing products per year. • May be applied on the day of harvest (0-day PHI). • DO NOT exceed one application per year. 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Cucurbits, Crop Group 9* Cantaloupe Cucumber Honeydew Muskmelon Watermelon Pumpkin Squash Zucchini And cultivars and/or hybrids of these.	Alternaria leaf blight (<i>A. cucumerina</i>) Alternaria leaf spot (<i>A. alternata</i>) Gummy stem blight (<i>Didymella bryoniae</i>) Powdery mildew (<i>Sphaerotheca fuliginea</i> , <i>Erysiphe cichoracearum</i>)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*Complete List of Cucurbit vegetable Crop Group 9:			
Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter-melon, Chinese cucumber); muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon); pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon.			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

(continued)

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed in the table above for Cucurbit vegetable Crop Group 9.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** make more than two applications by air.
- **DO NOT** apply more than 56 oz./A of **Button WDG Fungicide** per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil).
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year.
- May be applied up to 1 day before harvest (1-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Grapes and Small Fruit Vine Climbing Subgroup 13-07F (except fuzzy kiwifruit) Grapes Amur river grape Hardy kiwifruit Maypop Schisandra berry And cultivars and/or hybrids of these.	Botrytis (grey mold) (<i>B. cinerea</i>) Sour Rot (caused by a fungal complex)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	<p>Begin applications of Button WDG Fungicide at early bloom. Continue applications on a 21-day interval based on disease pressure. Up to three additional applications may be made at berry touch, veraison, or preharvest. Botrytis Bunch Rot is most effectively controlled by ground application, using sufficient water volume to provide thorough coverage. Thorough coverage of bunches is essential.</p> <p>For sour rot, make an application at veraison followed by 1 - 2 additional applications.</p> <p>Resistance Management: After 2 applications of Button WDG Fungicide, alternate with another fungicide with a different mode of action for 2 applications.</p>
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for Grapes and Small Fruit Vine Climbing Subgroup 13-07F (except fuzzy kiwifruit). • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 21 days • DO NOT make more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.4 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • DO NOT apply within 7 days of harvest (7-day PHI). • Minimum Re-treatment Interval: 21 days 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Herbs (Dried and fresh)*	Alternaria leaf spot (<i>Alternaria</i> spp.) Botrytis leaf blight (<i>Botrytis</i> spp.) Fusarium blight (<i>Fusarium</i> spp.)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Apply in a minimum spray volume of 30 gal./A to obtain thorough coverage. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*Complete List of Dried and Fresh Herbs			
Angelica, Balm, Basil, Borage, Burnet, Chamomile, Catnip, Chervil, dried leaves, Chives, Clary, Coriander, leaves (cilantro), Costmary, Culantro, leaves, Curry, leaves, Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage, leaves, Marigold, Marjoram, Nasturtium, Parsley, dried leaves, Pennyroyal, Rosemary, Rue, Sage, Savory, summer and winter, Sweet bay, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for dried and fresh herbs. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT make more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Kohlrabi [Not for use in California]	Powdery mildew (<i>Erysiphe polygoni</i>)	10 - 12 (0.23 - 0.28 lb. cyprodinil/Acre and 0.16 - 0.19 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development.
	Alternaria leaf blight (<i>Alternaria</i> spp.) Suppression: Cercospora leaf spot (<i>Cercospora brassicicola</i>)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for kohlrabi. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT make more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Leafy Greens, Crop Subgroup 4-16A*, except Parsley Leaf petiole vegetables Crop Subgroup 22B** Celtuce Fennel, Florence, fresh leaves, and stalk Celery Lettuce, head, and leaf Spinach And cultivars and/or hybrids of these. [Not for use in California]	Alternaria leaf spot (<i>Alternaria</i> spp.) Septoria leaf spot (<i>Septoria lactucae</i>) Gray mold (<i>Botrytis cinerea</i>) Sclerotinia rot (<i>Sclerotinia</i> spp.) Basal rot (<i>Phoma exigua</i>) Suppression: Powdery mildew (<i>Erysiphe cichoracearum</i>)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. For control of Sclerotinia, make the first application at thinning and again two weeks later. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*Complete List of Leafy Greens Crop subgroup 4-16A			
Amaranth, Chinese; amaranth, leafy; aster, Indian; blackjack; cat's whiskers; cham-chwi; cham-na-mul; chervil, fresh leaves; chipilin; chrysanthemum, garland; cilantro, fresh leaves; corn salad; cosmos; dandelion, leaves; dang-gwi, leaves; dillweed; dock; dol-nam-mul; ebolo; endive; escarole; farnflower; feather cockscomb; good king henry; huauzontle; jute, leaves; lettuce, bitter; lettuce, head; lettuce, leaf; orach; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; spinach; spinach, Malabar; spinach, New Zealand; spinach, tanier; Swiss chard; violet, Chinese, leaves; cultivars, varieties, and hybrids of these commodities			
**Leaf petiole vegetables Crop subgroup 22B Includes			
Cardoon; celery; celery, Chinese; fuki; rhubarb; udo; zuiki; cultivars, varieties, and hybrids of these commodities			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

(continued)

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed in the table above for Leafy Greens, Crop Subgroup 4-16A, except Parsley and Leaf petiole vegetables Crop Subgroup 22B.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** make more than two applications by air.
- **DO NOT** apply more than 56 oz./A of **Button WDG Fungicide** per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil).
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year.
- May be applied on the day of harvest (0-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Leaves of Root and Tuber Vegetables Crop Group 2* Beet, garden Beet, sugar Carrot Parsnip Radish Sweet Potato Turnip Yam (true)	Alternaria leaf blight <i>(Alternaria dauci)</i> Powdery mildew <i>(Erysiphe spp.)</i>	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*Complete List of Root and Tuber Vegetables, Leaves			
Beet, garden; Beet, sugar; Burdock, edible; Carrot; Cassava; Celeriac; Chicory; Dasheen; Parsnip; Radish; Radish (oriental); Rutabaga; Salsify (including black and Spanish); Sweet potato; Tanier; Turnip; Turnip-rooted chervil; Yam (true)			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

(continued)

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed in the table above for Leaves of Root and Tuber Vegetables Crop Group 2.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** make more than two applications by air.
- Radish ONLY - **DO NOT** make more than two applications per year.
- Radish ONLY - **DO NOT** apply more than 28 oz./A of **Button WDG Fungicide** per year (0.66 lb. cyprodinil and 0.44 lb. fludioxonil).
- Radish ONLY - **DO NOT** apply more than 0.66 lb. a.i./A of cyprodinil-containing products and 0.44 lb. a.i./A of fludioxonil-containing products per year.
- **DO NOT** apply more than 56 oz./A of **Button WDG Fungicide** per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil).
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI).
- **DO NOT** allow cattle or other livestock to feed upon the leaves of root and tuber vegetables.

Crop	Disease	Product Rate oz./Acre	Use Directions
Onions and Garlic Bulb Vegetables Crop Group 3-07A and 3-07B* Garlic Onion, bulb Onion, green Onions grown for seed And cultivars and/or hybrids of these.	Botrytis leaf blight or blast <i>(Botrytis spp.)</i> Stemphylium leaf blight <i>(Stemphylium vesicarium)</i> Purple blotch <i>(Alternaria porri)</i> Suppression: Neck rot <i>(Botrytis spp.)</i> Black mold <i>(Aspergillus niger)</i>	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. For optimal effect on neck rot, apply on a 7-day schedule at the 14 oz. rate. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
	Soilborne diseases White rot <i>(Sclerotium cepivorum)</i>	7 - 14 (0.16 - 0.33 lb. cyprodinil/Acre and 0.11 - 0.22 lb. Fludioxonil/Acre)	Apply at the time of planting as an in-furrow spray.

*Complete List of Bulb Vegetables

Bulb Onion: Chinese onion; Dry Bulb onion; Daylily bulb; Fritillaria bulb; Garlic; Great-headed garlic; Lily bulb; Pearl onion; Potato onion; Serpent garlic; Shallot;

Green Onion: Beltville bunching onion; Chinese chive fresh leaves; Fresh chive leaves; Fritillaria leaves; Fresh onion; Green onion; Hosta elegans; Kurrat; Lady's leek; Leek; Macrostem onion; Shallot fresh leaves; Tree tops onion; Welsh onion tops; Wild leek

Application Instructions

Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

(continued)

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed in the table above for Onions and Garlic Bulb Vegetables Crop Group 3-07A and 3-07B.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** make more than two applications by air.
- **DO NOT** apply more than 56 oz./A of **Button WDG Fungicide** per year.
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 1.0 lb. a.i./A of fludioxonil-containing products per year.
- For in-furrow applications, **DO NOT** apply more than the maximum single application rate listed in the table.
- **DO NOT** apply within 7 days of harvest (7-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Pistachio	Botrytis (<i>Botrytis</i> spp.) Alternaria (<i>Alternaria alternata</i>)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Make the first application during early bloom and repeat applications at 14-day intervals if conditions remain favorable for disease development. Resistance Management: After 2 applications, alternate with another fungicide with a different mode of action for 2 applications.
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for pistachio. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 14 days • DO NOT make more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • DO NOT apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Potatoes Tuberous and Corm Vegetables Crop Subgroup 1C* Sweet Potatoes [Not for use in California]	Brown spot <i>(Alternaria alternata)</i> Early blight <i>(A. solani)</i> Powdery mildew <i>(Erysiphe</i> <i>cichoracearum)</i> Septoria leaf spot <i>(Septoria lycopersici)</i> Tan spot <i>(Botrytis cinerea)</i>	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 appli- cations of Button WDG Fungicide , alter- nate with another fungicide with a different mode of action for 2 applications.
*Complete List of Vegetables, tuberous and corm subgroup 1C			
Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna, Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen (Taro), Ginger, Leren, Tanier, Turmeric, Yam (bean and true), and cultivars and/or hybrids of these			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for Potatoes Tuberous and Corm Vegetables Crop Subgroup 1C and Sweet Potatoes. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT make more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • DO NOT apply within 14 days of harvest (14-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Root Vegetables Subgroup 1B (except Sugar beet)* Carrot Beet, garden Ginseng Horseradish Parsnip Radish Radish (oriental) Rutabaga Turnip	Alternaria leaf blight <i>(Alternaria dauci)</i> Powdery mildew <i>(Erysiphe spp.)</i>	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*Additional Root Vegetables Subgroup 1B			
Burdock, edible, Celeriac, Chicory, Salsify (including black and Spanish), Skirret, Turnip-root parsley, and Turnip-rooted chervil			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

(continued)

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed in the table above for Root Vegetables Subgroup 1B (except Sugar beet).
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** make more than two applications by air.
- **Radish ONLY** - Make no more than two applications per year.
- **Radish ONLY** - **DO NOT** apply more than 28 oz./A of **Button WDG Fungicide** per year (0.66 lb. cyprodinil and 0.44 lb. fludioxonil).
- **Radish ONLY** - **DO NOT** apply more than 0.66 lb. a.i./A of cyprodinil-containing products and 0.44 lb. a.i./A of fludioxonil-containing products per year.
- **DO NOT** apply more than 56 oz./A of **Button WDG Fungicide** per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil).
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI).
- **DO NOT** allow cattle or other livestock to feed upon the leaves of root and tuber vegetables.

Crop	Disease	Product Rate oz./Acre	Use Directions
Strawberry and Berry, Low Growing Subgroup 13-07G (except Cranberry)* And cultivars and/or hybrids of these.	Gray mold (<i>Botrytis cinerea</i>) Powdery mildew (<i>Sphaerotheca macularis</i>) Anthracnose (<i>Colletotrichum</i> spp.)	11 - 14 (0.26 - 0.33 lb. cymodini/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin application at or before bloom and continue on a 7- to 10-day interval. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
	Root and crown anthracnose at planting (<i>Colletotrichum</i> spp.)	5 - 8 oz. per 100 gal. water (0.12 - 0.19 lb. cymodini/100 gal. water and 0.08 - 0.12 lb. Fludioxonil/100 gal. water)	Apply as a preplant dip to strawberry roots and crowns at the rate of 5 to 8 oz. per 100 gallons of water for suppression of root and crown rot caused by anthracnose. Wash transplants to remove excess soil prior to dipping. Completely immerse planting stock in dip solution. Dip or expose plants for a minimum of 2 minutes or a maximum of 5 minutes. Completely drain the transplants after dip. DO NOT reuse solution. Dispose of dip solution according to local regulations. Plant treated plants as quickly as possible. For continued anthracnose control, follow with foliar applications of beginning 2 - 3 weeks after transplant.
*Complete List of Low Growing Berries			
Bearberry; bilberry; cloudberry; muntries; partridgeberry; Strawberry and cultivars and/or hybrids of these			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

(continued)

Specific Use Restrictions

- **Maximum Single Application Rate:** **DO NOT** exceed the maximum rate listed in the table above for Strawberry and Berry, Low Growing Subgroup 13-07G (except Cranberry) and cultivars and/or hybrids of these.
- **DO NOT** apply more than 4 applications per year at the highest rate.
- **Minimum Application Interval:** 7 days
- **DO NOT** make more than two applications by air.
- Make only one pre-plant dip application per crop.
- **DO NOT** apply more than 56 oz./A of **Button WDG Fungicide** per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil).
- **DO NOT** apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year.
- May be applied on the day of harvest (0-day PHI).

Crop	Disease	Product Rate oz./Acre	Use Directions
Sugar Apple [Not for use in California]	Anthrachnose (<i>Colletotrichum</i> spp.) Botrytis fruit rot (<i>Botrytis</i> spp.) Alternaria fruit rot (<i>Alternaria</i> spp.) Leaf and fruit spot (<i>Alternaria</i> spp.)	14 (0.33 lb. cyprodinil/ Acre and 0.22 lb. Fludioxonil/Acre)	Begin applications prior to disease development. Continue applications throughout the season on a 7-day interval, following resistance management guidelines. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
Application Instructions			
Apply specified rate per acre as a directed foliar spray. Apply by ground using a minimum of 75 gallons/A spray volume. An adjuvant may be added at specified rates.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • DO NOT apply via aerial application. • DO NOT apply via irrigation, including chemigation. • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for sugar apple. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.32 lb. a.i./A of cyprodinil-containing products and 0.88 lb. a.i./A of fludioxonil-containing products per year. • DO NOT use an organo-silicone adjuvant. • May be applied on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Tomatoes and Fruiting Vegetable Crop Group 8-10* Eggplant Okra Pepper, bell Pepper, nonbell	Early blight (<i>Alternaria solani</i>) Grey mold (<i>Botrytis cinerea</i>) Powdery mildew (<i>Leveillula taurica</i>) Target spot ¹ (<i>Corynespora cassiicola</i>) ¹ [Not for use in California]	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*Complete List of Fruiting Vegetable Crop Group 8-10			
African eggplant; Bush tomato; Cocona; Currant tomato; Eggplant; Garden huckleberry; Goji berry; Groundcherry; Martynia; Naranjilla; Okra; Pea eggplant; Pepino; Pepper, bell; Pepper, nonbell; Roselle; Scarlet eggplant; Sunberry; Tomatillos; Tomato; Tree tomato and cultivars and/or hybrids of these			
Application Instructions			
Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for Tomatoes and Fruiting Vegetable Crop Group 8-10. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT make more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • DO NOT apply more than a maximum total of 4 applications (air plus ground plus chemigation) per year. • May be applied on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Tropical and Subtropical, Small fruit, inedible peel subgroup 24A* Lychee Longan Spanish lime And cultivars and/or hybrids of these [Not for use in California]	Botrytis fruit rot <i>(Botrytis spp.)</i> Alternaria fruit rot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i> Bipolaris bud and fruit rot <i>(Bipolaris spp.)</i>	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Make the first application during early bloom and repeat on 7- to 10-day intervals if conditions re- main favorable for disease development. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*Complete List of Subgroup 24A Crops			
Aisen; bael fruit; Burmese grape; cat's-eyes; inga; longan; Lychee; madras-thorn; manduro; matisia; mesquite; mongongo, fruit; pawpaw, small-flower; satinleaf; Sierra Leone-tamarind; Spanish lime; velvet tamarind; wampi; white star apple; cultivars, varieties, and hybrids of these commodities.			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for Tropical and Subtropical, Small fruit, inedible peel subgroup 24A. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • Make no more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • DO NOT apply more than a maximum total of 4 applications (air plus ground) per year. • May be applied on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Specific Tropical Fruits* Avocado Dragon Fruit Guava Mamey sapote Mango Papaya Passionfruit Starfruit	Botrytis fruit rot (<i>Botrytis</i> spp.) Alternaria fruit rot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Bipolaris bud and fruit rot (<i>Bipolaris</i> spp.)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Make the first application during early bloom and repeat on 7- to 10-day intervals if conditions remain favorable for disease development. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
*List of Specific Tropical Fruits			
Acerola; Avocado; Black Sapote; Canistel; Dragon Fruit; Feijoa; Guava; Jaboticaba; Mamey Sapote; Mango; Papaya; Passionfruit; Pulasan; Rambutan; Sapodilla; Star apple; Starfruit; Wax Jambu			
Application Instructions			
Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for specific tropical fruits. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • DO NOT make more than two applications by air. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • DO NOT apply more than a maximum total of 4 applications (air plus ground) per year. • May be applied on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz./Acre	Use Directions
Watercress [Not for use in California]	Cercospora leafspot (<i>Cercospora</i> spp.) Sclerotinia white mold (<i>Sclerotinia</i> spp.) Rhizoctonia rot (<i>Rhizoctonia solani</i>)	11 - 14 (0.26 - 0.33 lb. cyprodinil/Acre and 0.17 - 0.22 lb. Fludioxonil/Acre)	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Button WDG Fungicide , alternate with another fungicide with a different mode of action for 2 applications.
Application Instructions			
Applications may be made by ground or chemigation. Good coverage is essential for good disease control. For chemigation apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions			
<ul style="list-style-type: none"> • Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above for watercress. • DO NOT apply more than 4 applications per year at the highest rate. • Minimum Application Interval: 7 days • Applications can be made to a dry bed only. No direct applications to water. • DO NOT apply more than 56 oz./A of Button WDG Fungicide per year (1.3 lb. cyprodinil and 0.9 lb. fludioxonil). • DO NOT apply more than 1.3 lb. a.i./A of cyprodinil-containing products and 0.9 lb. a.i./A of fludioxonil-containing products per year. • May be applied on the day of harvest (0-day PHI). 			

CROP USE DIRECTIONS

FOR POST-HARVEST APPLICATIONS

Pomegranates

Use **Button WDG Fungicide** as a post-harvest dip for the control of Botrytis fruit rot and Gray mold in pomegranates.

Application Method	Disease	Rate (oz.)	Use Directions
In-Line Dip/Drench	Botrytis fruit rot Gray mold	19.2 oz./100 gal. (0.45 lb. cyprodinil/100 gal. and 0.30 lb. Fludioxonil/100 gal.)	Mix 19.2 oz. of Button WDG Fungicide in 100 gal. of water, wax/emulsion, or aqueous dilution of wax/oil emulsion. Dip for approximately 30 seconds and allow fruit to drain.
Application Instructions			
For maximum decay control, treat fruit once before storage and once after storage, just prior to marketing. Ensure the Button WDG Fungicide solution remains in suspension by using agitation.			
Specific Use Restrictions			
DO NOT make more than two post-harvest applications of fludioxonil-containing products to the fruit.			

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container, when not in use. Store in a cool, dry area that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

CONTAINER HANDLING:

Less than or equal to 50 pounds:

Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

WARRANTY AND DISCLAIMER STATEMENT

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury is not expected when applying end-use products, according to label directions and restrictions, that are formulated using this manufacturing use only product. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Sharda USA LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SHARDA USA LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

NOTES

CYPRODINIL	GROUP 9	FUNGICIDE
FLUDIOXONIL	GROUP 12	FUNGICIDE

Button WDG Fungicide

ACTIVE INGREDIENTS:

Cyprodinil: 4-cyclopropyl-6-methyl-N-phenylpyrimidin-2-amine	WT. BY %
Fludioxonil: 4-(2,2-difluoro-1,3-benzodioxol-4-yl)-1H-pyrole-3-carbonitrile	37.5%
	25.0%

OTHER INGREDIENTS: 37.5%

TOTAL: 100.0%

Button WDG Fungicide is a water-dispersible granule containing 37.5% cyprodinil and 25% fludioxonil.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID - IF ON SKIN OR CLOTHING: • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice. **IF IN EYES:** • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOTLINE NUMBERS - Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at **1-800-222-1222**. For information about this product, contact the National Pesticides Information Center (NPIC) at **1-800-858-7378**, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>.

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION - Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. **ENVIRONMENTAL HAZARDS** - This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. **Groundwater Advisory** - This chemical has properties and characteristics

associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. **Surface Water Advisory** - This chemical may contaminate water through drift of spray in wind. This chemical has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this chemical. A level, well maintained vegetative buffer strip between areas to which this chemical is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this chemical will be reduced by avoiding applications when conditions favor runoff (for example when soils are saturated and/or significant rainfall is forecast in the next 48 hours). Sound erosion control practices will reduce this chemical's contribution to surface water contamination. **Physical or Chemical Hazards - DO NOT** use or store near heat or open flame.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE:** Keep this product in its tightly closed original container, when not in use. Store in a cool, dry area that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility. **CONTAINER HANDLING:** Less than or equal to 50 pounds: Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

See label booklet for complete Precautionary Statements and Directions For Use.

Manufactured For:

Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 82633-133-83529

EPA Est. No. AG 72159-GA-001; GH 70815-GA-002;

MA 83411-MN-001; TX 07401-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: 0.75 lb.

OPEN HERE