

Gamla

INSECTICIDE

For both indoor and outdoor use.

For Use In, On and Around Buildings and Structures for the Control of Listed Pests: on Non-Residential Lawns and Turf, Ornamental Trees and Shrubs Around Non-Residential, Institutional, Commercial, Agricultural, and Industrial Buildings; Livestock/Poultry Housings and Pet Kennels.

ACTIVE INGREDIENT:	WT. BY %
Gamma-cyhalothrin: Cyclopropanecarboxylic acid, 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl, cyano(3-phenoxyphenyl)methyl ester*	5.9%
OTHER INGREDIENTS:	94.1%
TOTAL:	100.0%
Contains 0.5 lbs. of active ingredient per gallon. Contains petroleum distillate. *Synthetic pyrethroid, CAS No. 76703-62-3, capsule suspension (microencapsulated).	

KEEP OUT OF REACH OF CHILDREN / MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

See label booklet for complete First Aid,
Precautionary Statements, Directions For Use,
and Storage and Disposal.

EPA Reg. No. 83529-328

Manufactured For:

Sharda USA LLC



7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Est. No. (AG) 72159-GA-001; (MC) 89332-GA-001; (SC) 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: **1 Quart**

FIRST AID	
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	• Immediately call a poison control center or doctor for treatment advice. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give any liquid to the person. • DO NOT give anything by mouth to an unconscious person.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	
NOTE TO PHYSICIAN	
Induced vomiting as first aid for this substance may result in increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent. Vomiting should be induced only under professional supervision.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION / PRECAUCIÓN

Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Skin exposure may also result in a sensation described as tingling, itching, burning or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2 to 30 hours, without damage. Wash exposed areas thoroughly with soap and water. Relief from skin sensation may be obtained by applying an oil-based cream.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, made of barrier laminate, nitrile rubber $\geq$ 14 mils, neoprene rubber $\geq$ 14 mils, or Viton® $\geq$ 14 mils
- Shoes plus socks
- Coveralls
- Protective eyewear
- Respirator - wear minimum of a NIOSH-approved elastomeric half mask respirator with organic vapor (OV) cartridges and combination N, R, or P filters; OR a NIOSH-approved gas mask with OV canisters; OR a NIOSH-approved powered air purifying respirator with OV cartridges and combination HE filters.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment wash waters.

To protect the environment, **DO NOT** allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help to avoid run off to water bodies or drainage systems.

DO NOT allow applications to contact water inhabited by fish, such as aquariums and ornamental fish ponds that are located in and around structures being treated.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. **Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms.**



PROTECTION OF POLLINATORS

APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.

Look for the bee hazard icon in the **DIRECTIONS FOR USE** for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators. Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar. Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or direct to EPA at: beekill@epa.gov.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT use or store near heat or open flame. **DO NOT** use Gamla in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For both indoor and outdoor use.

SHAKE WELL BEFORE USING. Avoid freezing. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

USE PRECAUTIONS AND RESTRICTIONS

- **DO NOT** use or dilute this product with oil.
- **DO NOT** allow to enter indoor or outdoor drains.
- Follow proper disposal procedures on this label.
- All outdoor spray applications **MUST** be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - o Application to pervious surfaces such as soil, non-residential lawn, turf, and other vegetation;
 - o Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or non-residential lawn);
 - o Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
 - o Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
 - o Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
 - o Applications to vertical surfaces directly above pervious surfaces, such as soil, non-residential lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.
- Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).
- For soil or foliar applications, **DO NOT** apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

- LET TREATED SURFACES DRY BEFORE ALLOWING HUMANS AND PETS TO CONTACT SURFACES.
- **DO NOT** apply the product into fish pools, ponds, streams, or lakes.
- Application is prohibited directly into sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. **DO NOT** allow the product to enter any drain during or after application.
- **DO NOT** apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- **DO NOT** apply or irrigate to the point of runoff.
- Apply this product directly to non-residential lawn or garden area. Water treated area as directed on this label. **DO NOT** water to the point of run-off.
- **DO NOT** make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- **DO NOT** apply when the wind speed is greater than 10 mph.
- Treat surfaces to ensure thorough coverage but avoid runoff.
- To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.
- Removable chemical extraction probes (also known as "stingers") used in suction/ extraction systems must be rinsed within the pesticide container prior to removal.
- This product is **RESTRICTED** for use in New York state.
- **Nursery (ornamentals, trees, container stock)**
 - o **DO NOT** apply as foliar broadcast application using a mechanically-pressurized handgun to nurseries.
 - o **DO NOT** apply as drench/soil/ground-direct application methods using a mechanically pressurized handgun to nurseries.
- **Indoor Residential Sites:**
 - o **DO NOT** apply to carpets as a pin stream or coarse spray treatment.
 - o **DO NOT** apply this product in any room being used as living, eating, sleeping, or recovery area by patients, the elderly, or infirm when they are in the room.
 - o **DO NOT** apply to institutions (including libraries, sports facilities, etc.) in the immediate area when occupants are present.
 - o **DO NOT** apply to classrooms when in use.

- o **DO NOT** apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- o **DO NOT** allow applications to contact water inhabited by fish, such as in aquariums and ornamental fishponds that are located in/around structures being treated.



• **Foliar Application**

Follow application instructions as indicated in **Bee Hazard Direction for Use**.

For commercially grown ornamentals not under contract for pollination services but are attractive to pollinators:



- **DO NOT** apply this product while bees are foraging.
- This product is toxic to bees exposed to residue for more than 38 hours following treatment.
- **DO NOT** apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by the appropriate State or Federal authorities.

INSECTICIDE RESISTANCE MANAGEMENT

For resistance management, **Gamla** contains a Group 3a insecticide. Any insect/mite population may contain individuals naturally resistant to **Gamla** and other Group 3a insecticides. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide/acaricide resistance, take the following steps:

- Rotate the use of **Gamla** or other Group 3a insecticides/acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides/acaricides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - o Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.

- o Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
- o When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
- o Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- o The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity.
- o Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Sharda USA LLC. You can also contact your pesticide distributor or university extension specialist to report resistance.

MIXING INSTRUCTIONS

Dilute **Gamla** with water for application using hand-held or power-operated application equipment as a coarse spray for crack and crevice or spot and general surface (non-food/non-feed areas only) treatments. Application equipment that delivers low volume treatments, such as the Micro-Injector® or Actisol® applicator, may also be used to make crack and crevice or spot and general surface treatments. Fill applicator tank with the desired volume of water and add **Gamla**. Close and shake before use in order to ensure proper mixing. Shake or re-agitate applicator tank before use if application is interrupted. Mix only amount of treatment volume as required. **Gamla** may be applied by using a paintbrush or other porous applicator attached to a handle as a general surface treatment.

TANK MIXING

Gamla may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. A small volume mixing test with the other products is recommended to ensure compatibility. If other chemicals are added to the applicator tank, **Gamla** should be added last. If mixed with EC formulations, use within 24 hours. Fill tank to desired volume and continue to agitate while making applications.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Gamla may be tank mixed with an Insect Growth Regulator (IGR). Observe all restrictions and precautions which appear on the labels of these products.

FOAM APPLICATIONS

Gamla may be converted to a foam and the foam used to treat premise voids to control or prevent pests including ants, bees, termites (above ground only), wasps, or other arthropods harboring in walls, under slabs, or in other void areas.

INDOOR PREMISE PEST CONTROL

Gamla is for use as a general surface (non-food/non-feed areas), crack and crevice, or spot treatment in, on, and around buildings and structures and their immediate surroundings, and on modes of transport. Areas of use: industrial buildings, patios, porches, closets, furniture, apartment buildings, mobile homes, laboratories, buses, greenhouses (non-commercial), stores, factories, warehouses, wineries, vessels, railcars, trucks, trailers, aircraft (cargo and other non-cabin areas only), schools, nursing homes, hospitals, mausoleums, restaurants, hotels, correctional facilities, livestock/poultry housing, pet kennels, food granaries, food grain mills and food manufacturing, processing, and servicing establishments.

Non-food/non-feed areas of food/feed handling establishments: garbage rooms, lavatories, entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

For indoor applications, **Gamla** can be reapplied at 21-day intervals if necessary.

RATES FOR PESTS (HAND APPLICATION EQUIPMENT)

PESTS		DILUTION RATE (% a.i. and fl. oz.)
Ants	Fleas ²	0.0075% a.i.
Bedbugs (adult)	Flies (including House flies, stable flies, little house flies)	0.16 fl. oz. (5 ml.)/gal. water
Bees		0.5 fl. oz. (15 ml.)/
Boxelder	Lesser Grain Borers	3 gals. water
Bugs	Millipedes	Up to
Carpenter Bees	Mosquitoes	0.015% a.i.
Carpet Beetles	Red Flour Beetles	0.33 fl. oz. (10 ml.)/
Centipedes	Rice Weevils	gal. water
Cigarette Beetles	Saw-toothed Grain Beetles	
Clover Mites	Silverfish	
Cockroaches ¹	Sowbugs	
Confused Flour Beetles	Spiders	
Crickets	Ticks	
Earwigs	Wasps	
Firebrats		
Cockroaches ¹	Scorpions	0.03% a.i.
Crickets ⁶	Sowbugs	0.65 fl. oz. (20 ml.)/
Flies ⁵	Spiders ⁶	gal. water
Litter beetles ³ (Darkling, Hide, and Carrion beetles)	Spider Mites (Two-spotted, Spruce)	2.0 fl. oz. (60 ml.)/
Mosquitoes ⁴	Ticks ⁶	3 gals. water
Pillbugs		
Termites (above ground only)		0.03% a.i.
		0.65 fl. oz. (20 ml.)/
		gal. water
		2.0 fl. oz. (60 ml.)/
		3 gals. water

(continued)

RATES FOR PESTS (HAND APPLICATION EQUIPMENT) *(continued)*

¹For cockroaches, the recommended rate for maintenance treatments is 0.0075% a.i. and for clean-out treatments is 0.015% a.i.

For control of severe infestations, use 0.03% rate.

²For outdoor use only and use 0.015% rate.

³Use the 0.015% (0.33 fl. oz. (10 ml.)/gal. water) rate for light beetle infestation.

⁴For residual control, use 0.03% rate.

⁵In all states except California, rates for flies may be increased to 0.03% when environmental conditions are severe and/or populations are high.

⁶For clean out/severe infestations, use 0.03% rate*

Conversion Rate: 1 fluid ounce (fl. oz.) equals 29 milliliters (ml.).

One gallon finished dilution typically treats 500 - 1000 sq. ft.

Pest-Specific Use Directions

Ants: Apply to any trails around doors and windows and other places where ants may be found. Locate and treat nests. Where ants are trailing inside apply as a residual surface treatment to active areas such as baseboards, corners, around pipes, behind cabinets, behind and under refrigerators, sinks, furnaces, and stoves, cracks and crevices. When combining baits and residual surface insecticides, apply surface insecticides in cracks and crevices along baseboards and infested surfaces, and outside barrier treatments. Use baits in other areas that are untreated by residual insecticides, also see **Outdoor Surfaces and Perimeters**.

Bedbugs: Clean floors and surfaces by vacuuming. Apply as a coarse low pressure spray to harborage areas including crevices, baseboards, loose plaster, behind bed frames and headboards, beneath beds and furniture, and to bedsprings and bed frames. **DO NOT** apply to furniture surfaces or mattresses where people will be laying or sitting. Infested bedding should not be treated, but should be removed, placed in sealed plastic bags, and taken for laundering and drying at high temperatures.

Cockroaches, Crickets, Earwigs, Firebrats, Silverfish, and Spiders: Apply as a coarse low pressure spray to areas where these pests hide, such as baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, cabinets, behind and under refrigerators, sinks, furnaces and stoves, the underside of shelves, drawers, and similar areas. Pay particular attention to cracks and crevices, also see **Outdoor Surfaces and Perimeters**.

(continued)

RATES FOR PESTS (HAND APPLICATION EQUIPMENT) *(continued)*

Pest-Specific Use Directions *(continued)*

Bees, Flies, Mosquitoes, and Wasps: Apply directly to walls, ceilings, window screens, and other resting areas as a residual surface treatment. May be used inside residential buildings as well as in and around carports, garages, and storage sheds; also see **Outdoor Surfaces and Perimeters**. Use caution when treating nests of stinging insects as **Gamla** does not provide instant knockdown. Protective equipment for the applicator may be required. Treat bee, wasp, and hornet nests late in the day when most insects will be present. Allow 2 - 3 days for the colony to die and repeat if necessary.

Carpenter Bees: Apply coarse spray to thoroughly wet wood surfaces where bees have been previously active or to provide protection against further damage. Apply early in the spring to prevent bees from invading wood. When bees have infested wood surface applications can help control embedded larvae and bees that emerge from the wood.

Pantry Pests (i.e., Carpet beetle, Cigarette beetle, Confused flour beetle, Lesser grain borer, Red flour beetle, Rice weevil, and Saw-toothed grain beetle): Apply to cupboards, shelving, and storage areas. Remove all utensils, uncovered foodstuffs (or any having original package opened), and shelf paper before making application. Allow treated surfaces to dry and cover shelves with clean paper before replacing any utensils, foodstuff, or other items. Any foodstuff accidentally contaminated with treatment solution should be destroyed.

Boxelder Bugs, Centipedes, Millipedes, Pillbugs, and Sowbugs: Apply around doors and windows and other places where these pests may be found or where they may enter premises. Treat baseboards, storage areas, and other locations. Apply barrier treatments to prevent infestation as described below; also see **Outdoor Surfaces and Perimeters**.

Fleas and Ticks: To control nuisance fleas and ticks (e.g. dog ticks) apply to kennels, yards, runs, and other areas where pets may frequent. To control ticks, apply using a coarse fan spray to vegetation brush, branches, rock walls, and other areas near habitation where ticks may harbor or frequent. Treat entire area rather than spot treatments, and retreat monthly to maintain control. **DO NOT** apply to pasture or cropland, and **DO NOT** allow animals and people access to treated areas until the deposit has dried. Applications can begin in the spring and can continue until frost to control both larvae and adult ticks; also see **Outdoor Surfaces and Perimeters**.

(continued)

RATES FOR PESTS (HAND APPLICATION EQUIPMENT) *(continued)*

Pest-Specific Use Directions *(continued)*

Cluster Flies: Apply in late summer or early fall before flies are observed alighting on surfaces. Apply thoroughly on siding, under eaves, and around windows and doors, paying particular attention to south-facing surfaces, but **DO NOT** allow runoff to occur. Heavy precipitation prior to frost may require retreatments to maintain protection. In winter and spring when flies become active and are emerging, interior crack and crevice and void treatments can help reduce the infestation, along with ULV or general surface application in infested attics or unoccupied lofts.

Litter Beetles (Darkling, Hide, and Carrion Beetles) and Flies in Animal Housing (Such as Poultry Houses): To control adult litter beetles, apply **Gamla** to walls and floors and cleanout, before reintroduction of animals. This will suppress beetles that escaped earlier treatment and will help delay onset of future infestations. Pay attention to area where beetles frequently occur, such as walls, supports, cages, stalls, and around feeders. To help control flies, apply a directed application to horizontal surfaces and overhead areas and allow to dry before reintroduction of animals; also see **Livestock/Poultry Housing Structures and Pet Kennels**.

* **Gamla** may be used on termites to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. These applications are not a substitute for mechanical alteration to control subterranean termites, soil treatment labeled termiticides, or fumigation for extensive infestation of drywood termites or other wood-infesting insects but are considered supplemental. This product is not recommended as sole protection against termites. For active termite infestations, get a professional inspection.

Application Within Food Handling Establishments

For places other than private residences in which exposed food is held processed prepared or served areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing foods, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups), and serving areas.

Use as a crack and crevice or spot treatment in and around both food and nonfood areas. Apply in small amounts directly into cracks and crevices, using equipment capable of delivering a pin stream of insecticide, in points between different elements of construction, between equipment and floor, openings leading to voids and hollow spaces in walls, equipment and bases. Food contact surfaces and equipment should be cleaned with an effective cleaning compound and rinsed with potable water before using.

Limit individual spot treatments to an area no larger than 20% of the treated surface. Individual spot treatments may not exceed 2 sq. ft. Take extreme care that the product is not introduced into the air. Avoid contamination of food and food processing surfaces.

Application Within Food Serving Areas

For facilities where food is served: dining rooms. Apply as a crack and crevice or spot treatment to selective surfaces: baseboards, under elements of construction, and into cracks and crevices. **DO NOT** treat surfaces likely to be contacted by food.

Restrictions for food handling establishments

- **DO NOT** apply when facility is in operation or foods are exposed. Food must be covered or removed in area being treated.
- **DO NOT** apply directly to food or allow applications to contaminate food.
- **DO NOT** make application of this product in Food Areas and/or Food Serving Areas of Food Handling Establishments other than as a spot and/or crack and crevice treatment. Use of application equipment such as the Micro-Injector® or Actisol® applicator in food areas should be limited to crack and crevice treatment only.

Livestock/Poultry Housing Structures and Pet Kennels

Gamla may be used as a general surface and/or crack and crevice treatment. Treat interior and exterior perimeter applications in and around livestock, poultry and pet housing structures for optimum control. Suitable cleaning practices of the structure must also be followed along with applications of **Gamla** to effectively control the listed insects.

Litter Beetles: To control adult litter (darkling, hide and carrion beetles) beetles in animal housing, including poultry housing, apply **Gamla** to walls and floors at each grow-out or sanitation procedure, before reintroduction of poultry or other animals. Re-apply after each grow-out or sanitation procedure to control beetles that escaped earlier treatment and delay the onset of further infestations. Beetles frequently occur in areas such as on walls, supports, cages, stalls, and around feeders, so pay careful attention to these places. For poultry houses where birds are grown on litter, treat the floor area. For poultry houses where birds are grown in cages, treat the walls, posts, cage framing, and into cracks and crevices around insulation. Also apply as a perimeter treatment around the outside of the building foundations to prevent beetles from entering from outside. To prevent populations from reaching a problem level, a year-round maintenance program is highly recommended.

RESTRICTIONS to Livestock/Poultry Housing Structures and Pet Kennels:

- **DO NOT** make applications of **Gamla** in areas where animals, except for cattle and calves are present in the facility. Allow treated surfaces to dry completely before restocking the facility.
- **DO NOT** make applications to any animal feedstuffs water or watering equipment.
- **DO NOT** contaminate any animal food feed or water in and around livestock, poultry or pet housing when making applications.

OUTDOOR SURFACES AND PERIMETERS

To control flies, fleas, mosquitoes, ants, bees, wasps, centipedes, cockroaches, crickets, pillbugs, scorpions, sowbugs, spiders, termites* (above ground only), ticks and other listed pests, apply **Gamla** with either hand or power application equipment as a residual treatment to ornamental plants next to foundations of buildings and to external surfaces of buildings, porches, screens, door and window frames, eaves, patios, garages, refuse dumps and other areas where insects rest or congregate. **Gamla** may also be applied to lawns around non-residential buildings, or to refuse dumps or other non-residential areas where these insect pests congregate. Remove all exposed food and cooking utensils. Cover all food handling surfaces or wash thoroughly after treatment and before use. **DO NOT** use on edible crops.

For Perimeter treatments, apply a continual band of insecticide around building foundations and around windows, doors, service line entrances, eaves, vents, and other areas to reduce the potential for entry by crawling pests. Before application, remove debris and leaf litter from next to the foundation, cut back vegetation and branches that touch the foundation, and move or rake back rocks, deep mulch, or other potential pest harborage next to the foundation. Apply the band up to 7 ft. wide around the structure (or according to state regulations governing commercial pest control) and upwards along the foundation to 2 ft. and around windows, doors, and roof overhangs. Apply as a coarse spray to thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active.

The amount of concentrate is dependent upon pest species (see pest table and comments), infestation levels, and service interval desired.

Rate Table for Perimeter Barrier Applications

Application Rate: Fl. oz. (ml.) of Gamla	Gals. of Water ¹	Area of Coverage (sq. ft.)
0.16 fl. oz. (5 ml.)	1 - 5 gals.	800 - 1600 sq. ft.
0.33 fl. oz. (10 ml.)	1 - 5 gals.	800 - 1600 sq. ft.
0.65 fl. oz. (20 ml.)	1 - 5 gals.	800 - 1600 sq. ft.
¹ Application volume may be greater than 5 gals./800 - 1600 sq. ft. if required under heavy vegetative or landscaping materials in order to obtain desired coverage.		

Examples of Dilutions for Perimeter Barrier Applications

Application Volume Gals. of Solution per 1,000 sq. ft.	Application Rate Fl. oz. (ml.) of Gamla per 1,000 sq. ft.	Fl. oz. (ml.) of Gamla to Dilute in Water According to Spray Tank Volumes		
		5 gals.	10 gals.	50 gals.
1 gal. per 1,000 sq. ft.	0.16 fl. oz. (5 ml.)	0.80 fl. oz. (25 ml.)	1.6 fl. oz. (50 ml.)	8 fl. oz. (250 ml.)
	0.33 fl. oz. (10 ml.)	1.65 fl. oz. (50 ml.)	3.3 fl. oz. (100 ml.)	16 fl. oz. (500 ml.)
	0.65 fl. oz. (20 ml.)	3.3 fl. oz. (100 ml.)	6.5 fl. oz. (200 ml.)	33 fl. oz. (1000 ml.)
2 gals. per 1,000 sq. ft.	0.16 fl. oz. (5 ml.)	0.40 fl. oz. (12.5 ml.)	0.80 fl. oz. (25 ml.)	4 fl. oz. (125 ml.)
	0.33 fl. oz. (10 ml.)	0.80 fl. oz. (25 ml.)	1.65 fl. oz. (50 ml.)	8 fl. oz. (250 ml.)
	0.65 fl. oz. (20 ml.)	1.65 fl. oz. (50 ml.)	3.3 fl. oz. (100 ml.)	16 fl. oz. (500 ml.)

(continued)

Examples of Dilutions for Perimeter Barrier Applications (continued)

Application Volume Gals. of Solution per 1,000 sq. ft.	Application Rate Fl. oz. (ml.) of Gamla per 1,000 sq. ft.	Fl. oz. (ml.) of Gamla to Dilute in Water According to Spray Tank Volumes		
		5 gals.	10 gals.	50 gals.
5 gals. per 1,000 sq. ft.	0.16 fl. oz. (5 ml.)	0.16 fl. oz. (5 ml.)	0.33 fl. oz. (10 ml.)	1.6 fl. oz. (50 ml.)
	0.33 fl. oz. (10 ml.)	0.33 fl. oz. (10 ml.)	0.65 fl. oz. (20 ml.)	3.3 fl. oz. (100 ml.)
	0.65 fl. oz. (20 ml.)	0.65 fl. oz. (20 ml.)	1.65 fl. oz. (50 ml.)	6.5 fl. oz. (200 ml.)

Example calculation: to apply the mid-rate of **Gamla** at a volume of 5 gals./1,000 sq. ft. mix 3.3 oz of concentrate in 50 gallons of water. The percent active ingredient in the finished **Gamla** dilution can be calculated with the following formula:

mL needed to add times 5.9% active in concentrate, divided by gal. finished dilution times 3785 ml./gal. = % active in dilution (Example: 3.3 oz. in 50 gals. is 100 ml. times 5.9 equals 590, and 50 gals., times 3785 is 189,250. Dividing 590 by 189,250 equals 0.003% active in the tank dilution.)

Restrictions for outdoor surfaces and perimeters

- **DO NOT** use water-based sprays of **Gamla** in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard. Thoroughly wash out sprayer and screen with water and detergent before using **Gamla**. **Gamla** has not stained or caused damage to painted or varnished surfaces, plastics fabrics, or other surfaces where water applied alone causes no damage. However, treat a small area and allow to dry, to determine whether staining will occur.

NON-RESIDENTIAL LAWNS/TURFGRASS AND ORNAMENTALS

Gamla may be used for applications to maintain non-residential indoor or outdoor areas where turf and ornamentals are grown: non-residential landscapes around institutional, commercial and industrial buildings. Application rates for turf and ornamentals are lower than premise pest control rates, reflecting that treatment intervals are more frequent.

Applicators must ensure that they are certified by the necessary pesticide certification categories to allow application of **Gamla** away from structures such as to turf and ornamental plantings. Premise pest control certification categories may limit the distance away from structures for pesticide application. Consult your state extension office or pesticide regulatory officials for further information.

IMPORTANT: Time application to flowering plants during periods when pollinating insects are not present such as early morning or late evening.

RESTRICTIONS:

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product to edible crops.
- **DO NOT** apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms **DO NOT** apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, estuaries, and commercial fish farm ponds.
- **DO NOT** make outdoor broadcast applications to turf and ornamentals when wind speed is 10 mph or greater.
- In the state of New York, **DO NOT** apply within 100 ft. of coastal marshes or streams that drain into coastal marshes.

MIXING INSTRUCTIONS

Gamla is to be mixed with water and may be used in all types of standard application equipment. Fill applicator tank with the desired volume of water and add **Gamla**. It is suggested that the water be 5 - 7 pH. Adjust water pH with a buffering agent if necessary. Slowly add **Gamla** to applicator tank water with maximum agitation. Close and shake or re-agitate applicator tank before use if application is interrupted. Make up only amount of treatment volume as required.

TANK MIXING

Gamla may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. A small volume mixing test with the other products is recommended to ensure compatibility. If other chemicals are added to the applicator tank, **Gamla** should be added last. Fill tank to desired volume and continue to agitate while making applications. If mixed with EC formulations use within 24 hours. Observe all restrictions and precautions which appear on the labels of these products.

INSECT CONTROL ON ORNAMENTAL PLANTS

Gamla is for use to control the pests listed in the table below: non-residential landscaped areas and landscaped areas around institutional, commercial and industrial buildings.

NOTE: While phytotoxicity testing has been earned out on a wide range of ornamental plants under various environmental conditions and no phytotoxicity has been observed certain cultivars may be sensitive to the final spray solution.

It is advised to pre-spray a selection of ornamental plants and observe them for 7 - 10 days prior to treating large areas if local use experience is unavailable.

RESTRICTIONS:

- **DO NOT** apply more than 0.18 lbs. of the Gamma-Cyhalothrin (46.1 fl. oz. of concentrate) per acre per year.

TANK DILUTION RATES FOR ORNAMENTAL PESTS

PEST		AMOUNT OF GAMLA	APPLICATION INSTRUCTIONS
Ants (Including Imported fire Ants)	Crickets	1.3 - 4.4 fl. oz. per 100 gals.	Application to ornamentals should be started prior to the establishment of high insect pest populations. Apply at 7-day intervals if retreatment is necessary. Limit more frequent treatment to spot treatments. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at recommended rates may enhance the control of insects on certain species of ornamentals having waxy, hard to wet foliage. For spot treatments use 0.44 fl. oz. Gamla per 1 - 2.5 gals. of water. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area. Scale: Thoroughly cover the plant with Gamla spray including trunks, stems, twigs, and foliage for control of scale insects (crawler stage). Bagworm: Apply Gamla when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young.
Armyworms	Cutworms	OR	
Azalea caterpillars	Eastern tent caterpillars	39 - 132 ml. per 100 gals.	
Aphids	Elm leaf beetles		
Bagworms	European sawflies		
Black vine weevils (adult)	Fall webworms		
Boxelder bugs	Flea beetles		
Budworms	Forest tent caterpillars		
California oakworms	Gypsy moth larvae		
Cankerworms	Japanese beetles (adults)		
Cockroaches	June beetles (adults)		

TANK DILUTION RATES FOR ORNAMENTAL PESTS (continued)

PEST		AMOUNT OF GAMLA	APPLICATION INSTRUCTIONS
Lace bugs	Pinetip moths	1.3 - 4.4 fl. oz. per 100 gals.	Application to ornamentals should be started prior to the establishment of high insect pest populations. Apply at 7-day intervals if retreatment is necessary. Limit more frequent treatment to spot treatments. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at recommended rates may enhance the control of insects on certain species of ornamentals having waxy, hard to wet foliage. For spot treatments use 0.44 fl. oz. Gamla per 1 - 2.5 gals. of water. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area. Scale: Thoroughly cover the plant with Gamla spray including trunks, stems, twigs, and foliage for control of scale insects (crawler stage). Bagworm: Apply Gamla when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young.
Leaf-feeding caterpillars	Plant bugs	OR	
Leafhoppers	Root weevils	39 - 132 ml. per 100 gals.	
Leafminers	Sawflies		
Leafminers (adults)	Scale insects (crawlers)		
Leaf rollers	Spiders		
Leaf skeletonizers	Spittlebugs		
Midges	Striped beetles		
Mosquitoes	Striped oakworms		
Oleander moth larvae	Thrips		
Pillbugs	Tip moths		
Pine sawflies	Tussock moth larvae		
Pine shoot beetles	Wasps		
Broad mites	Mealybugs	2.6 - 4.4 fl. oz. per 100 gals.	
Brown soft scales	Pine needle scales (crawlers)	OR	
California red scales (crawlers)	Spider mites	78 - 132 ml. per 100 gals.	
Clover mites	Whiteflies		

Gamla Mixing Chart for Ornamental Insect Pest Control

Rate of Gamla	25 Gals.	50 Gals.	100 Gals.	200 Gals.	300 Gals.
1.32 oz. ¹	0.30 oz.	0.70 oz.	1.32 oz.	2.6 oz.	3.96 oz.
2.64 oz. ²	0.70 oz.	1.30 oz.	2.64 oz.	5.3 oz.	7.92 oz.
4.4 oz. ³	1.10 oz.	2.20 oz.	4.40 oz.	8.8 oz.	13.2 oz.

¹ Equivalent to 3.2 ml. per 1,000 sq. ft. (4.7 oz. per acre) when applied at 8 gals. per 1,000 sq. ft.
² Equivalent to 6.3 ml. per 1,000 sq. ft. (9.3 oz. per acre) when applied at 7 gals. per 1,000 sq. ft.
³ Equivalent to 10.6 ml. per 1,000 sq. ft. (15.6 oz. per acre) when applied at 8 gals. per 1,000 sq. ft.

NON-RESIDENTIAL LAWN/TURFGRASS INSECT PEST CONTROL

Gamla is for use to control the pests listed in the table below around non-residential, institutional, commercial and industrial buildings.

RESTRICTIONS:

- **KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL SPRAY HAS DRIED FOLLOWING APPLICATION.**
- **DO NOT** apply when turfgrass is waterlogged or when soils are saturated with water (i.e. will not accept irrigation).
- **DO NOT** apply more than 0.18 lbs. of a.i. (46.1 fl. oz. of concentrate) per acre per year.

RATES FOR NON-RESIDENTIAL LAWN AND TURFGRASS PESTS

PEST		AMOUNT OF GAMLA	APPLICATION INSTRUCTIONS
Ants (Including Imported fire Ants)	Millipedes Mites Mosquitoes (adult)	3.0 - 6.1 ml. per 1,000 sq. ft.	Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. To keep pest populations under control, use higher labeled rates as pest pressure increases. Apply at 7-day intervals if retreatment is necessary. Limit more frequent treatment to spot treatments. For spot treatments, use 0.20 fl. oz. of Gamla per 1 - 2.5 gals. of water.
Armyworms	Pillbugs	OR	
Centipedes	Sod webworms	4.4 - 8.8 fl. oz. per acre	
Crickets	Sow bugs		
Cutworms	Ticks (including species which transmit Lyme disease)		
Earwigs			
Fleas (adult)			
Grasshoppers			
Japanese beetles (adult)			
Bluegrass billbugs (adult)	Grubs (suppression)	6.1 ml. per 1,000 sq. ft.	
Black turfgrass	Hyperodes	OR	
Ataenius (adult)	weevils (adult)	8.8 fl. oz. per acre	
Chiggers	Mole crickets ¹		
Chinch Bugs ¹	(nymphs and young adults)		
Fleas (adult)			
¹ Not for use on mature adult mole crickets and chinch bugs in New York state.			
*Example: To treat the listed turf pests at the 3.5 ml. per 1,000 sq. ft. rate, determine the gallons of dilution per 1,000 sq. ft. needed to adequately cover the turf. At a 5 gallon per 1,000 sq. ft. dilution, take the 3.5 ml. and divide it by the 5 (gallon) to get 0.7 ml. to be added per gallon. For a 50 gallon tank, therefore, this would be 0.7 ml. X 50 = 35 ml. (or 1.25 oz.) in 50 gallons of water.			
Conversion Rate: 1 fluid ounce (fl. oz.) equals 29 milliliters (ml.).			

(continued)

RATES FOR NON-RESIDENTIAL LAWN AND TURFGRASS PESTS *(continued)*

Pest-Specific Application Instructions

Surface Insect Control (armyworm, cutworms, fleas, etc.): Apply **Gamla** at recommended rates in 2 - 5 gals. of water per 1,000 sq. ft. The use of spreader-sticker may be useful if high rainfall amounts are forecast, otherwise the addition of adjuvants is not necessary under normal conditions for surface insect control in turf. Delay watering or mowing for 12 - 24 hours to control surface-feeding insect pests.

Thatch Inhabiting Insect Control (chinch bugs, billbugs, etc.): Apply **Gamla** at recommended rates in 2 - 10 gals. of water per 1,000 sq. ft. The use of a nonionic wetting agent, penetrant or similar adjuvant is recommended at label rates. Lightly irritate after application with up to 1/2 inch of water to move the **Gamla** into the thatch layer. If irrigation is not available, then use high water application rates.

Subsurface Insect Control (mole crickets, grubs, etc.): Apply **Gamla** at recommended rates in 4 - 10 gals. of water per 1,000 sq. ft. The use of a nonionic wetting agent, penetrant, or similar adjuvant is strongly recommended following label rates. Use the highest water application rates possible with your sprayer. Apply **Gamla** to turf wet with dew, rain, or irrigation. Water-in immediately after application 1/4 - 1/2 inch of water.

Fire Ant Control: Treat individual mounds with a drench application using a watering can. Use 0.20 fl. oz. of **Gamla** per 2.5 gals. of water. Thoroughly soak each mound and a 3 ft. diameter circle around each mound. Gently apply the mixture to avoid disturbing the mound. Disturbing the mound may cause the ants to migrate and reduce the effectiveness of the treatment. Apply in early morning or late evening hours. You may make applications on a monthly basis to maintain protection of treated areas.

Mosquito Control: Apply as a general spray around landscape plantings, turf, and building foundations to control mosquitoes. Apply **Gamla** at recommended rates in 2 - 5 gals. of water per 1,000 sq. ft.

Gamla Mixing Chart for Non-residential Turf Insect Pest Control

Rate of Gamla Insecticide	App. Rate per 1,000 sq. ft. of Turf (oz.)					
Fl. oz. per Acre	2 gals.	4 gals.	5 gals.	6 gals.	8 gals.	10 gals.
4.4 fl. oz. per acre	5.1	2.5	2.0	1.7	1.3	1.0
6.6 fl. oz. per acre	7.6	3.8	3.0	2.5	1.9	1.5
8.8 fl. oz. per acre	10.1	5.1	4.0	3.4	2.5	2.0
Conversion rate: 1 fluid ounce (fl. oz.) equals 29 milliliters (ml.).						

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage and disposal. **DO NOT** pour or dispose down-the-drain or sewer. Call your local solid waste agency for local disposal options.

PESTICIDE STORAGE: Store this product in a cool, dry place in its original container only. **DO NOT** store this product near fertilizers, seeds, or other pesticides. If this product is spilled, sweep up the spillage and dispose pursuant to the below **PESTICIDE DISPOSAL** instructions.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.



(continued)

STORAGE AND DISPOSAL *(continued)*

CONTAINER HANDLING:

Less Than or Equal to 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

Greater Than 5 Gallons: Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For Bulk and Mini-Bulk Containers: Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

PRIMEROS AUXILIOS	
SI ENTRA EN CONTACTO CON LOS OJOS:	• Mantenga los ojos abiertos y enjuáguelos lenta y cuidadosamente con agua, durante 15 a 20 minutos. • Si utiliza lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
SI SE INGIERE:	• Llame de inmediato a un centro de control de envenenamientos o a un médico para consejo de tratamiento. • Si la persona puede tragar, haga que beba un vaso de agua lentamente. • NO induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. • NO administre nada por boca a una persona que haya perdido el conocimiento.
NÚMERO DE TELÉFONO DIRECTO	
Cuando llame a un centro de control de envenenamiento, o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. Para información de emergencia sobre este producto, llame al centro de control de envenenamientos a 1-800-222-1222.	
NOTA AL MÉDICO	
El vómito inducido como primeros auxilios para esta sustancia puede aumentar el riesgo de neumonía química o edema pulmonar causado por la aspiración del disolvente de hidrocarburos. El vómito debe inducirse únicamente bajo supervisión profesional.	

DECLARACIONES PREVENTIVAS

RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS

PRECAUCIÓN

Causa irritación moderada en los ojos. Dañino si se ingiere. Evite el contacto con los ojos, la piel o la ropa. Lávese bien con agua y jabón después de manipular y antes de comer, beber, masticar chicle, usar tabaco o usar el baño. Quítese y lave la ropa contaminada antes de reutilizarla.

El contacto prolongado y frecuente con la piel puede causar reacciones alérgicas en algunos individuos. La exposición cutánea también puede causar una sensación de hormigueo, picazón, ardor o sensación de pinchazos. La sensación puede aparecer inmediatamente o hasta 4 horas después de la exposición y puede durar de 2 a 30 horas sin causar daños. Lave a fondo las áreas expuestas con agua y jabón. El alivio de la sensación en la piel se puede obtener aplicando una crema a base de aceite.

EQUIPO DE PROTECCIÓN PERSONAL (PPE, POR SUS SIGLAS EN INGLÉS)

Los mezcladores, cargadores, aplicadores y otros manipuladores deben usar:

- Camisa de manga larga y pantalones largos
- Guantes resistentes a productos químicos fabricados con laminados impermeables, goma de nitrilo ≥ 14 mils, goma de neopreno ≥ 14 mils, o Viton® ≥ 14 mils
- Zapatos y calcetines
- Overol (mameluco)
- Gafas protectoras
- Respirador – utilice como mínimo, un respirador con purificador de aire y máscara para medio rostro o semimáscara (elastomérico) aprobado por NIOSH que esté equipado con un cartucho químico para vapores orgánicos (OV) en combinación con un filtro purificador o material de filtración de las series N, R, o P; o una mascarilla de filtrado para gases aprobada por NIOSH que esté equipada con cánters OV; o un respirador purificador de aire motorizado aprobado por NIOSH equipado con cartuchos OV en combinación con filtros HE.

Siga las instrucciones del fabricante para la limpieza y el mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE por separado del resto de la ropa.

RIESGOS AMBIENTALES

Este pesticida es extremadamente tóxico para los peces y los organismos acuáticos y tóxico para la vida silvestre. **NO** aplique directamente al agua, en áreas donde haya aguas superficiales, o en áreas intermareales que se encuentren bajo el nivel de altura máxima promedio del agua. **NO** aplique cuando las condiciones meteorológicas favorezcan la deriva desde las áreas tratadas. La deriva y la escorrentía pueden ser peligrosas para los organismos acuáticos en áreas adyacentes. **NO** contamine el agua al desechar el agua de lavado del equipo.

Para proteger el ambiente, **NO** permita que el pesticida entre o sea descargado en desagües pluviales, zanjas de drenaje, cunetas, o cuerpos de agua superficiales. Aplicar este producto cuando el clima esté tranquilo y sin lluvia pronosticada en las próximas 24 horas, ayudará a asegurar que el pesticida no será llevado fuera del área tratada debido a viento o lluvia. Lavar el equipo de aplicación dentro del área de tratamiento ayudará evitar escorrentías a cuerpos de agua o sistemas de drenaje.

NO permita que las aplicaciones entren en contacto con aguas habitadas por peces, tales como acuarios y estanques ornamentales, que se encuentren dentro o en los alrededores de las estructuras que están siendo tratadas.

DECLARACIÓN CONSULTIVA SOBRE ORGANISMOS NO OBJETIVO: Este producto es altamente tóxico para las abejas y otros insectos polinizadores expuestos al tratamiento directo o a residuos en/sobre cultivos en floración o malezas. **NO** aplique este producto ni deje que se disperse sobre cultivos en flor o malas hierbas mientras las abejas estén visitando activamente la zona de tratamiento. **Proteja a los insectos polinizadores siguiendo las instrucciones de la etiqueta destinadas a minimizar la deriva y reducir el riesgo para estos organismos.**



PROTECCIÓN DE POLINIZADORES

EXISTEN RESTRICCIONES DE APLICACIÓN PARA ESTE PRODUCTO DEBIDO AL RIESGO PARA LAS ABEJAS Y OTROS INSECTOS POLINIZADORES. SIGA LAS RESTRICCIONES DE APLICACIÓN INDICADAS EN LAS INSTRUCCIONES DE USO PARA PROTEGER A LOS POLINIZADORES.

Busque el icono de peligro para las abejas en las **INSTRUCCIONES DE USO** para cada sitio de aplicación a fin de conocer las restricciones de uso específicas y las instrucciones para proteger a las abejas y otros insectos polinizadores.

Este producto puede matar a las abejas y a otros insectos polinizadores. Las abejas y otros insectos polinizadores se alimentan en las plantas cuando estas florecen, liberan polen o producen néctar. Las abejas y otros insectos polinizadores pueden estar expuestos a este plaguicida por:

- El contacto directo durante aplicaciones foliares o el contacto con residuos en las superficies de las plantas después de las aplicaciones foliares.
- La ingestión de residuos en el néctar y el polen cuando el plaguicida se aplica como tratamiento de semillas, al suelo, mediante inyección en árboles, así como en aplicaciones foliares.

Al usar este producto, tome medidas para:

- Minimizar la exposición de este producto a las abejas y otros insectos polinizadores cuando estén forrajeando en plantas atractivas para los polinizadores alrededor del sitio de aplicación.
- Minimizar la deriva de este producto hacia las colmenas o hacia hábitats atractivos para los polinizadores fuera del sitio de aplicación. La deriva de este producto hacia las colmenas o fuera del sitio hacia hábitats atractivos.

Puede encontrar información sobre la protección de las abejas y otros insectos polinizadores en el sitio web de Administración Ambiental de Plaguicidas en:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>

Los incidentes relacionados con plaguicidas (por ejemplo, la muerte de abejas) deben reportarse inmediatamente a la agencia principal estatal o tribal correspondiente. Para obtener información de contacto de su estado, visite: www.aapco.org/officials.html.

PELIGROS FÍSICOS Y QUÍMICOS

NO utilice ni almacene cerca del calor o llamas abiertas. **NO** utilice aerosoles a base de agua de **Gamla** en conductos, carcasas de motores, cajas de conexiones, cajas de interruptores u otros equipos eléctricos debido al posible riesgo de descarga eléctrica.

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal.

ALMACENAMIENTO Y DESECHO

NO contamine el agua, los alimentos ni los forrajes durante el almacenamiento o el desecho. **NO** verter ni desechar en el desagüe ni en el alcantarillado. Llame a su agencia local de residuos sólidos para conocer las opciones de eliminación local.

ALMACENAMIENTO DE PESTICIDAS: Almacene este producto en un lugar fresco y seco, únicamente en su envase original. **NO** lo guarde cerca de fertilizantes, semillas ni otros pesticidas. En caso de derrame, recoja el derrame y deséchelo siguiendo las instrucciones de eliminación de pesticidas que se indican a continuación.

DESECHO DE PESTICIDAS: Los residuos de pesticidas son tóxicos. La disposición inadecuada del pesticida sobrante, mezcla de pulverización, o líquido de enjuague (rinsate), constituye una violación de la Ley Federal. Si estos residuos no pueden ser eliminados de acuerdo con las instrucciones de la etiqueta, contacte la agencia estatal de plaguicidas o de control ambiental de su estado, o el representante para el Manejo de Residuos Peligrosos de la Oficina Regional de la EPA más cercana para obtener instrucciones sobre métodos adecuados de eliminación.



(continuado)

ALMACENAMIENTO Y DESECHO (continuado)

MANIPULACIÓN DE ENVASES:

Menos o igual a 5 galones: Envase no rellenable. **NO** reutilice ni rellene este envase. Ofrezcalo para reciclaje si está disponible. Enjuague tres veces el envase (o equivalente) inmediatamente después de vaciarlo. Enjuague tres veces siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezcla, y drene por 10 segundos después de que el flujo empiece a gotear. Llene el envase hasta 1/4 de su capacidad con agua y tape de nuevo. Agite por 10 segundos. Vierta el líquido de enjuague (rinsate) en el equipo de aplicación o tanque de mezcla, o almacene el enjuague para uso posterior o eliminación. Drene por 10 segundos después de que el flujo empiece a gotear. Repita este procedimiento dos veces más. Luego ofrézcalo para reciclaje si está disponible o el reacondicionamiento si es adecuado, o perfore y deseche en un área adecuada para desechos sanitarios o por la incineración.

Más de 5 galones: Envase rellenable. Rellene este envase sólo con pesticida (plaguicida). **NO** reutilice este envase para ningún otro propósito. La limpieza del envase antes de su eliminación final es la responsabilidad de la persona que se deshace del envase. La limpieza antes de rellenarlo es responsabilidad de la persona que lo rellena. Para limpiar el envase antes de su eliminación final, vacíe el contenido restante de este envase en el equipo de aplicación o en el tanque de mezcla. Llene el envase con un 10% de agua. Agite vigorosamente o haga recircular el agua con la bomba durante 2 minutos. Vierta o bombee el líquido de enjuague en el equipo de aplicación o en el sistema de recolección de líquido de enjuague. Repita este procedimiento dos veces más.

Para contenedores y mini-contenedores a granel: Envase rellenable. Rellene este envase sólo con pesticida (plaguicida). **NO** reutilice este envase para ningún otro propósito. La limpieza del envase antes de su eliminación final es la responsabilidad de la persona que se deshace del envase. La limpieza antes de rellenarlo es responsabilidad de la persona que lo rellena. Para limpiar el envase antes de su eliminación final, vacíe el contenido restante de este envase en el equipo de aplicación o en el tanque de mezcla. Llene el envase con un 10% de agua. Agite vigorosamente o haga recircular el agua con la bomba durante 2 minutos. Vierta o bombee el líquido de enjuague en el equipo de aplicación o en el sistema de recolección de líquido de enjuague. Repita este procedimiento dos veces más. Luego ofrézcalo para reciclaje si está disponible o perfore y deseche en un área adecuada para desechos sanitarios, o por otros procedimientos aprobados por las autoridades estatales y locales o mediante incineración.

¡EL ENVASE NO ES SEGURO PARA ALIMENTOS, FORRAJES, NI AGUA POTABLE!

NOTES

NOTES

NOTES

GAMMA-CYHALOTHRIN **GROUP 3A** **INSECTICIDE**

Gamla

For both indoor and outdoor use.

For Use In, On and Around Buildings and Structures for the Control of Listed Pests: on Non-Residential Lawns and Turf, Ornamental Trees and Shrubs Around Non-Residential, Institutional, Commercial, Agricultural, and Industrial Buildings; Livestock/Poultry Housings and Pet Kennels.

ACTIVE INGREDIENT:

Gamma-cyhalothrin: Cyclopropanecarboxylic acid, 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl, cyano(3-phenoxyphenyl)methyl ester*

WT. BY %

..... 5.9%

OTHER INGREDIENTS:

..... 94.1%

TOTAL: **100.0%**

Contains 0.5 lbs. of active ingredient per gallon.

Contains petroleum distillate.

*Synthetic pyrethroid, CAS No. 76703-62-3, capsule suspension (microencapsulated).

KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand this label, find someone to explain it to you in detail.)

See label booklet for complete First Aid, Precautionary Statements, Directions For Use, and Storage and Disposal.

EPA Reg. No. 83529-328

EPA Est. No. (AG) 72159-GA-001; (MC) 89332-GA-001; (SC) 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Manufactured For:

Sharda USA LLC 

2717 Lancaster Pike, Suite A, Hockessin, Delaware 19707

Net Contents: **1 Quart**

OPEN HERE 