

Conduct

HERBICIDE

For Use on Residential and Non-Residential Turfgrass to Control Specified Broadleaf and Grass Weeds.

ACTIVE INGREDIENT:	WT. BY %
Dimethylamine Salt of Quinclorac: 3,7-dichloro-8-quinolinecarboxylic acid	18.92%
OTHER INGREDIENTS:	81.08%
TOTAL:	100.00%
Contains 1.50 lbs./gal. of quinclorac: 3,7-dichloro-8-quinolinecarboxylic acid equivalent.	

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

See inside booklet for complete Precautionary Statements,
Directions For Use, and Storage and Disposal.

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 83529-172

EPA Est. No. **AG** 72159-GA-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: **1 Gallon**

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, made of butyl rubber \geq 14 mils, natural rubber \geq 14 mils, neoprene rubber \geq 14 mils, or nitrile rubber \geq 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Keep out of lakes, ponds, and streams. **Do not** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. **Do not** contaminate water by cleaning of equipment or disposal of rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with this labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the specified area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Sharda USA LLC does not recommend or authorize the use of this product in manufacturing, processing, or preparing custom blends with other products for application to turfgrass. **Do not** use to formulate or reformulate any other pesticide product that is not registered by EPA.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any butyl rubber \geq 14 mils, natural rubber \geq 14 mils, neoprene rubber \geq 14 mils, or nitrile rubber \geq 14 mils
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses.

Professional applications to residential and non-residential turfgrass (excluding sod farms) are not within the scope of the Worker Protection Standard.

Do not enter or allow others to enter the treated area until sprays have dried.

PRODUCT INFORMATION

Conduct may be applied post-emergence to residential and non-residential turfgrass (refer to **Table 1**) for the control of many broadleaf and grass weeds on the following sites: Airports, Athletic fields, Cemeteries, Golf Courses, Grounds or lawns (around Residential and Commercial Establishments), Houses of Worship, Military Institutions, Multifamily Dwellings, Parks, Picnic Grounds, Roadsides, Schools, and Sod Farms.

Restrictions:

- In New York, **Conduct** can only be applied as a spot application.
- **DO NOT** apply more than 128 fl. oz. (1.5 lbs. ae) of **Conduct** per acre (or 2.9 fl. oz. per 1,000 sq. ft.) in 1 year.
- **DO NOT** apply more than 64 fl. oz. (0.75 lb. ae) of **Conduct** per acre (or 1.45 fl. oz. per 1,000 sq. ft.) in a single application.
- **DO NOT** apply to golf course collars or greens.
- **DO NOT** make applications of **Conduct** to drought-stressed turfgrass and/or drought-stressed weeds.
- **DO NOT** apply to fine fescue unless it is part of a seed blend.
- **DO NOT** apply to Bahiagrass, carpetgrass, centipedegrass, colonial and seaside bentgrass, dichondra, St. Augustinegrass, or lawns or turfgrass where desirable clovers are present.
- **DO NOT** apply to exposed feeder roots of trees or ornamentals or within the dripline of trees and other ornamental species.
- **DO NOT** apply into any ornamental bed.
- **DO NOT** apply within 4 weeks after seedling emergence of creeping bentgrass, fine fescue blends, Kentucky bluegrass, and perennial ryegrass.
- **DO NOT** apply **Conduct** prior to and within 2 weeks after seeding seashore paspalum.
- **DO NOT** use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- **DO NOT** plant eggplants or tobacco within 12 months in areas treated with **Conduct**.
- **DO NOT** plant tomatoes or carrots within 24 months in areas treated with **Conduct**.
- Apply as a medium or coarser spray (ASABE standard 572).
- **DO NOT** release spray at a height greater than 30 inches above the ground.
- **DO NOT** apply when wind speeds are greater than 10 mph at the application site.
- Use a lawn-type sprayer with coarse spray because wind drift is less likely.
- Avoid mist and spray onto vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants, especially plants belonging to the *Solanaceae* family, such as tomatoes, eggplants, and bell peppers.
- **DO NOT** discard rinsate on or near desirable plants.
- **DO NOT** apply this product by air or through any type of irrigation system or equipment.

RESISTANCE MANAGEMENT

For resistance management, **Conduct** is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to **Conduct** and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of **Conduct** or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout for herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species;
 - Surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance management and/or integrated weed-management recommendations for specific crops and weed biotypes.

APPLICATION INFORMATION

In New York, **Conduct** can only be applied as a spot application.

Apply **Conduct** to actively growing weeds as post-emergence broadcast or spot sprays using the turfgrass species, rate, and growth stages indicated in **Table 1**, **Table 2**, and **Table 3**. **DO NOT** exceed the labeled application rate or fail to comply with use restrictions listed in the above **Restrictions** section, under **PRODUCT INFORMATION**.

For best results, weeds should not be under stress from lack of water, excessive water, low fertility, mowing shock, excessive hot or cold temperatures, or injury from other herbicide applications.

To achieve consistent weed control, use methylated seed oil (MSO). Refer to **Table 2** and **Table 3** for rates.

Adding adjuvants may cause slight leaf burn, but new growth is normal, and turfgrass vigor is not reduced. The potential for leaf burn is increased when relative humidity and temperature are high. Additional stress from low mowing heights may also increase the possibility of turfgrass injury. Chelated iron or sprayable soluble nitrogen fertilizer will reduce a slight yellowing that may occur on some turfgrass species. Not all chelated iron or sprayable nitrogen fertilizers are compatible with **Conduct**. Always perform a compatibility test to ensure proper mixing. See **Compatibility Test For Mix Components** section of label for directions.

Broadcast Applications

Apply with properly calibrated ground equipment in sufficient water per acre to provide uniform spray distribution (at least 20 gallons of water per acre or at least 0.5 gallon per 1,000 sq. ft.). Use low-pressure sprayers at 20 - 40 PSI.

Maintain continuous agitation during spraying with good mechanical or bypass agitation. Nozzle screens must be no finer than 50-mesh (100-mesh is finer than 50-mesh). Check sprayer routinely to determine proper calibration. Flat fan, flood, or cone nozzles may be used. Arrange nozzles for uniform coverage for turfgrass and weeds to be controlled. Adjust boom height, nozzle selection, and pressure to provide uniform coverage and minimize spray drift.

Avoid overlaps that will increase rates above those labeled for use. Avoid application when winds may cause drift.

DO NOT apply when wind speeds are greater than 10 mph at the application site.

Spot Applications

Post-emergence spot applications may be made to susceptible weeds in turfgrass that is tolerant to **Conduct** (see **Table 1** and **Table 2**). Apply 1.45 fl. oz. (0.75 lb. ae/A) of **Conduct** per 1,000 sq. ft. of treated area. See **Table 5** for spot spray mix instructions.

Spot Treatment in New York: Spray individual weeds only. Adjust the sprayer to coarse spray to minimize wind drift. Apply to the center of the weed and spray to lightly cover.

Mowing Information

DO NOT mow 2 days before or after applying **Conduct** to maximize weed control and minimize potential turfgrass injury. Clippings from the first 3 mowings after application must be left on the treated area.

Irrigation and Rainfall

If soil moisture is not sufficient prior to **Conduct** application, irrigation may improve weed control. For best results, **DO NOT** water or irrigate for 24 hours after application. If rainfall does not occur in 2 - 7 days after application, irrigation of at least half inch is desirable.

Extended Grass Control

To extend grass control, **Conduct** can be tank mixed with Pendulum® 3.3 EC herbicide (EPA Reg. No. 241-341, pendimethalin) or Pendulum® AquaCap™ herbicide (EPA Reg. No. 241-416, pendimethalin) to provide residual control of annual grasses. Consult the respective tank mix labels for additional weeds controlled.

Seeding/Overseeding/Sprigging

The use of **Conduct** before or after seeding or overseeding a turfgrass area will not significantly interfere with the turfgrass seed germination and growth of those grass types identified as tolerant or moderately tolerant in **Table 1**. See **Table 4** for seeding, overseeding, or sprigging application timing.

ADJUVANTS

Additives in Spray Mix to Achieve Control

Methylated seed oil is the preferred adjuvant for post-emergence applications. However, if an MSO is not available in your region, the use of a crop oil concentrate (COC) or other high quality surfactant must be used in the spray tank at the time of application. Refer to actual product label for use rates and use directions.

Additives must not be used when tank mixing with emulsifiable concentrate (EC) products or turfgrass phytotoxicity may occur.

Methylated seed oil or crop oil concentrate used as the adjuvant with **Conduct** must meet all the following criteria:

- Nonphytotoxic
- Contain only EPA-exempt ingredients
- Provide good mixing quality in the jar test
- Successful in local experience

The exact composition of suitable products will vary; however, any methylated seed oil or crop oil concentrate used should contain emulsifiers to provide good mixing quality.

When an adjuvant (or a specific adjuvant product, such as a drift control agent) is to be used with this product, the use of a Chemical Producers and Distributors Association (CPDA) certified adjuvant is recommended.

Consult your local Sharda USA LLC representative or distributor for instructions for your area.

TURFGRASS TANK MIXES

Read and follow the applicable Restrictions and Limitations and Directions For Use on all products involved in tank mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

To increase spectrum of control of broadleaf weed species, a tank mix with 2,4-D; triclopyr; or other broadleaf herbicides may be used. For extended residual control, apply **Conduct** with Pendulum® 3.3 EC herbicide or Pendulum® AquaCap™ herbicide.

For sedge control, applications of **Conduct** with Basagran® T&O herbicide (EPA Reg. No. 7969-326, sodium bentazon), Image® 70 DG herbicide (EPA Reg. No. 241-319, imazaquin), or MSMA may be made. Combinations with MSMA will aid in control of certain grass weeds, such as Bahiagrass or kikuyugrass. Consult labels for turfgrass tolerance when tank mixing. Make separate applications if all target weeds are not at the correct growth stage for treatment at the same time.

Physical incompatibility, reduced weed control, or turfgrass injury may result from mixing **Conduct** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers.

Before tank mixing, a simple jar test is required to ensure compatibility of herbicides or other pesticides and/or additives.

Cleaning Spray Equipment

Clean application equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product.

Compatibility Test for Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of labeled rate per acre:

1. Water - For 20 gallons per acre spray volume, use 3.3 cups (800 mL) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
2. Products in PVA bags - Cut an opening in the water-soluble PVA bag just large enough to use a teaspoon for measuring purposes. Use the opened, water-soluble PVA bag first when preparing spray solution. Boron-containing fertilizers can be incompatible with PVA material. Include PVA material if a boron fertilizer is intended to be used. Cap the jar and invert 10 cycles.
3. Water-dispersible products (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions) - Cap the jar and invert 10 cycles.
4. Water-soluble products (**Conduct**) - Cap the jar and invert 10 cycles.
5. Emulsifiable concentrates (methylated seed oil) - Cap the jar and invert 10 cycles.
6. Water-soluble additives - Cap the jar and invert 10 cycles.

Let the solution stand for 15 minutes.

Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface nor thick (clabbered) texture. For water-dispersible granule (WG) or wettable powder (WP) products, a fine precipitate that is easily resuspended is normal; large, non-dispersible particles (>300 microns) that precipitate on standing are a sign of tank mix incompatibility. Do not use any spray solution that could clog spray nozzles.

Mixing Instructions - Mixing Order

1. Water - Begin by agitating a thoroughly clean sprayer tank 3/4 full of clean water.
2. Agitation - Maintain constant agitation throughout mixing and application.
3. Inductor - If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA bags - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water-dispersible products (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
6. Water-soluble products (such as **Conduct**)
7. Emulsifiable concentrates (such as methylated seed oil or crop oil concentrate)
8. Water-soluble additives (such as chelated iron or soluble nitrogen fertilizer when applicable; not all chelated iron or sprayable nitrogen fertilizers are compatible with **Conduct**. Always perform a compatibility test to ensure proper mixing. See **Compatibility Test for Mix Components** section above for directions.)
9. Remaining quantity of water.

Maintain constant agitation during application.

Backpack Sprayer

Begin with a clean spray tank. Fill the spray tank 1/2 full with clean water and add the required amount of **Conduct** to the sprayer. Cap sprayer and agitate to ensure mixing. Uncap sprayer and add appropriate amount of methylated seed oil. Cap sprayer and agitate once again. Uncap sprayer and finish filling tank to desired level. During application, agitate the mixture on occasion to ensure mixing. If the mixture is allowed to settle for any period of time, thorough agitation is essential before spraying is resumed.

APPLICATION DIRECTIONS

Table 1. Turfgrass Tolerance (Established)

Highly Tolerant	Moderately Tolerant	Susceptible
Bermudagrass, Common*	Bentgrass, Creeping*	Bahiagrass
Bluegrass, Annual	Bermudagrass, Hybrid*	Bentgrass, Colonial
Bluegrass, Kentucky	Bluegrass, Rough (<i>Poa Trivialis</i>)	Bentgrass, Seaside
Buffalograss	Fescue, Chewing's	Carpetgrass
Fescue, Tall	Fescue, Fine**	Centipedegrass
Ryegrass, Annual	Fescue, Hard	Dichondra
Ryegrass, Perennial	Fescue, Red	St. Augustinegrass
Zoysiagrass	Paspalum, Seashore	

*Yellowing that occurs on these species can be reduced by the addition of chelated iron or sprayable soluble nitrogen fertilizer. See **APPLICATION INFORMATION** and **ADJUVANTS**.

Apply **Conduct to fine fescue only when it is part of a blend.

Restriction:

- **DO NOT** use on golf course greens and collars.

Table 2. Conduct Application to Established Creeping Bentgrass

Turfgrass Species	Application Rate/Timing	Additive Rate
Bentgrass, Creeping* ¹	Conduct must be applied in 2 - 3 split applications at 0.5 - 1.0 fl. oz. per 1,000 sq. ft. (0.25 - 0.51 lb. ae/A). Time sequential application(s) 14 - 21 days apart.	Methylated seed oil at 0.55 fl. oz. per 1,000 sq. ft. (1.5 pts./A)
Bentgrass, Creeping* ²	Conduct must be applied in 2 to 3 split applications at 0.65 - 1.0 fl. oz. per 1,000 sq. ft. (0.33 - 0.51 lb. ae/A). Time sequential application(s) 14 - 21 days apart.	
<p>*Yellowing that occurs on these species can be reduced by the addition of chelated iron or sprayable soluble nitrogen fertilizer. See APPLICATION INFORMATION and ADJUVANTS. ¹DO NOT apply this rate in California. ²This rate range for use only in California.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT exceed 128 fl. oz. (1.5 lbs. ae) of Conduct per acre (2.9 fl. oz. per 1,000 sq. ft.) per year. • DO NOT use on golf course greens and collars. 		

Table 3. Application Rates and Timing for Post-Emergence Weed Control in Turfgrass*

Weed Species Controlled	Conduct Rate	Additive Rate
Grass Weeds	Broadcast Application	Methylated seed oil at 0.55 fl. oz. per 1,000 sq. ft. (1.5 pts./A)
Barnyardgrass (<i>Echinochloa crus-galli</i>)	64 fl. oz. per acre or	
Crabgrass, Large (<i>Digitaria sanguinalis</i>) ^{1,4}	1.45 fl. oz. per 1,000 sq. ft. (0.75 lb. ae/A)	
Crabgrass, Smooth (<i>Digitaria ischaemum</i>) ^{1,4}		
Foxtail, Giant (<i>Setaria faberii</i>)	Spot Application	
Foxtail, Green (<i>Setaria viridis</i>) ¹	1.45 fl. oz. per 1,000 sq. ft. of treated area	
Foxtail, Yellow (<i>Setaria glauca</i>) ¹	(0.75 lb. ae/A)	
Kikuyugrass (<i>Pennisetum clandestinum</i>) ^{2,3}	Refer to footnotes in Table 2 and Table 3 for specific turfgrass or weed instructions.	
Signalgrass, Broadleaf (<i>Brachiaria platyphylla</i>) ¹		
Torpedograss (<i>Panicum repens</i>) ³		
<p>*New York: Conduct can only be applied as a spot application. ¹Under certain conditions, Conduct application to annual grasses at 2- to 4-tiller may not provide complete control. A sequential application will be required for grass control in these situations. Optimum control is achieved when applications of Conduct + methylated seed oil are applied either before second tiller or as grass weeds mature. ²Tank mix partner or sequential application required. ³Make 2 sequential applications of 1.0 fl. oz. (0.51 lb. ae/A) of Conduct per 1,000 sq. ft. and an additional sequential application up to 0.90 fl. oz. (0.46 lb. ae/A) of Conduct per 1,000 sq. ft. at 14- to 21-day intervals. ⁴Biotypes of large and smooth crabgrass in California have shown varied response to Conduct. If control failure occurs following a full or split application, do not reapply Conduct. Change to a herbicide with a different mode of action.</p>		

Table 3. Application Rates and Timing for Post-Emergence Weed Control in Turfgrass* (continued)

Weed Species Controlled	Conduct Rate	Additive Rate
Broadleaf Weeds		
Bindweed, Field (<i>Convolvulus arvensis</i>)	<p>Broadcast Application 64 fl. oz. per acre or 1.45 fl. oz. per 1,000 sq. ft. (0.75 lb. ae/A)</p> <p>Spot Application 1.45 fl. oz. per 1,000 sq. ft. of treated area (0.75 lb. ae/A)</p> <p>Refer to footnotes in Table 2 and Table 3 for specific turfgrass or weed instructions.</p>	<p>Methylated seed oil at 0.55 fl. oz. per 1,000 sq. ft. (1.5 pts./A)</p>
Clover, Hop (<i>Trifolium aureum</i> Pollich)		
Clover, Red (<i>Trifolium pratense</i>)		
Clover, White (<i>Trifolium repens</i>)		
Daisy, English (<i>Bellis perenne</i>) ^{2,5}		
Dandelion, Common (<i>Taraxacum officinale</i>) ²		
Dollarweed (<i>Hydrocotyle umbellata</i>)		
Geranium, Carolina (<i>Geranium carolinianum</i>)		
Horseweed (<i>Conyza canadensis</i>) ²		
Medic, Black (<i>Medicago lupulina</i>)		
Morningglory spp. (<i>Ipomoea</i> sp.)		
Speedwell, Common (<i>Veronica officinalis</i>)		
Speedwell, Slender (<i>Veronica filiformis</i>)		
Speedwell, Thymeleaf (<i>Veronica serpyllifolia</i>)		
Violet, Wild (<i>Viola</i> sp.)		
<p>*New York: Conduct can only be applied as a spot application. ²Tank mix partner or sequential application required. ⁵Not for use to control this weed in California.</p>		

Table 4. Seeding/Overseeding/Sprigging Timing Chart*

Variety	Before Seeding ¹	At-Seeding	7 Days After Emergence	14 Days After Emergence	28 Days After Emergence
Annual Bluegrass	OK	OK	OK	OK	OK
Annual Ryegrass	OK	OK	OK	OK	OK
Buffalograss	OK	OK	OK	OK	OK
Common Bermudagrass ² (for sprigging see footnote 2)	OK	OK	OK	OK	OK
Creeping Bentgrass	OK	NO	NO	NO	OK
Fine Fescue (in blend)	OK	NO	NO	NO	OK
Hybrid Bermudagrass ² (for sprigging see footnote 2)	OK	OK	OK	OK	OK
Kentucky Bluegrass	OK	NO	NO	NO	OK
Perennial Ryegrass	OK	OK	NO	NO	OK
Seashore Paspalum ^{2,3} (for sprigging see footnote 2)	NO	NO	NO	OK	OK
Tall Fescue	OK	OK	OK	OK	OK
Zoysiagrass ² (for sprigging see footnote 2)	OK	OK	OK	OK	OK

*No adjuvant or additive should be used when **Conduct** applications are made on newly emerged turfgrass seedlings until 28 days after emergence. With the exception of seashore paspalum, a **Conduct** application rate of 1.45 fl. oz./1,000 sq. ft. (0.75 lb. ae/A) can be made to all other turfgrass species in **Table 4**.

¹**Conduct** can be applied 7 days or greater prior to seeding.

²**Conduct** can be used any time prior to, at or after sprigging as indicated by turfgrass species in **Table 4**.

³0.75 fl. oz. - 1.45 fl. oz./1,000 sq. ft. (0.37 - 0.75 lb. ae/A) application can be made at times indicated in **Table 4**.

Time **Conduct** application around the seeding operations using the chart above as a reference point.

Table 5. Spot Spraying with Conduct

*Spray Mix Volume (Gallons)	Conduct Product in Mix (Tablespoons)	MSO Adjuvant in Mix (Tablespoons)
1	3	1.5
2	6	3.0
3	9	4.5

*Apply at the rate of 1 gallon per 1,000 sq. ft.

1 tablespoon = 0.5 fl. oz. (0.25 lb. ae/A) of **Conduct** product.

Notes: For consistent results, make **Conduct** application to newly germinated crabgrass, to 1-tiller crabgrass, and when crabgrass has matured to 5-tillers or greater. Under certain conditions, applications of **Conduct** made to annual grasses 2-tiller to 4-tiller may not provide complete control. A sequential application will be required for grass control in these situations.

Spot Treatment in New York: Spray individual weeds only. Adjust the sprayer to coarse spray to minimize wind drift. Apply to center of the weed and spray lightly to cover.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry, well-ventilated area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Less Than or Equal to 5 Gallons. Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

Greater Than 5 Gallons. Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times.

Greater Than 5 Gallons. Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration.

For Bulk and Mini-Bulk Containers. Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

NOTES

NOTES

QUINCLORAC GROUP 4 HERBICIDES

Conduct

For Use on Residential and Non-Residential Turfgrass to Control Specified Broadleaf and Grass Weeds.

ACTIVE INGREDIENT:	WT. BY %
Dimethylamine Salt of Quinclorac:	
3,7-dichloro-8-quinolinecarboxylic acid	18.92%
OTHER INGREDIENTS:	81.08%
TOTAL:	100.00%

Contains 1.50 lbs./gal. of quinclorac:
3,7-dichloro-8-quinolinecarboxylic acid equivalent.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID - IF SWALLOWED: • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. • **DO NOT** give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice. **IF IN EYES:** • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. **IF INHALED:** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER - Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at **1-800-222-1222**.

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION - Harmful if swallowed. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS - This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Keep out of lakes, ponds, and streams. **Do not** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. **Do not** contaminate water by cleaning of equipment or disposal of rinseate.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry, well-ventilated area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER HANDLING: Less Than or Equal to 5 Gallons.** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

See label booklet for additional
Precautionary Statements and Directions For Use.

Manufactured For: Sharda USA LLC

7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 83529-172 EPA Est. No. [AG] 72159-GA-001;

[MC] 89332-GA-001; [SC] 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: **1 Gallon**

OPEN HERE