

Sequence SC

FUNGICIDE

For control of certain diseases of Almonds, Apples, Apricots*, Bananas*, Beech nuts*, Brazil nuts*, Butternuts*, Cashews*, Cherries, Chestnuts*, Chinquapins*, Filberts*, Hickory nuts*, Macadamia nuts*, Nectarines, Peaches, Peanuts*, Pears, Pecans*, Plums*, Plums-chickasaw*, Plums-damson*, Plums-Japanese*, Plumcots*, Prune plums*, and Walnuts.

*Not registered for use by California.

ACTIVE INGREDIENT:	WT. BY %
Dodine (n-dodecylguanidine acetate)	39.6%**
OTHER INGREDIENTS:	60.4%
TOTAL:	100.0%

**Equivalent to 3.4 pounds Dodine per gallon

KEEP OUT OF REACH OF CHILDREN / MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

See label booklet for complete Precautionary Statements, Directions for Use, and Storage and Disposal.

FIRST AID - IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to by a poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBERS - For general information about this product, contact the National Pesticides Information Center (NPIC) at **1-800-858-7378**, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>. Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at **1-800-222-1222**.

NOTE TO PHYSICIAN - Probable mucosal damage may contraindicate the usage of gastric lavage.

PRIMEROS AUXILIOS - SI ENTRA EN CONTACTO CON LOS OJOS: Mantenga los ojos abiertos y enjuáguelos lenta y cuidadosamente con agua, durante 15 a 20 minutos. Si utiliza lentes de contacto, retírelos después de los primeros 5 minutos; luego continúe enjuagando los ojos. Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento. **SI CAE EN LA PIEL O LA ROPA:** Quítelse la ropa contaminada. Enjuague la piel inmediatamente con bastante agua por 15 - 20 minutos. Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento. **SI SE INGERIE:** Llame de inmediato a un centro de control de envenenamientos o a un médico para consejo de tratamiento. Si la persona puede tragar, haga que beba un vaso de agua lentamente. **NO** induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. **NO** administre nada por boca a una persona que haya perdido el conocimiento. **SI ES INHALADO:** Trásfale a la persona al aire fresco. Si la persona no está respirando llame al 911 o a una ambulancia, luego dé respiración artificial, preferiblemente de boca a boca, si es posible. Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.

NÚMERO DE TELÉFONO DIRECTO - Para información general sobre este producto, contacte al Centro Nacional de Información de Pesticidas (NPIC, por sus siglas en inglés) a **1-800-858-7378**, Lunes a Viernes, 8 AM a 12 PM PST (hora del Pacífico), o a <http://npic.orst.edu>. Cuando llame a un centro de control de envenenamiento, o a un médico, o intente obtener tratamiento, tenga a la mano el recipiente o la etiqueta del producto. Para información emergencia sobre este producto, llame al centro de control de envenenamientos a **1-800-222-1222**.

NOTA AL MÉDICO - El probable daño de las mucosas puede contraindicar el uso de un lavado gástrico.

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 82633-104-83529

EPA Est. No. **AG** 72159-GA-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER/PELIGRO

Corrosive. Causes irreversible eye damage. Causes skin irritation. **DO NOT** get in eyes, on skin, or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid breathing spray mist or vapor. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate or viton.

Applicators mixers, loaders, applicators, and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material (barrier laminate, butyl rubber $\geq$ 14 mils, nitrile rubber $\geq$ 14 mils, neoprene rubber $\geq$ 14 mils, polyvinyl chloride (PVC) $\geq$ 14 mils, and/or viton $\geq$ 14 mils)
- Goggles or face shield
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron for mixers and loaders

Mixers, loaders, and applicators for mechanically pressurized handguns to bananas, olives, Pome Fruits – Crop Group 11-10, Stone Fruits – Crop Group 12-12, and Tree Nuts – Crop Group 14-12 must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Mixers and loaders for aerial applications to bananas, olives, Pome Fruits – Crop Group 11-10, Stone Fruits – Crop Group 12-12, and Tree Nuts – Crop Group 14-12 must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

See **ENGINEERING CONTROLS** for additional requirements.

USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.249(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Surface Water and Erosion Control Statement: This product may contaminate water through runoff and has the potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated including plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants,
- Chemical-resistant gloves made of any waterproof material, chemical-resistant footwear plus socks,
- Protective eyewear, and
- Chemical-resistant headgear (if overhead exposure).

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

USE RESTRICTIONS

- **DO NOT** use this product for any crop or any pest not specifically listed on this label. For crops and pests listed on the label, follow closely the listed application timings.
- **DO NOT** apply at any application timing not listed on the label.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply in greenhouses.
- **DO NOT** graze livestock in treated fields, orchards or treated groves.

FUNGICIDE RESISTANCE MANAGEMENT

For resistance management, **Sequence SC** contains a Group U12 fungicide. Any fungal population may contain individuals naturally resistant to **Sequence SC** and other Group U12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides/bactericides are used repeatedly in the same fields. Appropriate resistance management strategies must be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of **Sequence SC** or other Group U12 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Sharda USA LLC at <http://www.shardausa.com/>. You can also contact your pesticide distributor or university extension specialist to report resistance.

DILUTE AND CONCENTRATE SPRAYS

Sequence SC may be used effectively in dilute and concentrate sprays when applied properly to deliver the labeled rate (pints per acre). Trees must be covered uniformly. **DO NOT** excessively drench fruit and foliage with dilute sprays. Avoid heavy deposits when using concentrate sprays.

Use **Sequence SC** in a minimum of 15 gallons of water per acre for all ground applications, and in a minimum of 10 gallons of water per acre for all air applications.

INSTRUCTIONS FOR MIXING

Add **Sequence SC** directly to the spray tank when half filled with water. Keep agitator running during filling and spraying.

COMPATIBILITY

Experience indicates that **Sequence SC** can be used effectively at specified dosages with most of the commonly used fungicides, herbicides, insecticides and acaricides. However, some incompatibilities were identified, especially with some copper, oil, sulfur, foliar nutrients and chlorpyrifos formulations. Therefore, we always advise testing the compatibility of the products used in combination with **Sequence SC** at small scale before spraying or to ask your local distributor or consultant about potential incompatibility.

Observe all precautions, restrictions and limitations on labeling of all products used in mixtures. Follow the most restrictive label limitations and precautions. No label dosage rates may be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity), and method of application (e.g., ground, aerial airstream) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product.

WIND SPEED

DO NOT apply at wind speeds greater than 10 mph at the application site.

DROPLET SIZE

Apply as a medium or coarser spray (ASAE standard 572), except when making orchard airblast applications. For orchard airblast applications, apply at the largest droplet size that provides sufficient coverage and control.

GROUND APPLICATION

Release Height - **DO NOT** release spray at a height greater than 4 feet above the ground or crop canopy.

AIRBLAST

Sprays must be directed into the crop canopy. Turn off outward pointing nozzles at row ends and when spraying outer rows.

AERIAL APPLICATION

- **Release Height** - **DO NOT** release spray at a height greater than 10 feet above the ground or crop canopy.
- **Boom Length** - The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- **Swath Adjustment** - When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for the displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Leave at least one swath unsprayed at the downwind edge of the treated field.

CROP-SPECIFIC INSTRUCTIONS**ALMONDS, BEECH NUTS*, CHESTNUTS*, AND CHINQUAPINS***

***Not registered for use by California**

DISEASE	APPLICATION DIRECTIONS	USE RATE
Brown rot blossom blight – (<i>Monilinia laxa</i> / <i>Monilinia fructicola</i>)	For disease suppression at pink bud, full bloom, and petal fall.	3 pints/A (1.3 lbs. a.i./acre)
Scab (<i>Venturia carpophila anamorph</i> , <i>Caldosporium carpophilum</i>)	Use at petal fall. Repeat application 2 and 5 weeks later.	
Shot hole (<i>Wilsonomyces carpophilus</i>)	Use at leaf emergence (bloom through petal fall). Repeat application 2 and 5 weeks later.	
Anthraxnose (<i>Gnomonia leptostyla</i>)	Use when first leaves unfold. Repeat application at 10- to 14-day intervals.	

RESTRICTIONS:

- **DO NOT** apply more than 9 pints of **Sequence SC** per acre per year (3.9 lbs. a.i./acre/year).
- **DO NOT** apply more than 3 pints (1.3 lbs. a.i./acre) of **Sequence SC** per acre per application.
- **DO NOT** apply more than 3 applications per year.
- **DO NOT** apply less than 10 days from the last application.
- **DO NOT** apply later than 5 weeks after petal fall.

APPLES

DISEASE	APPLICATION DIRECTIONS
Scab (<i>Venturia inaequalis</i>)	Use 1.5 pints per acre of Sequence SC beginning at green tip, and 7 days later to clean the orchard of over-wintering scab at the start of the season. Alternatively, apply Sequence SC preventively at weekly intervals from green tip to petal fall. DO NOT apply more than 2 applications per year. For resistance management, it is advised to tank-mix Sequence SC with a fungicide including captan or mancozeb based formulation, at labeled rates. When tank-mixing Sequence SC with a fungicide including captan or mancozeb, fill the spray tank with clean water half of final desired volume. Start agitation. Slowly add Sequence SC at the specified rate. Once mixed, slowly add a fungicide including, captan or mancozeb based formulation, at the specified rate. Add water to desired final volume. Mix thoroughly and apply at once. Maintain sufficient agitation during mixing and spraying operations to ensure a uniform spray mixture.

RESTRICTIONS:

- **DO NOT** apply more than 1.5 pints (0.65 lb. a.i./acre) of **Sequence SC** per acre in any single application.
- **DO NOT** apply more than 3 pints (1.3 lbs. a.i./acre) of **Sequence SC** per acre per year.
- **DO NOT** apply more than 2 applications per year.
- **DO NOT** apply less than 7 days from the last application.
- **DO NOT** apply after petal fall.

APRICOTS*, NECTARINES, AND PEACHES

***Not registered for use by California**

DISEASE	APPLICATION DIRECTIONS	USE RATE
Leaf Curl (<i>Taphrina deformans</i>)	Use as a dormant or delayed dormant application. Additional applications may be made just before buds swell in the spring.	3 pints/A (1.3 lbs. a.i./acre)
Brown rot blossom blight (<i>Monilinia laxa</i> / <i>Monilinia fructicola</i>) (States West of the Mississippi River)	For disease suppression , use at pink bud. Repeat application at full bloom and again at petal fall.	
Scab (<i>Venturia carpophila anamorph</i> ; <i>Cladosporium carpophilum</i>)	Use at petal fall.	
Shot hole (<i>Wilsonomyces carpophilus</i>)	Use at dormant state, and repeat application at petal fall.	
RESTRICTIONS: • DO NOT apply more than 9 pints of Sequence SC per acre per year (3.9 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence SC per acre per application. • DO NOT apply more than 3 applications per year. • DO NOT apply less than 7 days from the last application. • DO NOT apply after petal fall.		

BANANAS*

***Not registered for use by California**

DISEASE	APPLICATION DIRECTIONS
Sigatoka	Air application: Use 0.85 to 1.5 pints per acre of Sequence SC as needed to maintain Sigatoka control in a mixture of 19 to 25 pints water/acre and 4 pints mineral oil/acre. Repeat application thereafter at 7- to 15-day interval. It is advised for subsequent cover sprays to use Sequence SC in rotation with other labeled fungicides that have a different mode of action.
RESTRICTIONS: • DO NOT apply more than 7.5 pints of product per acre per year (3.2 lbs. a.i./acre/year). • DO NOT apply more than 1.5 pints (0.65 lb. a.i./acre) of Sequence SC per acre per application. • DO NOT apply more than 5 applications per year. • DO NOT apply less than 7 days from the last application.	

BRAZIL NUTS*, BUTTERNUTS*, CASHEWS*, FILBERTS*, HICKORY NUTS*, MACADEMIA NUTS*, AND WALNUTS

***Not registered for use by California**

DISEASE	APPLICATION DIRECTIONS	USE RATE
Brown rot blossom blight (<i>Monilinia laxa</i> / <i>Monilinia fructicola</i>)	For disease suppression , use at pink bud, full bloom, and petal fall.	3 pints/A (1.3 lbs. a.i./acre)
Scab (<i>Venturia carpophila anamorph</i> , <i>Cladosporium carpophilum</i>)	Use at petal fall. Repeat application 2 and 5 weeks later.	
Shot hole (<i>Wilsonomyces carpophilus</i>)	Use at leaf emergency (bloom through petal fall). Repeat application 2 and 5 weeks later.	
Anthrachnose (<i>Gnomonia leptostyla</i>)	Use when first leaves unfold. Repeat application at 10- to 14-day intervals.	
RESTRICTIONS: • DO NOT apply more than 9 pints of Sequence SC per acre per year (3.9 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence SC per acre per application. • DO NOT apply more than 3 applications per year. • DO NOT apply less than 10 days from the last application. • DO NOT apply within 7 days of harvest.		

CHERRIES (SOUR AND SWEET)

DISEASE	APPLICATION DIRECTIONS
Leaf Spot (<i>Blumeriella jaapii</i> anamorph, <i>Coccomyces hiemalis</i>)	Use 1.5 - 3 pints of Sequence SC per acre at petal fall or when first leaves unfold. Repeat application there- after at 7- to 10-day intervals. If conditions favor leaf spot, continue applications through harvest. For resis- tance management, it is advised to tank-mix with a fungicide including a captan based formulation, at labeled rate. Fill the spray tank with clean water at half of the final desired volume. Start agitation. Slowly add Sequence SC at the specified rate. Slowly add a fungicide including, captan, at the labeled rate. Add water to desired final volume. Mix thoroughly and apply at once. Maintain sufficient agitation during mixing and spraying operations to ensure a uniform spray mixture. Post-Harvest Applications to Trees: Apply 1.5 - 3 pints of Sequence SC per acre immediately after har- vest and as needed.
Brown rot blossom blight (<i>Monilinia laxa</i> / <i>Monilinia fructicola</i>) (States West of the Mississippi River)	For disease suppression, use 3 pints of Sequence SC per acre at the time flower buds begin to open (white bud-popcorn bloom stages). Repeat application at full bloom and again at petal fall. Applications of Sequence SC to cherries when temperatures and humidity are excessively high may cause spotting to foliage or fruit.
RESTRICTIONS: • DO NOT apply more than 12 pints of Sequence SC per acre per year (5.2 lbs. a.i./acre/year). • DO NOT apply more than 6 applications per year (2 applications at 3 pints of Sequence SC per acre (1.3 lbs. a.i./acre), and 4 applications at 1.5 pints of Sequence SC per acre (0.65 lb. a.i./acre)). • DO NOT apply less than 7 days from the last application. • DO NOT apply within 7 days of harvest.	

PEANUTS*

*Not registered for use by California

DISEASE	APPLICATION DIRECTIONS
Early Leaf Spot (<i>Cercospora arachidicola</i>) and Late Leaf Spot (<i>Cercosporidium personatum</i>)	Ground and Air Applications: Sequence SC alone - Use a maximum of 0.9 - 1.5 pints Sequence SC per acre in sufficient water to achieve thorough plant coverage. Begin with applications, first approximately 35 days after planting, and repeat at 10- to 14-day intervals. Sequence SC in tank-mix - When tank-mixing with white mold products including tebuconazole, dose rate can be reduced down to 0.8 pints of Sequence SC per acre. Tank-mix incompatibilities were observed with some tebuconazole based formulations. Therefore, we always advise to test the compatibility of the products used in combination with Sequence SC at small scale before spraying or to ask your local distributor about potential incompatibility. The tank-mix fungicide partner must be used at the labeled rate. Applications are made at 10- to 14-day intervals, beginning approximately 55 days after planting. DO NOT apply within 14 days of harvest.
RESTRICTIONS: • DO NOT apply more than 4.5 pints of Sequence SC per acre per year (1.95 lbs. a.i./acre/year). • DO NOT apply more than 1.5 pints (0.65 lb. a.i./acre) of Sequence SC per acre per application. • DO NOT apply more than 5 applications per year when using lower rates. • DO NOT apply less than 10 days from the last application. • DO NOT apply within 14 days of harvest. • DO NOT feed hay or vines (peanuts) to livestock.	

PEARS

DISEASE	APPLICATION DIRECTIONS
Scab (<i>Venturia pirina</i>) and leaf spot (<i>Fabraea</i> spp.)	Protection Schedule – Ground and Air application: Use 3 pints per acre of Sequence SC at 7-day intervals to maintain scab control beginning at green tip. Fill the spray tank with clean water half of final desired volume. Start agitation. Slowly add Sequence SC at the specified rate. Add water to desired final volume. Mix thoroughly and apply at once. Maintain sufficient agitation during mixing and spraying operations to ensure a uniform spray mixture.
RESTRICTIONS: • DO NOT apply more than 9 pints of Sequence SC per acre per year (3.9 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence SC per acre per application. • DO NOT apply more than 3 applications per year. • DO NOT apply less than 7 days from the last application. • DO NOT apply within 7 days of harvest.	

PECANS*

***Not registered for use by California**

DISEASE	APPLICATION DIRECTIONS
Scab, liver spot, brown leaf spot, downy leaf spot, leaf blotch, downy mildew	Ground and Air application: Sequence SC alone - Use 3 pints of Sequence SC per acre in sufficient water to obtain good coverage of spray at pre- and post-pollination periods. Apply every 10 to 21 days or as needed as determined by rainfall, from bud break through shell hardening. Sequence SC in tank-mix - In case of tank-mix with triphenyltin hydroxide or fenbuconazole use 1.5 pints of Sequence SC per acre. Sequence SC has been used effectively on most pecan varieties. Severe foliage injury has been experienced on Barton, Moore, Shawnee and Van Deman in the Southeast. Foliar bronzing has sometimes been observed on other varieties, especially when tank-mixed with certain miticides.
RESTRICTIONS: • DO NOT apply more than 18 pints of Sequence SC per acre per year (7.8 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence SC per acre per application. • DO NOT apply more than 6 applications per year. • DO NOT apply less than 10 days from the last application. • DO NOT apply after shucks have started to open.	

PLUMS*, PLUMS-CHICKASAW*, PLUMS-DAMSON*, PLUMS-JAPANESE*, PLUMCOTS*, PRUNE PLUMS*

***Not registered for use by California**

DISEASE	APPLICATION DIRECTIONS	USE RATE
Leaf Curl (<i>Taphrina deformans</i>)	Use as a dormant or delayed dormant application. Additional applications may be made just before buds swell in the spring.	3 pints/A (1.3 lbs. a.i./acre)
Brown rot blossom blight (<i>Monilinia laxa</i> / <i>Monilinia fructicola</i>) (States West of the Mississippi River)	For disease suppression , use at pink bud. Repeat application at full bloom and again at petal fall.	
Scab (<i>Venturia carpophila</i> anamorph, <i>Cladosporium carpophilum</i>)	Use at petal fall. Repeat application at 2 and 5 weeks after petal fall.	
Shot hole (<i>Wilsonomyces carpophilus</i>)	Use at dormant stage, and repeat application at petal fall.	
RESTRICTIONS: • DO NOT apply more than 9 pints of Sequence SC per acre per year (3.9 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence SC per acre per application. • DO NOT apply more than 3 applications per year. • DO NOT apply less than 7 days from the last application. • DO NOT apply within 7 days of harvest.		

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container. **DO NOT** store in areas where temperatures are below 32 degrees F. Store in a cool, dry (preferably locked) area away from other pesticides, fertilizer, food, or feed that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Rigid Plastic Less Than or Equal to 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by state and local authorities.

Rigid Plastic Greater Than 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of empty container in a sanitary landfill or by other procedures allowed by state and local authorities.

Rigid Plastic Bulk and Mini-Bulk Containers: Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

DECLARACIONES PREVENTIVAS

RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS

PELIGRO

Corrosivo. Causa daño ocular irreversible. Irrita la piel. **NO** permita que caiga en los ojos, la piel o la ropa. Dañino si se ingiere. Dañino si se absorbe por la piel. Dañino si se inhala. Evite respirar la niebla o el vapor pulverizado. Lávese con abundante agua y jabón luego de manejar (pesticidas) y antes de comer, beber, masticar chicle, utilizar tabaco, o usar al baño. Quítese y lave la ropa contaminada antes de reutilizarla.

EQUIPO DE PROTECCIÓN PERSONAL (PPE, POR SUS SIGLAS EN INGLÉS)

Algunos materiales que son resistentes químicamente a este producto son laminado impermeable o Vitón.

Los mezcladores, cargadores, aplicadores, y otros manipuladores deben usar:

- Overol (mameluco) sobre camisa de manga corta y pantalón corto
- Calcomines y zapatos resistentes a productos químicos
- Guantes resistentes a productos químicos hechos de material resistente al agua (laminados impermeables; goma de butilo ≥ 14 mils; goma de nitrilo ≥ 14 mils; goma de neopreno ≥ 14 mils; cloruro de polivinilo ≥ 14 mils, o Vitón ≥ 14 mils)
- Gafas de seguridad o máscara protectora
- Equipo de protección para la cabeza resistente a productos químicos, para la exposición por encima de la cabeza
- Delantal resistente a productos químicos para mezcladores y cargadores

Los operarios que mezclan, cargan y aplican productos con pistolas pulverizadoras a presión mecánica en bananos, olivos, frutas de pepita (Grupo de Cultivos 11-10), frutas de hueso (Grupo de Cultivos 12-12) y frutos secos (Grupo de Cultivos 14-12) deben usar, como mínimo, una mascarilla filtrante de partículas aprobada por NIOSH con cualquier filtro N, R o P; O un respirador elastomérico de partículas aprobado por NIOSH con cualquier filtro N, R o P; O un respirador purificador de aire motorizado aprobado por NIOSH con filtros HE.

Los operarios que mezclan y cargan productos para aplicaciones aéreas en bananos, olivos, frutas de pepita (Grupo de Cultivos 11-10), frutas de hueso (Grupo de Cultivos 12-12) y frutos secos (Grupo de Cultivos 14-12) deben usar, como mínimo, una mascarilla filtrante de partículas aprobada por NIOSH con cualquier filtro N, R o P; O un respirador elastomérico de partículas aprobado por NIOSH con cualquier filtro N, R o P; O un respirador purificador de aire motorizado aprobado por NIOSH con filtros HE.

Consulte la declaración de medidas de control de ingeniería para conocer los requisitos adicionales.

REQUISITOS DE SEGURIDAD DEL USUARIO

Deseche la ropa y otros materiales absorbentes que hayan sido empapados o muy contaminados con el concentrado de este producto. **NO** vuelva a reutilizarlos. Siga las instrucciones del fabricante para la limpieza/mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE separadamente de otra ropa para lavar.

DECLARACIÓN DE CONTROL TÉCNICO

Los pilotos deben utilizar una cabina cerrada que satisfaga la definición que figura en la WPS para pesticidas (plaguicidas) agrícolas [40 CFR 170.249(d)(6)].

RIESGOS AMBIENTALES

Este pesticida es tóxico hacia peces e invertebrados acuáticos. **NO** aplique directamente a cuerpos de agua, en áreas donde haya aguas superficiales, o en áreas intermareales que se encuentren bajo el nivel de altura máxima promedio del agua. **NO** contamine cuerpos de agua cuando disponga de las aguas de lavado de equipo o aguas de enjuague. La deriva o el rociado y las escorrentías pueden ser peligrosas para los organismos acuáticos en aguas adyacentes a las áreas tratadas.

Declaración sobre el Agua Superficial y el Control de la Erosión: Este producto puede contaminar el agua a través de la escorrentía y existe la posibilidad de que dicha escorrentía ocurra durante varias semanas posteriores a su aplicación. Los suelos mal drenados y los suelos con capas freáticas poco profundas son más propensos a producir escorrentías que contengan este producto. Una franja de protección o amortiguamiento vegetativo, nivelada y bien mantenida entre las áreas donde este producto es utilizado y cuerpos de agua superficiales tales como lagos (lagunas), ríos y manantiales, reducirá el potencial de contaminación del agua por escorrentía de lluvia. Escorrentías de este producto pueden ser reducidas al evitar aplicaciones cuando estén pronosticados lluvia dentro de 48 horas. Las prácticas adecuadas de control de la erosión reducirán la contribución de este producto a la contaminación de las aguas superficiales.

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal.

ALMACENAMIENTO Y DESECHO

NO contamine agua, comida ni forrajes mediante el almacenamiento y desecho.

ALMACENAMIENTO DE PESTICIDAS: Mantenga este producto en su envase original, bien cerrado. **NO** lo almacene en áreas donde la temperatura sea inferior a 32°F (0°C). Consérvelo en un lugar fresco y seco (preferiblemente bajo llave), alejado de otros pesticidas, fertilizantes, alimentos o forrajes, y fuera del alcance de niños y animales.

DESECHO DE PESTICIDAS: Los residuos de pesticidas son muy peligrosos. Los residuos resultantes del uso de este producto pueden eliminarse en el sitio o en una instalación de eliminación de residuos aprobada. La disposición inadecuada del pesticida sobrante, mezcla de aerosol o aguas de enjuague de contenedores, constituye una violación de la Ley Federal. Si estos residuos no pueden ser eliminados de acuerdo con las instrucciones de la etiqueta, contacte la agencia para el manejo de pesticidas o control ambiental de su estado, o el representante para el Manejo de Residuos Peligrosos de la Oficina Regional de la EPA más cercana para obtener indicaciones del proceso a seguir para su disposición.

MANIPULACIÓN DE ENVASES:

Contenedores de plástico rígido de 5 galones o menos: Recipiente no rellenable. **NO** reutilice ni rellene este recipiente. Ofrezca para el reciclaje si está disponible. Enjuague tres veces (o equivalente) el recipiente inmediatamente después de vaciarlo. Enjuague tres veces siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezclar, y drene por 10 segundos después de que el flujo empiece a gotear. Llene el recipiente 1/4 lleno con agua y tape de nuevo. Agite por 10 segundos. Eche el enjuague en el equipo de aplicación o tanque de mezclar, o almacene el enjuague para uso más tarde o eliminación. Drene por 10 segundos después de que el flujo empiece a gotear. Repita este procedimiento dos veces más. Luego ofrezca para el reciclaje si está disponible o el reacondicionado si está adecuado, o perfore y deseche en un área adecuada para desechos sanitarios o por otros procedimientos aprobados por las autoridades estatales y locales.

Contenedores de plástico rígido de más de 5 galones: Recipiente no rellenable. **NO** reutilice ni rellene este recipiente. Ofrezca para el reciclaje si está disponible. Enjuague tres veces (o equivalente) el recipiente inmediatamente después de vaciarlo. Enjuague tres veces siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezclar. Llene el recipiente 1/4 lleno con agua. Reemplace y asegure los cierres. Vuelque el recipiente al lado y hágalo rodar por 30 segundos, asegurando por lo menos una revolución completa. Ponga el recipiente vertical e inclínelo hacia adelante y hacia atrás varias veces. Dé vuelta al recipiente hacia el lado opuesto e inclínelo hacia adelante y hacia atrás varias veces. Eche el enjuague en el equipo de aplicación o tanque de mezclar, o almacene el enjuague para uso más tarde o eliminación. Repita este procedimiento dos veces más. Luego ofrezca para el reciclaje si está disponible o el reacondicionado si está adecuado, o perfore y deseche en un área adecuada para desechos sanitarios o por otros procedimientos aprobados por las autoridades estatales y locales.

Contenedores de plástico rígido a granel y mini-granel: Recipiente rellenable. Rellene este recipiente únicamente con pesticida. **NO** reutilice este recipiente para ningún otro propósito. La limpieza del recipiente antes de su eliminación final es la responsabilidad de la persona que se deshace del recipiente. La limpieza antes de rellenar es la responsabilidad de la persona que lo va a rellenar. Para limpiar el recipiente antes de su eliminación final, vacíe el contenido restante de este recipiente en el equipo de aplicación o en el tanque de mezcla. Llene el recipiente con un 10% de agua. Agitar vigorosamente o haga recircular el agua con la bomba durante 2 minutos. Vierta o bombee el agua de enjuague en el equipo de aplicación o en el sistema de recolección de agua de enjuague. Repita este procedimiento dos veces más. Luego ofrezca para el reciclaje si está disponible o el reacondicionado si está adecuado, o perfore y deseche en un área adecuada para desechos sanitarios o por otros procedimientos aprobados por las autoridades estatales y locales.

¡EL RECIPIENTE NO ES SEGURO PARA ALIMENTOS, FORRAJES, NI AGUA POTABLE!

NOTES

NOTES

Sequence SC

For control of certain diseases of Almonds, Apples, Apricots*, Bananas*, Beech nuts*, Brazil nuts*, Butternuts*, Cashews*, Cherries, Chestnuts*, Chinquapins*, Filberts*, Hickory nuts*, Macadamia nuts*, Nectarines, Peaches, Peanuts*, Pears, Pecans*, Plums*, Plums-chickasaw*, Plums-damson*, Plums-Japanese*, Plumcots*, Prune plums*, and Walnuts.

*Not registered for use by California.

ACTIVE INGREDIENT:	WT. BY %
Dodine (n-dodecylguanidine acetate)	39.6%**
OTHER INGREDIENTS:	60.4%
TOTAL:	100.0%

**Equivalent to 3.4 pounds Dodine per gallon

KEEP OUT OF REACH OF CHILDREN DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

FIRST AID - IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to by a poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBERS - For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>. Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222.

NOTE TO PHYSICIAN - Probable mucosal damage may contraindicate the usage of gastric lavage.

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - DANGER/PELIGRO - Corrosive. Causes irreversible eye damage. Causes skin irritation. **DO NOT** get in eyes, on skin, or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid breathing spray mist or vapor. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS - This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when

disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **Surface Water and Erosion Control Statement:** This product may contaminate water through runoff and has the potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

DIRECTIONS FOR USE - It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE:** Keep this product in its tightly closed original container. **DO NOT** store in areas where temperatures are below 32 degrees F. Store in a cool, dry (preferably locked) area away from other pesticides, fertilizer, food, or feed that is inaccessible to children and animals. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER HANDLING: Rigid Plastic Less Than or Equal to 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by state and local authorities. **Rigid Plastic Greater Than 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of empty container in a sanitary landfill or by other procedures allowed by state and local authorities. **Rigid Plastic Bulk and Mini-Bulk Containers:** Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities. **CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

Manufactured For: Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 82633-104-83529

EPA Est. No. **(AG)** 72159-GA-001; **(MC)** 89332-GA-001; **(SC)** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: **2.5 Gals.*** ☐ **265 Gals.**

* Unless alternate checked

OPEN HERE