

RESTRICTED USE PESTICIDE / PESTICIDA DE USO RESTRINGIDO

DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS / DEBIDO A SU TOXICIDAD PARA PECES Y ORGANISMOS ACUÁTICOS

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification. / Para la venta al por menor y uso únicamente por aplicadores certificados o personas que estén bajo su directa supervisión y únicamente para los usos detallados en el certificado de la Certificación del Aplicador.

ESFENVALERATE GROUP 3A INSECTICIDE

Trikon

INSECTICIDE

For the control of insect pests on Fields Crops, Vegetable Crops, Fruit Crops, Tree Nut Crops.

ACTIVE INGREDIENT:	By Weight %
Esfenvalerate: (S)-cyano (3-phenoxyphenyl) methyl (S)-4-chloro-alpha-(1-methylethyl) benzeneacetate	8.4%
OTHER INGREDIENTS*:	91.6%
TOTAL:	100.0%
Contains 0.66 lb. active ingredient per gallon. *Contains petroleum distillate	

KEEP OUT OF REACH OF CHILDREN / MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand this label, find someone to explain it to you in detail.)

See label booklet for complete First Aid, Precautionary Statements,
Directions For Use, and Storage and Disposal.

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 82633-111-83529

EPA Est. No. **AG** 72159-GA-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

FIRST AID	
IF SWALLOWED:	• Immediately call a poison control center or doctor for treatment advice. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give any liquid to the person. • DO NOT give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN	
Contains petroleum distillate. Vomiting may cause aspiration pneumonia. If on skin, after drying, apply Vitamin E cream or oil, if available. If not available, apply vegetable oil liberally over painful areas. The oil or cream may be used repeatedly until relief is achieved.	
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378 , Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING/AVISO

May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. **DO NOT** get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber $\geq$ 14 mils, nitrile rubber $\geq$ 14 mils, polyvinyl chloride (PVC) $\geq$ 14 mils, or Viton $\geq$ 14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when cleaning equipment or when disposing of equipment wash-waters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

NON-TARGET ORGANISM ADVISORY STATEMENT

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or to residues in/on blooming crops or weeds. **Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.**

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all **Directions for Use** carefully before applying.

USE RESTRICTIONS

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DO NOT apply this product through any type of irrigation system. **Trikon** may be premixed in a supply tank with water, oil, fertilizer, or other appropriate tank mixed agricultural chemicals.

DO NOT connect an irrigation system (including greenhouse systems) used for **Trikon** application to a public water system.

DO NOT apply a combination of **Trikon** and 2,4-D herbicides in a volume of water less than 2 gallons per acre total spray. Always mix **Trikon** thoroughly in the total volume of spray water first, followed by the addition of the 2,4D herbicide.

DO NOT allow to enter indoor or outdoor drains.

No permita la entrada a desagües internos o externos.



Follow proper disposal procedures on this label.

Siga las indicaciones del etiquetado para el desecho apropiado del producto.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

PRODUCT INFORMATION

Trikon emulsifiable concentrate contains 0.66 pounds of active ingredient per gallon. For the applications given below, mix the required amount of **Trikon** in sufficient diluent to provide uniform coverage (refer to **Use Tables**). **Trikon** may be applied by ground or aerial application equipment. For aerial application use the following directions unless otherwise specified in this label: use a minimum of 2 gallons per acre (gpa) of water, except in tree and orchard crops use a minimum of 10 gpa.

Apply at the recommended rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

INSECT RESISTANCE MANAGEMENT

For resistance management, **Trikon** contains esfenvalerate and is classified in the pyrethroids, pyrethrins chemical class as a Group 3A insecticide, sodium channel modulators.

Any insect population may contain individuals naturally resistant to **Trikon** and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Trikon** or other Group 3A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture.
- In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.

- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact your local Sharda USA LLC representative at 1-(910) 859-3090 Monday through Friday, 9 AM to 4 PM EST, or <https://shardausa.com>.

INTEGRATED PEST MANAGEMENT

It is recommended to use Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

SPRAY PREPARATION

Spray equipment must be clean and free of previous pesticide deposits before applying **Trikon**.

TANK MIXING AND COMPATIBILITY

Unless directed otherwise in a specific crop section of this label, **DO NOT** tank mix **Trikon** with fungicides containing fentin hydroxide (triphenyltin hydroxide) such as "Super Tin" as crop injury may result.

This product can be mixed with pesticide products labeled for use on crops on this label in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Since formulations may be changed and new ones introduced, it is recommended that users premix a small quantity of a desired tank mix and observe for possible adverse changes (settling out, flocculation, etc.). Avoid mixtures of several materials and very concentrated spray mixtures. For best results, use of spray equipment having continuous agitation is recommended.

Trikon may be tank mixed with herbicide products when insect populations require control concurrent with the need for weed control. Follow all herbicide and **Trikon** label directions regarding proper usage.

Trikon may be used in combination with 2,4-D herbicides providing that the following mixing directions are followed: 1) **DO NOT** apply the combination in a volume of water less than 2 gallons per acre total spray. 2) Always mix **Trikon** thoroughly in the total volume of spray water first, followed by the addition of the 2,4D herbicide. Because of the availability of a great variety of 2,4-D herbicide products, a test for physical compatibility should be conducted before field mixtures of a particular combination are made.

CHEMIGATION

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) row, traveler, big gun, solid set, or hand move irrigation systems. **DO NOT** apply this product through any other type of irrigation system. **Trikon** may be premixed in a supply tank with water, oil, fertilizer, or other appropriate tank mixed agricultural chemicals. A pretest of physical compatibility for untried tank mixes is advised. Agitation may be necessary. Application should be in sufficient water and of sufficient duration to apply the recommended rate evenly to the entire treated area. No run-off can be permitted during chemigation. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

DO NOT connect an irrigation system (including greenhouse systems) used for **Trikon** application to a public water system.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut-off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

CROP ROTATION

ALL ROTATION CROPS MAY BE PLANTED IMMEDIATELY FOLLOWING LAST APPLICATION.

VEGETATIVE FILTER STRIPS

Construct and maintain a vegetative filter strip, according to the width specified below, of grass or other permanent vegetation between the field edge and nearby down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing esfenvalerate onto fields where a maintained vegetative filter strip of at least 25 feet exists between the field edge and where a down gradient aquatic habitat exists. This minimum required width of 25 feet may be reduced or removed under the following conditions:

- For Western irrigated agriculture, a maintained vegetative filter strip of at least 10 feet wide is required. Western irrigated agriculture is defined as irrigated farmland in the following states: WA, OR, CA, ID, NV, UT, AZ, MT, WY, CO, NM, and TX (west of I-35).
- For Western irrigated agriculture, if a sediment control basin is already present, a vegetative filter strip is not required.
- In all other areas, a vegetative filter strip with a minimum width of 25 feet is required, unless the following conditions are met. The vegetative filter strip requirement may be reduced from 25 feet to 15 feet if at least one of the following applies:
 - o The area of application is considered prime farmland (as defined in 7 CFR § 657.5).
 - o Conservation tillage is being implemented on the area of application. Conservation tillage is defined as any system that leaves at least 30% of the soil surface covered by residue after planting. Conservation tillage practices can include mulch-till, no-till, or strip-till.
 - o A functional terrace system is maintained on the area of application.
 - o Water and sediment control basins for the area of application are functional and maintained.
 - o The area of application is less than or equal to 10 acres.

For further guidance on vegetated filter strips, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. <https://www.regulations.gov/document/EPA-HQ-OPP-2008-0331-0175>.

BUFFER ZONES TO WATER BODIES

Ground Application

DO NOT apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Ultra Low Volume (ULV) Aerial Application

DO NOT apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Non-ULV Aerial Application

DO NOT apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzle and pressure that deliver medium or coarser droplets (ASABE S641).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wing-span for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% of the wingspan or less for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- If the windspeed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 - 15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Airblast Applications:

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to select nozzle and pressure that deliver medium or coarser droplets (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

- For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

- Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

- Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

- Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

SPRAY TANK CLEANOUT

Immediately following application of **Trikon**, thoroughly clean all mixing and spray equipment. Flush the tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Take all necessary precautions when cleaning equipment. **DO NOT** clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

POLLINATOR PROTECTION

Following best management practices can help reduce risk to terrestrial pollinators. Examples of best management practices include applying pesticides in the evening and at night when pollinators are not foraging and checking to confirm hive locations before spraying. For additional resources on pollinator best management practices, visit <https://www.epa.gov/pollinator-protection/find-best-management-practices-protect-pollinators>.

Managed pollinator protection plans are developed by states/tribes to promote communication between growers, landowners, farmers, beekeepers, pesticide users, and other pest management professionals to reduce exposure of bees to pesticides. If available, visit state plans for additional information on how to protect pollinators.

How to Report Bee Kills

It is recommended that users contact both the state lead agency and the U.S. Environmental Protection Agency to report bee kills due to pesticide application. Bee kills can be reported to EPA at beekill@epa.gov. To contact your state lead agency, see the current listing of state pesticide regulatory agencies at the National Pesticide Information Center's website: http://npic.orst.edu/reg/state_agencies.html.

CROP-SPECIFIC USES

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)	
		LB. A.I./A	FL. OZ./A			
Corn (field)*	Western Bean Cutworm	0.015 - 0.03	2.9 - 5.8	44 - 22	21	
	Armyworm (True Armyworm)	0.03 - 0.05	5.8 - 9.6	22 - 13		
	Black Cutworm (except CA)					
	Chinch Bug					
	Corn Earworm					
	Corn Leaf Aphid					
	Corn Rootworm (adult control)					
	Cutworm Flea Beetle					
	Grasshopper					
	Japanese Beetle (adult) (except CA)					
	Oat Bird-Cherry Aphid					
	Southwestern Corn Borer					
	Stalk Borer					
	European Corn Borer	0.04 - 0.05	7.8 - 9.6	16 - 13		
	Black Cutworm - Trikon may be applied at 3.2 - 9.6 fl. oz./acre (0.0165 - 0.05 lb. a.i./acre) for the control of black cutworm when applied at planting of corn (except CA).					
	Chinch Bug - Over the top sprays, as used for control of ear and stalk infesting corn pests, are not adequate for chinch bug control. It is very important that the spray be directed at the base of the plant through the use of drop nozzles or some other mechanism.					
Corn Earworm - First application should be at or before silking. Repeat applications may be applied if economically damaging populations exist. Subsequent applications should be made at 3 - 5 day intervals until silking is completed.						
Corn Leaf Aphid, Oat Bird-Cherry Aphid - For optimum results, direct the spray at the aphid population so as to achieve maximum coverage of the exposed insects. Aphids not contacted by the spray, such as in whorls and leaf axils, may not be adequately controlled.						
Corn Rootworm (Adult) - Apply at the first sign of silk feeding.						
Trikon may be tank-mixed with methyl parathion and applied on field corn where supplemental control of Adult Corn Rootworm is desired in conjunction with insects controlled by Trikon when used alone. Refer to the Trikon and methyl parathion labels for appropriate rates of the individual products for controlling the respective insects.						
Cutworm - Applications for cutworm control may be applied before, during, or after planting as required to protect emerging or emerged corn seedlings.						
European Corn Borer - First brood: Spray while eggs are in the blackhead stage or before the larvae enter the whorl. Application by ground equipment is suggested. Good coverage of both upper and lower leaf surfaces is essential. This can be accomplished with drop nozzles over the row and on each side of the corn plant. Multiple applications may be required when egg laying is prolonged or where moderate to heavy populations are present. A higher rate is recommended for moderate to heavy populations. Proper coverage by ground equipment usually requires 20 - 30 gallons of carrier. Once larvae enter the whorl, foliar sprays will not provide adequate control. Second brood: Make applications when sufficient egg masses are found. Spray when eggs are in the blackhead stage or starting to hatch. When egg laying is prolonged or a third generation is present, additional sprays may be required. A higher rate is recommended for moderate to heavy populations. Good coverage above, below, and in the ear zone is essential. This usually requires 2 - 3 gallons of carrier by air. If ground equipment is used, drop nozzles on each side of the plant will provide best coverage.						
Trikon may be tank mixed with methyl parathion and applied on field corn where supplemental control of European corn borer is desired in conjunction with insects controlled by Trikon when used alone. Refer to the Trikon and methyl parathion labels for appropriate rates of the individual products for controlling the respective insects.						
Grasshopper - For control of first and second instar grasshopper nymphal stages a rate range of 3.9 to 5.8 fluid ounces of product per acre (0.02 - 0.03 lb. a.i./A) can be used. Correct timing of spray applications to the first and second instar nymphal stages and thorough coverage is critical to achieve optimum control. For grasshopper nymph stages larger than second instar, use Trikon at use rates of 5.8 to 9.6 fluid ounces of product per acre (0.03 - 0.05 lb. a.i./A).						
Southwestern Corn Borer - For moderate to heavy infestations, higher rates (0.036-0.05 lb. a.i. per acre) are recommended.						
Stalk Borer, Flea Beetle - Application must be made early in migration from grassy areas to corn, before borers enter the plant.						
Western Bean Cutworm - Apply before larvae enter the ear.						
* DO NOT apply more than 0.25 lb. a.i. per acre per season.						

(continued)

CROP	INSECT	APPLICATION RATE				ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A				
Corn (field) At Plant	Cutworm	0.0023 lb. a.i. per 1,000 ft. of row		0.45 fl. oz. per 1,000 ft. of row			21
	Apply as an in-furrow, T-band, or band treatment using a minimum 4" band. Use the table below to determine the pounds active ingredient and fluid ounces of Trikon applied at 0.0023 lb. a.i. per 1000 ft. or row for various row spacings.						
	In-furrow Applications: Apply into the seed furrow through spray nozzles behind the planter furrow openers and in front of the press wheel.						
	Banded Applications: Apply at planting as a 4 - 7 inch T-band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel.						
	Apply a minimum spray volume of 3 gallons per acre.						
	DO NOT exceed 0.05 lb. a.i. per acre per season as an at-plant application.						
	DO NOT apply more than 0.25 lb. a.i. per acre per season including at-plant plus foliar applications of Trikon .						
	Row Spacings (inches)	40"	38"	36"	30"		
	Linear ft./A	13.068	13.756	14.520	17.424		
	Trikon lb. a.i./A	0.03	0.032	0.033	0.04		
	Trikon fl. oz./A	5.8	6.2	6.4	7.8		
Corn (Pop)	For specific insect control recommendations refer to Field Corn (above). Follow directions carefully.						1
	Multiple applications and/or shortened intervals between sprays must be used to insure proper insect control. DO NOT apply more than 0.5 lb. a.i. per acre per season.						
Corn (Seed)	For specific insect control recommendations refer to Field Corn (above). Follow directions carefully.						
	Multiple applications and/or shortened intervals between sprays must be used to insure proper insect control. DO NOT apply more than 0.25 lb. a.i. per acre per season.						

(continued)

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Cotton	Cotton Leaf Perforator	0.03	5.8	22	21
	Beet Armyworm* Black Cutworm (except CA) Boll Weevil Cabbage Looper Cotton Aphid* Cotton Bollworm Cotton Leafworm Cutworms Fleahoppers Grasshoppers Green Stink Bug (except CA) Leafhoppers Lygus Bugs Pink Bollworm Plant Bugs Saltmarsh Caterpillar Southern Green Stink Bug (except CA) Thrips (on seedling cotton) Tobacco Budworm Whitefly*	0.03 - 0.05	5.8 - 9.6	22 - 13	
	NOTE: For light infestation of the above insects	0.02	3.9	33	
*Aids in control. May be applied in water or nonvolatile vegetable oils. When applying Trikon in an oil carrier, apply a total spray volume of at least 1 qt. per acre. When applying Trikon in a water carrier, apply at least 1 gal. per acre by air (at least 3 gals. per acre in Arizona and 5 gals. per acre in California) or 4 gal. per acre by ground. DO NOT apply more than 0.5 lb. a.i. per acre per season. DO NOT graze livestock on treated fields or feed treated trash. Black Cutworm - Trikon may be applied at 3.2 - 9.6 fl. oz./acre (0.0165 - 0.05 lb. a.i./acre) for the control of black cutworm when applied at planting of cotton (except CA). Boll Weevil - To control Boll Weevil infestations, a 3 to 5 day interval between applications may be necessary. Heliothis spp. - Trikon can provide contact ovicidal effect on <i>Heliothis</i> spp. eggs when applied according to label directions for control of tobacco budworm; application should be timed to correspond with peak egg deposition to achieve maximum ovicidal effect. Use on this pest stage (egg) is not registered in California. DO NOT make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season.					

(continued)

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Peanuts	Corn Earworm Potato Leafhopper Red-necked Peanut Worm Velvetbean Caterpillar	0.015 - 0.03	2.9 - 5.8	44 - 22	21
	Beet Armyworm* Cutworms Granulate Cutworm Grasshoppers	0.033 - 0.05	5.8 - 9.6	22 - 13	
	Fall Armyworm* Lesser Cornstalk Borer*	0.05	9.6	13	
	*Aids in control. DO NOT feed or graze livestock on treated vines. DO NOT apply more than 0.15 lb. a.i. per acre per season.				
Sorghum (Grain) Except CA	Sorghum Midge	0.015 - 0.03	2.9 - 5.8	44 - 22	21
	Black Cutworm Chinch Bugs Corn Earworm (headworm) Cutworms	0.03 - 0.05	5.8 - 9.6	22 - 13	
	DO NOT apply more than 0.15 lb. a.i. per acre per season. When applying in nonvolatile vegetable oils use a total spray volume of 1 or more qts. per acre. Black Cutworm - Trikon may be applied at 3.2 - 9.6 fl. oz./acre (0.0165 - 0.05 lb. a.i./acre) for the control of black cutworm when applied at planting of sorghum. Chinch Bug Control - For optimum results, spray should be directed at base of plants.				

(continued)

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Soybean	Green Cloverworm Mexican Bean Beetle Potato Leafhopper Saltmarsh Caterpillar Velvetbean Caterpillar Woollybear Caterpillar	0.015 - 0.03	2.9 - 5.8	44 - 22	21
	Bean Leaf Beetle Beet Armyworm* Cabbage Looper Corn Earworm Cutworms Grasshoppers Green Stink Bug (except CA) Japanese Beetle (adult) Southern Green Stink Bug Soybean Aphid (except CA) Three-cornered Alfalfa Hopper	0.03 - 0.05	5.8 - 9.6	22 - 13	
Grasshopper - For control of first and second instar grasshopper nymphal stages a rate range of 3.9 to 5.8 fluid ounces of product per acre (0.02 - 0.03 lb. a.i./A) can be used. Correct timing of spray applications to the first and second instar nymphal stages and thorough coverage is critical to achieve optimum control. For grasshopper nymph stages larger than second instar, use Trikon at use rates of 5.8 to 9.6 fluid ounces of product per acre (0.03 - 0.05 lb. a.i./A). Soybean Aphid - Trikon provides control of soybean aphid, however under certain conditions such as rapid aphid population growth, or extremely high populations, a tank mixture may be considered. Trikon can be tank mixed with other insecticides such as chlorpyrifos (e.g. "Lorsban") or methomyl (e.g. DuPont™ Lannate®) to achieve rapid knockdown of soybean aphid. Because Lannate® is a fast-acting contact insecticide, for best results follow direct spraying of the target insect. When preparing a tank mixture, read and follow the label instructions for all products in the mixture regarding restrictions, requirements and proper usage. Use sufficient water to obtain thorough, uniform coverage. For aerial application use a minimum of 2 gallons per acre, and for ground application use a minimum of 10 gallons per acre. *Aids in control. When applying in nonvolatile vegetable oils, use a total spray volume of at least 1 qt. DO NOT feed or graze livestock on treated fields. DO NOT apply more than 0.2 lb. a.i. per acre per season.					
Sugar Beets	Beet Armyworm* Beet Webworm Cabbage Looper Cutworms Flea Beetle (except CA) Grasshoppers Leafhoppers Saltmarsh Caterpillar Sugar Beet Root Maggot (adult) (except CA)	0.03 - 0.05	5.8 - 9.6	22 - 13	21
	Grasshopper - For control of first and second instar grasshopper nymphal stages a rate of 3.9 to 5.8 fluid ounces of product per acre (0.02 - 0.03 lb. a.i./A) can be used. Correct timing of spray applications to the first and second instar nymphal stages and thorough coverage is critical to achieve optimum control. For grasshopper nymph stages larger than second instar, use Trikon at use rates of 5.8 to 9.6 fluid ounces of product per acre (0.03 - 0.05 lb. a.i./A). *Aids in control. DO NOT apply more than 0.15 lb. a.i. per acre per season. Apply with ground or air equipment using sufficient water to provide uniform coverage (minimum of 2 gals. of water per acre).				

(continued)

CROP	INSECT	APPLICATION RATE				ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)	
		LB. A.I./A	FL. OZ./A					
Sugar Beets At Plant	Cutworm	0.0023 lb. a.i. per 1,000 ft. of row	0.45 fl. oz. per 1,000 ft. of row				21	
	Apply as an in-furrow, T-band, or band treatment using a minimum 4" band. Use the table below to determine the pounds active ingredient and fluid ounces of Trikon applied at 0.0023 lb. a.i. per 1000 ft. of row for various row spacings.							
	In-Furrow Applications: Apply into the seed furrow through spray nozzles, behind the planter furrow openers and in front of the press wheel.							
	Banded Applications: Apply at planting as a 4 - 7 inch T-band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel.							
	Apply a minimum spray volume of 3 gallons per acre.							
	DO NOT exceed 0.05 lb. a.i. per acre per season as an at-plant application.							
		DO NOT apply more than 0.25 lb. a.i. per acre per season including at-plant plus foliar applications of Trikon .						
	Row Spacing (inches)	40"	38"	36"	30"	22"		
	Linear ft./A	13,068	13,756	14,520	17,424	23,760		
	Trikon lb. a.i./A	0.03	0.032	0.033	0.04	0.05		
	Trikon fl. oz./A	5.8	6.2	6.4	7.8	9.6		
Sugarcane	Sugarcane Borer	0.03 - 0.05	5.8 - 9.6			22 - 13	21	
	DO NOT apply more than 0.2 lb. a.i. per acre per season.							
Sunflower	Sunflower Beetle (except CA)	0.0075 - 0.03	1.45 - 5.8			88 - 22	28	
	Banded Sunflower Moth	0.03 - 0.05	5.8 - 9.6			22 - 13		
	Beet Armyworm*							
	Cutworms							
	Grasshoppers							
	Heliothis (complex)							
	Leafhoppers							
	Sunflower Maggot							
	Sunflower Moth							
	Sunflower Seed Weevil							
Sunflower Stem Weevil								
Grasshopper - For control of first and second instar grasshopper nymphal stages a rate range of 3.9 to 5.8 fluid ounces of product per acre (0.02 - 0.03 lb. a.i./A) can be used. Correct timing of spray applications to the first and second instar nymphal stages and thorough coverage is critical to achieve optimum control.								
For grasshopper nymph stages larger than second instar, use Trikon at use rates of 5.8 to 9.6 fluid ounces of product per acre (0.03 - 0.05 lb. a.i./A).								
*Aids in control.								
DO NOT apply more than 0.2 lb. a.i. per acre per season.								

FRUITS

DILUTE SPRAY: Apply specified dosage per 100 gallons of water in a uniform spray applied to the point of drip with conventional ground equipment. **DO NOT** exceed maximum number of gallons per acre indicated.

NOTE: In order to apply the correct amount of **Trikon** insecticide to your orchard you must know the number of gallons of water needed to spray one acre of your trees to the point of drip. If you **DO NOT** already know this gallonage, you should conduct a test to determine it. If you **DO NOT** know how to conduct such a test with your equipment, you should ask for assistance from your equipment dealer or State Extension specialist.

CONCENTRATE SPRAY: Apply specified dosage per acre in no less than 30 gals. of water per acre by ground equipment.

FOR AERIAL APPLICATION IN TREE AND ORCHARD CROPS: Use a minimum of 10 gallons of water per acre. When applying **Trikon** by air, consult your Cooperative Extension Service for further application guidelines.

CROP	INSECT	APPLICATION RATE			ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A	FL. OZ./100 GAL.		
Apples	Apple Aphid Apple Maggot Codling Moth Green Fruitworm Lesser Appleworm Mullein Plant Bug (except CA) Oblique Banded Leafroller Oriental Fruit Moth Periodical Cicada Plant Bugs (Tarnished Plant Bug, Stink Bugs) Plum Curculio Red-Banded Leafroller Rosy Apple Aphid San Jose Scale (fruit infestations only) Tentiform Leaf Miner Tufted Apple Bud Moth Variegated Leafroller White Apple Leafhopper	0.025 - 0.075	4.8 - 14.5	2.0 - 5.8	26 - 9	21
	Apple Ermine Moth (ID, OR, & WA only)			3.0		
	Tufted Apple Bud Moth (overwintering) (MD, NC, NJ, PA, VA, WV only)	0.04 - 0.075	8.0 - 14.5		16 - 9	
DO NOT feed or graze livestock on treated orchard floors. DO NOT apply more than 0.525 lb. a.i. per acre per season. For dilute spray apply 200 - 600 gals. per acre, but DO NOT apply more than 14.5 fl. oz. of Trikon per acre per treatment. Apple Ermine Moth - Apply with 2 to 4 gallons of superior spray oil in 100 gallons of water in a spray-to-wet application to insure thorough coverage of all stems and branches where Apple Ermine Moth hibernacula are found. When using on apple nursery stock, DO NOT treat bundled plants since it is difficult to achieve a full coverage application which could result in less than complete control. Make first application in the fall after 90% of leaf fall has occurred – usually after October 15. Make a second application 7 to 14 days later. NOTE: Overwintering larvae DO NOT die until approximately 30 days after application. Plant Bug, Rosy Apple Aphid Control - Time of application is critical in achieving control. Use prebloom and post bloom spray timings recommended by State Extension Services. Tufted Apple Bud Moth (overwintering) - For use on apple for the control of overwintering larvae of the tufted apple bud moth with directed ground application to the apple orchard floor. Make one application of Trikon at either pink stage of apple or at petal fall stage of apple. Use the lower rate on small larvae (pink stage of apple) and/or on lower populations. Use the higher rate on larger larvae (petal fall stage of apple) and/or on moderate to high populations. Apply specified dosage per acre to the orchard floor in no less than 30 gals. of water per acre by ground to obtain uniform coverage. Apply treatment in a band from trunk to drip line to allow coverage of areas where overwintering tufted apple bud moth are found. Beneficial Insects: Application of Trikon to the groundcover at the pink stage of apple development may be toxic to overwintering <i>Stethorus punctum</i>. <i>S. punctum</i> is a coccinellid insect and the major predator of spider mites in the MD, NC, NJ, PA, VA and WV fruit growing areas. This predator overwinters in the same areas of the orchard groundcover as the tufted apple bud moth and moves into apple trees from mid-April through mid-May when maximum daily temperatures exceed 68° F. Emergence from the groundcover is 20 - 70% complete by the pink stage and 90 - 100% complete by petal fall on the apple cultivar Yorking.						

(continued)

CROP	INSECT	APPLICATION RATE			ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A	FL. OZ./100 GAL.		
Blueberry (except CA)	Aphids (NJ only) Blueberry Spanworm Cherry Fruitworm* Cranberry Fruitworm Cranberry Weevil* Grasshoppers Japanese Beetle Leafhoppers Red Striped Fireworm*	0.025 - 0.05	4.8 - 9.6		26 - 13	14
	Blueberry Maggot Black Vine Weevil (adult control) Strawberry Root Weevil (adult control) (OR, WA only)	0.05	9.6		13	
	*Aids in control. DO NOT apply more than 0.2 lb. a.i. per acre per season (38.4 fluid ounces of product per acre per season). Use of ground application is recommended; for ground application use a minimum of 50 gals. water per acre. DO NOT apply this product through any type of irrigation system. NOTE: Trikon can act as a bee repellent, DO NOT apply within 7 days of pollination. Apply as a pre-bloom or post-bloom spray only. Black Vine Weevil & Strawberry Root Weevil (adult control) (OR, WA only) - Look for leaf notching beginning in late May to early June as the first sign of weevil feeding. Also check for adults on or just below the soil surface around the base of plants. Apply Trikon within two to three weeks of first sign of infestation. DO NOT apply by air; apply by ground using a minimum of 50 gallons of water per acre. Direct spray to provide full coverage of foliage and soil area around base of plants. Best results are from applications made after dark when temperatures are warm and weevils are actively feeding. Root weevils emerge over a several week period, make additional applications when signs of new feeding appear.					
Caneberries (blackberries, boysen-berries, dewberries, loganberries, raspberries, youngberries, and varieties of these) (except CA)	Aphids Oblique Banded Leafroller Orange Tortrix	0.025 - 0.05	4.8 - 9.6		26 - 13	7
	Adult Root Weevils* (OR & WA only)	0.05	9.6		13	
	*Aids in control. DO NOT apply by air. DO NOT apply more than 0.15 lb. a.i. per acre per season. DO NOT apply this product through any type of irrigation system. NOTE: Trikon can act as a bee repellent, DO NOT apply within 7 days of pollination. Apply as a pre-bloom or post-bloom spray only. Remove bees prior to application. For maximum safety to bees, apply Trikon in the evening after sunset. Adult Root Weevils (OR, WA only) - Look for leaf notching beginning in late May to early June as the first sign of weevil feeding. Also check for adults on or just below the soil surface around the base of plants. Apply Trikon within two to three weeks of infestation. Apply by ground using a minimum of 50 gallons of water per acre. Direct spray to provide full coverage of foliage and soil area around base of plants. Best results are from applications made after dark when temperatures are warm and weevils are actively feeding. Root weevils emerge over a several week period, make additional applications when signs of new feeding appear. Oblique Banded Leafroller, Orange Tortrix and Aphids - Apply as a full coverage spray in a minimum of 50 gallons of water with ground equipment only. Apply no earlier than 12 days before harvest and no later than 7 days before harvest.					

(continued)

CROP	INSECT	APPLICATION RATE			ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A	FL. OZ./100 GAL.		
Kiwifruit	Boxelder Bug (suppression only)	0.05	9.6		13	14
	Spray in sufficient water for thorough coverage. A maximum of 7 applications is allowed per season (total of 0.35 lb. a.i. per acre/season) with a minimum of 7 days between treatments.					
Pear	Codling Moth Green Fruitworm Leafrollers Pear Psylla Pear Slug Periodical Cicada Plum Curculio	0.025 - 0.075	4.8 - 14.5	2.0 - 5.8	26 - 9	28
	DO NOT apply more than 0.375 lb. a.i. per acre per season. DO NOT apply more than 0.225 lb. a.i. per acre between bloom and harvest. DO NOT feed or graze livestock on treated orchard floors. For dilute spray apply 200 - 600 gals. per acre, but DO NOT apply more than 14.5 fl. oz. of Trikon per acre per treatment.					
Pear (Dormant)	Pear Psylla	0.05 - 0.1	9.6 - 19.2	7.3 - 12.8	13.2 - 6.6	28
	Apply during dormant to prebloom (white bud) stage only. DO NOT apply more than 0.2 lb. a.i. per acre per season. DO NOT graze orchard floor. For dilute spray apply 150 - 250 gals. per acre but DO NOT apply more than 19.2 fl. oz. of Trikon per acre per treatment.					

(continued)

CROP	INSECT	APPLICATION RATE			ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A	FL. OZ./100 GAL.		
Stone Fruits (including apricots, cherries, nectarines, peaches, plums, prune plums)	American Plum Borer Black Cherry Aphid Cherry Fruit Fly Green Fruitworm Leafhoppers Leafrollers Lesser Peach Tree Borer Oriental Fruit Moth Peach Tree Borer Peach Twig Borer Periodical Cicada Plant Bugs (Tarnished Plant Bug, Stink Bugs) Plum Curculio Western Cherry Fruit Fly	0.025 - 0.075	4.8 - 14.5	2.0 - 5.8	26 - 9	14
	Peach Twig Borer (Dormant) (CA only)	0.04 - 0.075	8.0 - 14.5	3.1 - 5.8	16 - 9	
	Tufted Apple Bud Moth (overwintering) (Peach only) (MD, NC, NJ, PA, VA and WV only)	0.04 - 0.075	8.0 - 14.5		16 - 9	
	DO NOT apply more than 0.375 lb. a.i. per acre per season with no more than 0.3 lb. a.i. per acre per season between bloom and harvest. For dilute spray apply 200 - 400 gals. per acre, but DO NOT apply more than 14.5 fl. oz. of Trikon per acre per treatment. DO NOT graze livestock on treated orchard floors. Peach Tree Borer, American Plum Borer, Lesser Peach Tree Borer Control - Apply as directed trunk and scaffold limb spray. Thorough coverage of trunk and scaffold limbs is required. Peach Twig Borer (Dormant) - Make application with an EPA registered dormant oil; for specific recommendations on use of oil consult manufacturer's label. For best performance, ground application equipment is recommended. Peach Twig Borer, Plum Curculio, Oriental Fruit Moth, Cherry Fruit Fly, Western Cherry Fruit Fly, Leafrollers, Black Cherry Aphid, Periodical Cicada Control - Apply by ground sprayer to achieve thorough coverage of all aerial portions of the tree. Plant Bug Control - Time of application is critical in achieving control. Use prebloom and postbloom spray timings recommended by State Extension Services. Tufted Apple Bud Moth (overwintering) - For use on peach for the control of overwintering larvae of the tufted apple bud moth with directed ground application to the peach orchard floor. Make one application of Trikon at either popcorn stage of peach or at shuck split stage of peach. Use the lower rate on small larvae (popcorn stage of peach) and/or on lower populations. Use the higher rate on larger larvae (shucksplit stage of peach) and/or on moderate to high populations. Apply specified dosage per acre to the orchard floor in no less than 30 gals. of water per acre by ground to obtain uniform coverage. Apply treatment in a band from trunk to drip line to allow coverage of areas where overwintering tufted apple bud moth are found. Beneficial Insects: Application of Trikon to the groundcover at the popcorn stage of peach development may be toxic to overwintering <i>Stethorus punctum</i>. <i>S. punctum</i> is a coccinellid insect and the major predator of spider mites in the MD, NC, NJ, PA, VA and WV fruit growing areas. This predator overwinters in the same areas of the orchard groundcover as the tufted apple bud moth and moves into peach trees from mid-April through mid-May when maximum daily temperatures exceed 68° F. Emergence from the groundcover is 90 - 100% complete by shucksplit. Shucksplit is the preferred timing to minimize predator toxicity.					

TREE NUT CROPS

DILUTE SPRAY: Apply specified dosage per 100 gallons of water in a uniform spray applied to the point of drip with conventional ground equipment. **DO NOT** exceed maximum number of gallons per acre indicated.

NOTE: In order to apply the correct amount of **Trikon** insecticide to your orchard you must know the number of gallons of water needed to spray one acre of your trees to the point of drip. If you **DO NOT** already know this gallonage, you should conduct a test to determine it. If you **DO NOT** know how to conduct such a test with your equipment, you should ask for assistance from your equipment dealer or State Extension specialist.

CONCENTRATE SPRAY: Apply specified dosage per acre in no less than 30 gals. of water per acre, by ground equipment.

FOR AERIAL APPLICATION IN TREE AND ORCHARD CROPS: Use a minimum of 10 gallons of water per acre. When applying **Trikon** by air, consult your Cooperative Extension Service for further application guidelines.

CROP	INSECT	APPLICATION RATE			ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A	FL. OZ./100 GALS.		
Almonds	Navel Orangeworm Peach Twig Borer Peach Twig Borer (Dormant) (CA Only)	0.05 - 0.1	9.6 - 19.2	7.3 - 12.8	13.2 - 6.6	21
	DO NOT apply more than 0.2 lb. a.i. per acre per season. DO NOT graze livestock on treated orchard floors. Peach Twig Borer (Dormant) - Make application with an EPA registered dormant oil; for specific recommendations on use of oil consult manufacturer's label. For best performance, ground application equipment is recommended.					
Filberts	Filbertworm Oblique Banded Leafroller	0.05 - 0.1	9.6 - 19.2	7.3 - 12.8	13.2 - 6.6	21
	Make first application after emergence of filbert worm moths in early summer. DO NOT apply a second treatment earlier than three weeks after the first. DO NOT apply more than 0.2 lb. a.i. per acre per season. For dilute spray, apply 200 - 400 gals. per acre, but DO NOT apply more than 19.2 fl. oz. of Trikon per acre per treatment. DO NOT graze livestock on treated orchard floors.					
Pecans	Hickory Shuckworm Pecan Aphids Pecan Nut Casebearer Pecan Leaf Phylloxera Pecan Spittlebug Pecan Stem Phylloxera Pecan Weevil	0.025 - 0.075	4.8 - 14.5	2.0 - 5.8	26 - 9	21
	DO NOT feed or graze livestock on treated orchard floors. DO NOT apply more than 0.3 lb. a.i. per acre per season. For dilute spray apply 200 - 600 gals. per acre, but DO NOT apply more than 14.5 fl. oz. of Trikon per acre per treatment. Trikon may be tank-mixed with fungicides containing fentin hydroxide (triphenyltin hydroxide) such as "Super Tin". Refer to the Trikon and fentin hydroxide (triphenyltin hydroxide) labels for appropriate rates of the individual products for controlling the respective pests. Phylloxera - Correct timing of spray applications is critical in achieving optimum control of leaf and stem phylloxera. Consult local spray recommendations for correct times of application.					
Walnuts	Codling Moth Navel Orangeworm Walnut Aphid Walnut Husk Fly	0.05 - 0.1	9.6 - 19.2	4.0	13 - 6	21
	DO NOT apply more than 0.2 lb. a.i. per acre per season. For dilute sprays, apply 200 - 400 gals. per acre. DO NOT feed or graze livestock on treated crop floors. NOTE: Use of baits in Walnut Husk fly sprays is recommended where endorsed by local Agricultural Extension Service.					

VEGETABLE CROPS

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Artichoke	Artichoke Plume Moth	0.03 - 0.05	5.8 - 9.6	22 - 13	1
	DO NOT apply more often than each 14 days. Apply no more than 0.15 lb. a.i. per acre between bud formation and harvest of an individual fruit. Apply in a minimum of 10 gallons per acre by air and 50 - 200 gallons per acre by ground (use sufficient water to obtain coverage without excessive runoff).				
Beans, Dry (Including adzuki bean, blackeyed pea, broad bean (dry), chickpea, cow pea, crowder pea, field bean, kidney bean, lima bean (dry), mung bean, navy bean, pinto bean, southern pea, teparty bean) Peas, Dry Lentils	Leafhoppers (except CA) Mexican Bean Beetle Saltmarsh Caterpillar	0.015 - 0.03	2.9 - 5.8	44 - 22	21
	Beet Armyworm* Cabbage Looper Corn Earworm Corn Rootworm (Adults) Cowpea Curculio Cucumber Beetle Cutworms Grasshoppers Green Cloverworm Leafhoppers Painted Lady Butterfly (larvae) Pea Aphid Potato Leafhopper Soybean Aphid (except CA) Velvetbean Caterpillar Western Bean Cutworm	0.03 - 0.05	5.8 - 9.6	22 - 13	
	Pea Leaf Weevil (ID, OR & WA only) Pea Weevil (ID, OR & WA only)	0.025 - 0.05	4.8 - 9.6	26 - 13	
	*Aids in control. DO NOT apply more than 0.2 lb. a.i. per acre per season. DO NOT feed or graze livestock on treated vines. Pea Weevil & Pea Leaf Weevil (ID, OR & WA) - Time of application is critical in achieving control of pea weevil. For optimum results, apply at bloom prior to detecting adult pea weevils. Once adult pea weevil populations reach a level of 2 or more adults per 25 sweeps, control may be reduced. Grasshopper - For control of first and second instar grasshopper nymphal stages a rate of 3.9 to 5.8 fluid ounces of product per acre (0.02 - 0.03 lb. a.i./A) can be used. Correct timing of spray applications to the first and second instar nymphal stages and thorough coverage is critical to achieve optimum control. For grasshopper nymph stages larger than second instar, use Trikon at use rates of 5.8 to 9.6 fluid ounces of product per acre (0.03 - 0.05 lb a.i./A).				

(continued)

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Beans, Snap Also known as: (blue lake, bush, common, edible-podded, filet, flageolet, French, French horticultural, frijoles comunes, garden, green, haricot, haricot commun, Italian, judia comum, Kentucky wonder, magic, pole, romano, string, succulent, vainica, wax)	Leafhoppers (except CA) Mexican Bean Beetle Saltmarsh Caterpillar	0.015 - 0.03	2.9 - 5.8	44 - 22	3
	Beet Armyworm* Cabbage Looper Corn Earworm Corn Rootworm (adults) Cucumber Beetle Cucumber Beetle (adults) Cutworm (seedling spray) European Corn Borer Flea Beetle Grasshoppers Green Cloverworm Leafhopper Leafminer (Guam only) Pea Aphid Potato Leafhopper Soybean Aphid (except CA) Velvet Bean Caterpillar Western Bean Cutworm	0.03 - 0.05	5.8 - 9.6	22 - 13	
*Aids in control. DO NOT apply more than 0.2 lb. a.i. per acre per season. DO NOT allow livestock to graze treated bean fields. DO NOT harvest treated bean vines for livestock forage, fodder, or hay.					
Broccoli (including chinese broccoli), Cabbage, Cauliflower, Chinese Cabbage (tight headed varieties only, e.g. Napa cabbage)	Imported cabbageworm	0.015 - 0.03	2.9 - 5.8	44 - 22	3
	Alfalfa Looper Beet Armyworm* Cabbage Looper Cutworm Flea Beetle Grasshoppers	0.03 - 0.05	5.8 - 9.6	22 - 13	
*Aids in control. DO NOT apply more than 0.4 lb. a.i. per acre per season.					
Carrots	Aster Leafhopper Cutworms Leafhoppers	0.03 - 0.05	5.8 - 9.6	22 - 13	7
	Carrot Weevil	0.05	9.6	13	
DO NOT apply more than 0.5 lb. a.i. per acre per season. For aerial application apply in a minimum of 5 gals. water per acre. Thorough spray coverage of crown area is essential. Use of ground application is recommended. Carrot Weevil - Begin treatment when weevils become active.					

(continued)

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Collards	Alfalfa Looper Beet Armyworm* Cabbage Looper Cutworm Flea Beetle Grasshopper Imported Cabbageworm	0.03 - 0.05	5.8 - 9.6	22 - 13	7
	*Aids in control. DO NOT feed livestock on treated plant parts. DO NOT apply more than 0.2 lb. a.i. per acre per season. For aerial application apply in a minimum of 5 gals. water per acre.				
Cucumber, Melons (cantaloupe, honeydew melons, muskmelon, watermelon), Pumpkin, Squash (summer, winter)	Cabbage Looper Corn Earworm Cucumber Beetle (adults) Cutworms (seedling spray) Grasshoppers Leafhoppers Pickleworm Plant Bugs (Lygus Bugs, Stink Bugs) Rindworms Squash Bug Squash Vine Borer	0.03 - 0.05	5.8 - 9.6	22 - 13	3
	DO NOT apply more than 0.25 lb. a.i. per acre per season.				
Eggplant	Colorado Potato Beetle Corn Earworm European Corn Borer Flea Beetles Loopers	0.03 - 0.05	5.8 - 9.6	22 - 13	7
	Apply when insects are observed or when insect damage is observed. Repeat applications at 7 to 10 day intervals to achieve control. DO NOT apply more than 0.35 lb. a.i. per acre per season.				
Kohlrabi	Cabbage Looper	0.03 - 0.05	5.8 - 9.6	22 - 13	3
	DO NOT apply more than 0.4 lb. a.i. per acre per season.				
Lentils	See Beans, Dry				
Lettuce, Head AZ CA, CO, FL, NM & TX ONLY	Alfalfa Looper Beet Armyworm* Cabbage Looper <i>Heliothis</i> spp.	0.025 - 0.05	4.8 - 9.6	26 - 13	7
	*Aids in control. DO NOT apply more than 0.35 lb. a.i. per acre per season.				
Mustard Greens	Cabbage Looper Imported Cabbageworm	0.05	9.6	13	7
	DO NOT apply more than 0.2 lb. a.i. per acre per season.				
Okra (FL only)	Cabbage Looper Corn Earworm Southern Armyworm	0.03 - 0.05	5.8 - 9.6	22 - 13	1
Peas, Dry	See Beans, Dry				

(continued)

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Peas, Green	Green Cloverworm Pea Aphid	0.015 - 0.03	2.9 - 5.8	44 - 22	3
	Alfalfa Caterpillar Alfalfa Looper Armyworm Cabbage Looper Celery Looper Corn Earworm Cutworms Imported Cabbageworm	0.03 - 0.05	5.8 - 9.6	22 - 13	
	Pea Leaf Weevil (ID, OR & WA only) Pea Weevil (ID, OR & WA only)	0.025 - 0.05	4.8 - 9.6	26 - 13	
	DO NOT apply more than 0.1 lb. a.i. per acre per season. DO NOT feed treated pea vines to livestock. Pea Weevil & Pea Leaf Weevil (ID, OR & WA only) - Time of application is critical in achieving control of pea weevil. For optimum results, apply at bloom prior to detecting adult pea weevils. Once adult pea weevil populations reach a level of 2 or more adults per 25 sweeps, control may be reduced.				
Pepper	Beet Armyworm* Colorado Potato Beetle Corn Earworm Cucumber Beetle (adults) European Corn Borer Flea Beetles Loopers Pepper Weevil* Southern Armyworm	0.03 - 0.05	5.8 - 9.6	22 - 13	7
*Aids in control. Apply when insects are observed or when insect damage is observed. Repeat applications at 7 to 10 day intervals to achieve control. DO NOT apply more than 0.35 lb. a.i. per acre per season.					

(continued)

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Potato	Leafhoppers (except CA) Potato Psyllid	0.015 - 0.03	2.9 - 5.8	44 - 22	7
	Potato Tuberworm	0.015 - 0.05	2.9 - 9.6	44 - 13	
	Beet Armyworm* Buckthorn Aphid Cabbage Looper Colorado Potato Beetle Cucumber Beetle (adult control) Cutworms European Corn Borer Flea Beetles Grasshoppers Potato Aphid Potato Leafhopper Tarnished Plant Bug Western Yellow-Striped Armyworm	0.03 - 0.05	5.8 - 9.6	22 - 13	
	Colorado Potato Beetle (Long Island, NY only)	0.05	9.6	13	
	Grasshopper - For control of first and second instar grasshopper nymphal stages a rate range of 3.9 to 5.8 fluid ounces of product per acre (0.02-0.03 lb. a.i./A) can be used. Correct timing of spray applications to the first and second instar nymphal stages and thorough coverage is critical to achieve optimum control. For grasshopper nymph stages larger than second instar, use Trikon at use rates of 5.8 to 9.6 fluid ounces of product per acre (0.03 - 0.05 lb. a.i./A). Potato Tuberworm - For control of Potato Tuberworm apply Trikon when tuberworm larvae and/or moth counts reach locally established treatment threshold populations. Repeat applications of effective insecticides may be needed to keep tuberworm larvae populations as low as possible prior to harvest in order to reduce the risk of tuber damage. Failure to adequately control tuberworm larvae prior to crop senescence or vine kill increases the risk of tuber damage. *Aids in control. DO NOT apply more than 0.35 lb. a.i. per acre per season.				
Radishes	Armyworms Beetles	0.03 - 0.05	5.8 - 9.6	22 - 13	7
	DO NOT apply more than 0.1 lb. a.i. per acre per season.				

(continued)

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Sweet Corn	Western Bean Cutworm	0.015 - 0.03	2.9 - 5.8	44 - 22	1
	For additional information, consult Directions For Use under Corn (field) .				
	Armyworm Banded Cucumber Beetle Beet Armyworm* Chinch Bugs Corn Earworm Corn Leaf Aphid Corn Rootworms (adults) Cutworms European Corn Borer Fall Armyworm (except CA) 1 st and 2 nd instar Flea Beetles Grasshoppers Oat Bird-Cherry Aphid Sap Beetles (adults) Southwestern Corn Borer Stalk Borer Tarnished Plant Bug	0.03 - 0.05	5.8 - 9.6	22 - 13	
	Corn Silkwfly (except CA)**	0.05	9.6	13	
	*Aids in control. **Suppression only. For Ear Protection - Begin applications either just before or at time of silking. For additional information consult directions for use under Corn (field) . Corn Leaf Aphid & Oat Bird-Cherry Aphid - For optimum results, direct the spray at the aphid population so as to achieve maximum coverage of the exposed insects. Aphids not contacted by the spray, such as in whorls and leaf axils, may not be adequately controlled. Corn Silkwfly (except CA) - Direct application to the ear zone to obtain thorough coverage of the corn silk. Fall Armyworm (except CA) - 1 st and 2 nd instar fall armyworm only. Direct the application to the ear zone to obtain thorough coverage of the corn silk. DO NOT apply more than 0.5 lb. a.i. per acre per season.				
Tomato	Tobacco Hornworm Tomato Hornworm	0.015 - 0.03	2.9 - 5.8	44 - 22	1
	Beet Armyworm* Cabbage Looper Colorado Potato Beetle Cutworms Flea Beetle Grasshoppers Potato Aphid Southern Armyworm Tomato Fruitworm Tomato Pinworm Western Yellow-Striped Armyworm Whitefly	0.03 - 0.05	5.8 - 9.6	22 - 13	
	Vegetable Leafminer**	0.05	9.6	13	
	*Aids in control. ** Trikon is not recommended for use on the Vegetable Leafminer in Florida. DO NOT apply more than 0.5 lb. a.i. per acre per season.				

(continued)

CROP	INSECT	APPLICATION RATE		ACRES TREATED PER GAL. OF TRIKON	LAST APPLICATION (DAYS TO HARVEST)
		LB. A.I./A	FL. OZ./A		
Turnips	Armyworm Flea Beetle Imported Cabbageworm	0.03 – 0.05	5.8 – 9.6	22 - 13	7
	DO NOT apply more than 0.4 lb. a.i. per acre per season.				

SPECIALTY USES

CROP	INSECT	APPLICATION RATE		
		HIGH VOLUME SPRAYS		LOW VOLUME SPRAYS LB. A.I./A
		LB. A.I./100 GAL.	FL. OZ./100 GAL.	
Christmas tree plantings, Conifer plantations, Conifer seed orchards, Forest tree nurseries	Balsam Twig Aphid Balsam Woolly Adelgid Cranberry Girdler (adult control) European Pine Sawfly Nantucket Pine Tip Moth and other Pine Tip Moths (except CA) Northern Pine Weevil Pales Weevil Pine Chafer Pine Conelet Bug Pine Needle Midge Pineleaf Chermid Red Pine Sawfly Redheaded Pine Sawfly Spittlebugs Spruce Budworm	0.03 - 0.05	5.8 - 9.6	0.03 - 0.05
	Spray in sufficient gallonage to obtain good coverage of entire tree.			
	Coneworm Seed Chalcid Seedbug	9.6 fl. oz./100 gals. water for high volume sprayers. 52 fl. oz./100 gals. water for low volume sprayers. 0.19 lb. a.i./acre application in not less than 10 gals. of water for aerial applications.		
	Apply first application within 1 week of female flower closure or peak pollen flight for Webbing Coneworm control. For other Coneworms and Seedbugs , apply first application within 30 days following female flower closure. Repeat application at intervals of 4 weeks but DO NOT apply more than 1.6 lbs. a.i. per acre per year. For Seed Chalcid control, apply when all cones are pendant, and repeat at 1 - 2 week intervals for 2 or more sprays. Apply approximately 5 - 10 gals. of the 9.6 fl. oz./100 gals. dilution per tree with high volume sprayers. With low volume sprayers apply 100 gals. of the 52 fl. oz./100 gals. dilution per acre. DO NOT graze or harvest cover crop. Refer to Spray Recommendations and Precautions when applying to areas adjacent to water.			
NON-CROPLAND (Excluding Public Land Such As Forests, Parks, Or Recreational)	Grasshoppers Saltmarsh Caterpillar	0.015 - 0.03	2.9 - 5.8	44 - 22
	Army Cutworms Armyworms Chinch Bugs	0.03 - 0.05	5.8 - 9.6	22 - 13
	Spray non-cropland adjacent to tilled areas to control migrating insects (Grasshoppers , Armyworms) which are a threat to crops. DO NOT apply more than 0.5 lb. active ingredient per acre per year. DO NOT feed treated crop to livestock. Refer to Spray Recommendations and Precautions when applying to areas adjacent to water.			

STORAGE AND DISPOSAL

DO NOT pour or dispose down-the-drain or sewer. Call your local solid waste agency for local disposal options.



DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in cool dry place in original container. Keep container closed when not in use. Do not store near food or feed. Do not use or store around the home. Avoid contact with water. In case of spill or leak, soak up with sand, earth or synthetic absorbent (do not use alkaline absorbents) and dispose of wastes in compliance with local, State and Federal regulations.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Less Than or Equal to 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

Greater Than 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by other procedures allowed by State and local authorities.

For Bulk and Mini-Bulk Containers: Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

PRIMEROS AUXILIOS	
SI SE INGIERE:	• Llame de inmediato a un centro de control de envenenamientos o a un médico para consejo de tratamiento. • NO induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. • NO administre ningún líquido a la persona. • NO administre nada por boca a una persona que haya perdido el conocimiento.
SI CAE EN LA PIEL O LA ROPA:	• Quítese la ropa contaminada. • Enjuague la piel inmediatamente con bastante agua por 15 - 20 minutos. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
SI ENTRA EN CONTACTO CON LOS OJOS:	• Mantenga los ojos abiertos y enjuáguelos lenta y cuidadosamente con agua, durante 15 a 20 minutos. • Si utiliza lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
NOTA AL MÉDICO	
Contiene un destilado de petróleo. El vómito puede causar neumonía por aspiración. Si la zona está en contacto con la piel, después de secarla, aplique crema o aceite de vitamina E, si dispone de él. Si no dispone de él, aplique aceite vegetal generosamente sobre las zonas doloridas. Puede usar el aceite o la crema repetidamente hasta obtener alivio.	
NÚMERO DE TELÉFONO DIRECTO	
Cuando llame a un centro de control de envenenamiento, o a un médico, o intente obtener tratamiento, tenga a la mano el recipiente o la etiqueta del producto. Para información emergencia sobre este producto, llame al centro de control de envenenamientos a 1-800-222-1222 . Para información general sobre este producto, contacte al Centro Nacional de Información de Pesticidas (NPIC, por sus siglas en inglés) a 1-800-858-7378 , Lunes a Viernes, 8 AM a 12 PM PST (hora del Pacífico), o a http://npic.orst.edu .	

DECLARACIONES PREVENTIVAS
RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS
AVISO

Puede ser letal si se ingiere. Dañino si se absorbe por la piel. Causa irritación moderada en los ojos. **NO** permita que caiga en los ojos, la piel o la ropa. Lávese con abundante agua y jabón luego de manejar (pesticidas) y antes de comer, beber, masticar chicle, utilizar tabaco, o usar al baño. Quítese y lave la ropa contaminada antes de reutilizarla.

EQUIPO DE PROTECCIÓN PERSONAL (PPE, por sus siglas en inglés)

Los aplicadores y otros manipuladores de pesticidas (plaguicidas) deberán usar:

- Camisa de manga larga y pantalones largos
- Guantes resistentes a productos químicos fabricados con laminados impermeables, goma de butilo ≥ 14 mils, goma de nitrilo ≥ 14 mils, cloruro de polivinilo ≥ 14 mils, o Viton ≥ 14 mils
- Zapatos y calcetines
- Lentes o gafas protectoras

Deseche la ropa y otros materiales absorbentes que hayan sido empapados o muy contaminados con el concentrado de este producto. **NO** vuelva a reutilizarlos. Siga las instrucciones del fabricante para la limpieza/mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE separadamente de otra ropa para lavar.

DECLARACIÓN DE CONTROL TÉCNICO

Cuando los manipuladores de pesticidas usen sistemas cerrados, cabinas, o aeronaves, de manera que cumplan con los requisitos enumerados en la Ley de Protección al Trabajador (WPS, por sus siglas en inglés) [40 CFR170.240 (d) (4-6)], para pesticidas agrícolas, los requisitos de equipo de protección personal (PPE) para el manipulador de pesticidas pueden reducirse o modificarse como está especificado en la WPS.

RIESGOS AMBIENTALES

Este pesticida es extremadamente tóxico para peces e invertebrados acuáticos. **NO** aplique el producto directamente al agua, a zonas donde haya presencia de agua superficial ni a las zonas intermareales por debajo de la marca de pleamar media. **NO** aplicar cuando las condiciones meteorológicas favorezcan la dispersión del producto desde las áreas tratadas. La deriva o el rociado y las escorrentías pueden ser peligrosas para los organismos acuáticos en aguas adyacentes a las áreas tratadas. **NO** contaminar el agua al limpiar el equipo o al desechar las aguas de lavado.

Este producto es altamente tóxico para las abejas expuestas al tratamiento directo, o a los residuos en cultivos en floración o malezas. **NO** aplique este producto ni deje que se disperse sobre cultivos en flor o malas hierbas mientras las abejas estén visitando activamente la zona de tratamiento.

DECLARACIÓN CONSULTIVA SOBRE ORGANISMOS NO OBJETIVO

Este producto es altamente tóxico para las abejas expuestas al tratamiento directo, o a los residuos en cultivos en floración o malezas. **Proteja a los insectos polinizadores siguiendo las instrucciones de la etiqueta destinadas a minimizar la deriva y reducir el riesgo para estos organismos.**

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal.

ALMACENAMIENTO Y DESECHO

NO verter ni desechar en el desagüe ni en el alcantarillado. Llame a su agencia local de residuos sólidos para conocer las opciones de eliminación local.



NO contamine agua, comida ni forrajes mediante el almacenamiento y desecho.

ALMACENAMIENTO DE PESTICIDAS: Conservar en un lugar fresco y seco en su envase original. Mantenga el envase cerrado cuando no lo use. **NO** almacenar cerca de forraje o alimentos. **NO** utilizar ni almacenar en el hogar. Evite el contacto con el agua. En caso de derrame o fuga, absorba con arena, tierra o absorbente sintético (no utilice absorbentes alcalinos) y deseche los residuos de conformidad con las regulaciones locales, estatales y federales.

DESECHO DE PESTICIDAS: Los residuos resultantes del uso de este producto pueden eliminarse en el sitio o en una instalación de eliminación de residuos aprobada.

MANIPULACIÓN DE ENVASES:

Menor o igual a 5 galones: Recipiente no rellenable. **NO** reutilice ni rellene este recipiente. Ofrezca para el reciclaje si está disponible. Enjuague tres veces (o equivalente) el recipiente inmediatamente después de vaciarlo. Enjuague tres veces siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezclar, y drene por 10 segundos después de que el flujo empiece a gotear. Llene el recipiente 1/4 lleno con agua y tape de nuevo. Agite por 10 segundos. Eche el enjuague en el equipo de aplicación o tanque de mezclar, o almacene el enjuague para uso más tarde o eliminación. Drene por 10 segundos después de que el flujo empiece a gotear. Repita este procedimiento dos veces más. Luego ofrezca para el reciclaje si está disponible o el reacondicionado si está adecuado, o perforfe y deseche en un área adecuada para desechos sanitarios o por otros procedimientos aprobados por las autoridades estatales y locales.

Más de 5 galones: Recipiente no rellenable. **NO** reutilice ni rellene este recipiente. Ofrezca para el reciclaje si está disponible. Enjuague tres veces (o equivalente) el recipiente inmediatamente después de vaciarlo. Enjuague tres veces siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezclar. Llene el recipiente 1/4 lleno con agua. Reemplace y asegure los cierres. Vuelque el recipiente al lado y hágalo rodar por 30 segundos, asegurando por lo menos una revolución completa. Ponga el recipiente vertical e inclínelo hacia adelante y hacia atrás varias veces. Dé vuelta al recipiente hacia el lado opuesto e inclínelo hacia adelante y hacia atrás varias veces. Eche el enjuague en el equipo de aplicación o tanque de mezclar, o almacene el enjuague para uso más tarde o eliminación. Repita este procedimiento dos veces más. Deseche el recipiente vacío en un vertedero sanitario o mediante otros procedimientos aprobados por las autoridades locales y estatales.

Para Contenedores y mini-contenedores a granel: Recipiente rellenable. Rellene este recipiente únicamente con pesticida. **NO** reutilice este recipiente para ningún otro propósito. La limpieza del recipiente antes de su eliminación final es la responsabilidad de la persona que se deshace del recipiente. La limpieza antes de rellenar es la responsabilidad de la persona que lo va a rellenar. Para limpiar el recipiente antes de su eliminación final, vacíe el contenido restante de este recipiente en el equipo de aplicación o en el tanque de mezcla. Llene el recipiente con un 10% de agua. Agitar vigorosamente o haga recircular el agua con la bomba durante 2 minutos. Vierta o bombee el agua de enjuague en el equipo de aplicación o en el sistema de recolección de agua de enjuague. Repita este procedimiento dos veces más. Luego ofrezca para el reciclaje si está disponible o perforfe y deseche en un área adecuada para desechos sanitarios, o por otros procedimientos aprobados por las autoridades estatales y locales o la incineración.

NOTES

RESTRICTED USE PESTICIDE DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

ESFENVALERATE GROUP 3A INSECTICIDE

Trikon

For the control of insect pests on Fields Crops, Vegetable Crops, Fruit Crops, Tree Nut Crops.

ACTIVE INGREDIENT:	By Weight %
Esfenvalerate: (S)-cyano (3-phenoxyphenyl) methyl (S)-4-chloro-alpha-(1-methylethyl) benzeneacetate	8.4%
OTHER INGREDIENTS*	91.6%
TOTAL:	100.0%

Contains 0.66 lb. active ingredient per gallon.

*Contains petroleum distillate

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	• Immediately call a poison control center or doctor for treatment advice. • DO NOT induce vomiting unless told by a poison control center or doctor. • DO NOT give any liquid to the person. • DO NOT give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN	
Contains petroleum distillate. Vomiting may cause aspiration pneumonia. If on skin, after drying, apply Vitamin E cream or oil, if available. If not available, apply vegetable oil liberally over painful areas. The oil or cream may be used repeatedly until relief is achieved.	
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222. For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	

Manufactured For: Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 82633-111-83529

EPA Est. No. **AG** 72159-GA-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING/AVISO

May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. **DO NOT** get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when cleaning equipment or when disposing of equipment wash-waters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

NON-TARGET ORGANISM ADVISORY:

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or to residues in/on blooming crops or weeds. **Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all **Directions for Use** carefully before applying.

STORAGE AND DISPOSAL

DO NOT pour or dispose down-the-drain or sewer. Call your local solid waste agency for local disposal options. **DO NOT** contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Store in cool dry place in original container. Keep container closed when not in use. Do not store near food or feed. Do not use or store around the home. Avoid contact with water. In case of spill or leak, soak up with sand, earth or synthetic absorbent (do not use alkaline absorbents) and dispose of wastes in compliance with local, State and Federal regulations. **PESTICIDE DISPOSAL:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Less Than or Equal to 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by State and local authorities. **Greater Than 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by other procedures allowed by State and local authorities. **For Bulk and Mini-Bulk Containers:** Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

See inside label booklet for additional Precautionary Statements and Directions For Use.

Net Contents: **2.5 Gals.*** ☐ **265 Gals.**

* Unless alternate checked