

ACEQUINOCYL GROUP 20B ACARICIDE

Actor SC Miticide

INSECTICIDE

For the Control of Mites on: Small Fruit Vine Climbing except
Fuzzy Kiwifruit (Crop Subgroup 13-07F) and Tree Nut (Crop Group 14-12).

ACTIVE INGREDIENT:	WT. BY %
Acequinocyl: 3-dodecyl-1, 4-dihydro-1, 4-dioxo-2-naphthyl-acetate	30.0%
OTHER INGREDIENTS:	70.0%
TOTAL:	100.0%
This product contains 2.5 lbs. a.i. per gallon.	

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

See label booklet for complete First Aid, Precautionary Statements,
Directions For Use, and Storage and Disposal.

Manufactured For:

Sharda Cropchem Ltd. 

2nd Floor, Prime Business Park,
Dashrathlal Joshi Rd. Vile Parle (West),
Mumbai - 400056, India

EPA Reg. No. 82633-151

EPA Est. No.  72159-GA-001;  83411-MN-001;  89332-GA-001;
 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that
match the first two letters in the batch number.

Net Contents: **2.5 Gallons**

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBERS	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222. For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu.</p>	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Harmful if swallowed. Harmful if inhaled.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks
- Chemical-resistant gloves including barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride ≥ 14 mils, or viton ≥ 14 mils

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and change into clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates such as Eastern oysters and mysid shrimp. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate.

Surface Water Statement

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several days after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Carefully read and follow the Directions for Use and restrictions before applying this product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow entry into treated areas during restricted-entry interval (REI) of **12 hours**.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water is:

- Coveralls
- Shoes and socks
- Chemical-resistant gloves

MANDATORY SPRAY DRIFT MANAGEMENT

AIRBLAST APPLICATIONS:

- Spray must be directed into the canopy.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outwind pointing nozzles at row ends and when spraying outer rows.
- Do not apply during temperature inversions.

GROUND BOOM APPLICATIONS:

- User must only apply with the release height recommended by the manufacturer, but no more than 3 ft. above the ground or crop canopy.
- Applicators are required to use a fine or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size — Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT — Ground Boom

For ground equipment, the booms should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and a common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.
- Boomless Ground Applications: Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

PRODUCT INFORMATION

Actor SC Miticide is a suspension concentrate containing the active ingredient acequinocyl. When applied to the foliage as directed, **Actor SC Miticide** provides knock-down and residual control. **Actor SC Miticide** is not expected to have cross-resistance to existing acaricides/miticides and is very effective against various species mites including those resistant to other miticides. **Actor SC Miticide** is relatively harmless to most predacious mites and beneficial insects, making it compatible with IPM (Integrated Pest Management) and resistance management programs.

Restrictions (For All Crops):

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply by air.
- **DO NOT** apply within 75 feet of aquatic areas.
- Not for residential use.
- A 30-day plant-back interval is required for all crops not listed on this label.
- **DO NOT** combine **Actor SC Miticide** with Phos-ethyl (Fosetyl).

RESISTANCE MANAGEMENT GUIDANCE

For resistance management, **Actor SC Miticide** contains a Group 20B acaricides. Any mite population may contain individuals naturally resistant to **Actor SC Miticide** and other Group 20 acaricides. The resistant individuals may dominate the mite population if this group of acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay acaricide resistance, take the following steps:

- Rotate the use of **Actor SC Miticide** or other Group 20 acaricides within a growing season, or among growing seasons, with different groups that control the same pests. It is recommended not use **Actor SC Miticide** successive applications. Rotation with other acaricides having different modes of action is recommended.
- Use tank-mixtures with acaricides not in Group 20 when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for mites between the individual components of a mixture.
- In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
 - Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact your Sharda Cropchem Limited representative.

TANK MIXING INFORMATION

It is the pesticide user's responsibility to ensure that all products in the mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions and precautionary statement of each product in the tank mixture.

Actor SC Miticide can be combined with most commonly used pesticides with the exception of strongly alkaline material, which can reduce the activity of this product. **DO NOT** combine **Actor SC Miticide** with Phos-ethyl (Fosetyl). To obtain broad-spectrum pest control, **Actor SC Miticide** can be tank-mixed with most other miticide/insecticide products. However, all plant species and their varieties and cultivars have not been tested with possible tank-mix combinations, sequential pesticide treatments and adjuvants and surfactants. The user should conduct a spray mix compatibility and phytotoxicity trial under local conditions to ensure compatibility prior to any large-scale use.

Preparation of Spray Solution

Begin with clean equipment. Add sufficient clean water to the spray-tank for one-half of the mix load. Start tank agitation. Add and properly suspend the necessary amount of product before adding any other products. Agitate to ensure thorough mixing while adding the remaining required water. If allowed for use on your specific crop a high-quality wetting agent or other agricultural spray adjuvant may be added to spray solutions according to the manufacturer's use directions. **Actor SC Miticide** must be properly suspended and diluted prior to the addition of any adjuvant. Consult the adjuvant label or manufacturer for crop tolerance and safety information when used with this product. Maintain agitation during mixing and application. Allowing the spray mixture to stand without agitation may result in an improperly suspended mixture. If agitation is stopped for any reasons, thoroughly re-mix the spray solution prior to any further use. Apply with properly calibrated spray equipment.

APPLICATION INFORMATION

Apply **Actor SC Miticide** when threshold populations are observed and before large mite populations have been established.

Apply **Actor SC Miticide** in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for optimum control. To achieve adequate coverage, use proper spray pressures, nozzles, nozzle spacing, and water volume per acre.

When used as directed, **Actor SC Miticide** is effective for the control of a variety of listed mite species. **Actor SC Miticide** shows efficacy on all mite life stages of susceptible mites. Using **Actor SC Miticide** in successive miticide applications is not advised. Use **Actor SC Miticide** as part of a sound resistance management program that includes rotation with other miticide treatments having different modes of action.

ROTATIONAL CROP RESTRICTIONS

Food and feed crops not listed on this label must not be planted within 30 days of the last **Actor SC Miticide** application.

SPECIFIC USE DIRECTIONS

Small Fruit Vine Climbing except Fuzzy Kiwifruit (Subgroup 13-07f)

Amur River Grape; Gooseberry; Grape; Kiwifruit, Hardy; Maypop; Schisandra Berry; Cultivars, Varieties, and/or Hybrids of these.

Pest	Use Rate per Acre
Pacific Spider Mite (<i>Tetranychus pacificus</i>) Two-Spotted Spider Mite (<i>Tetranychus urticae</i>) Willamette Spider Mite (<i>Eotetranychus willamettei</i>)	16 fl. oz. (0.3 lb. a.i.)
<p>Restrictions:</p> <ul style="list-style-type: none"> • For Grapes (young vines, early season application prior to dense foliage) - DO NOT use less than 50 gals. of water per acre. • DO NOT use less than 100 gals. of water volume per acre (except for grapes). • Apply to every row to ensure coverage. DO NOT use in alternate row middle applications. • Allow a minimum of 21 days between treatments. • DO NOT make more than 2 applications of Actor SC Miticide per year. • DO NOT apply more than 32 fl. oz. (0.6 lb. a.i.) of Actor SC Miticide per acre per year. • Pre-harvest interval: 7 days • Restricted-entry Interval (REI): 12 hours 	

Tree Nut (Crop Group 14-12)

African Nut-Tree; Almond; Beechnut; Brazil Nut; Brazilian Pine; Bunya; Bur Oak; Butternut; Cajou Nut; Candlernut; Cashew; Chestnut; Chinquapin; Coconut; Coquito Nut; Dika Nut; Ginkgo; Guiana Chestnut; Hazelnut (Filbert); Heartnut; Hickory Nut; Japanese Horse-Chestnut; Macadamia Nut; Mongongo Nut; Monkey-Pot; Monkey Puzzle Nut; Okari Nut; Pachira Nut; Peach Palm Nut; Pecan; Pequi; Pili Nut; Pine Nut; Pistachio; Sapucaia Nut; Tropical Almond; Walnut, Black; Walnut, English; Yellowhorn; Cultivars, Varieties, and/or Hybrids of these.

Pest	Use Rate per Acre
Pacific Spider Mite (<i>Tetranychus pacificus</i>) Two-Spotted Spider Mite (<i>Tetranychus urticae</i>) Willamette Spider Mite (<i>Eotetranychus willamettei</i>)	16 fl. oz. (0.3 lb. a.i.)
<p>For optimal control, use 150 gallons of water per acre or higher.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT use less than 100 gals. of water volume per acre. Apply to every row to ensure coverage. DO NOT use in alternate row middle applications. • Allow a minimum of 21 days between treatments. • DO NOT make more than 2 applications of Actor SC Miticide per year. • DO NOT apply more than 32 fl. oz. (0.6 lb. a.i.) of Actor SC Miticide per acre per year. • Pre-harvest interval: 7 days • Restricted-entry Interval (REI): 12 hours 	

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep tightly closed. Store in a cool dry place. Protect from excessive heat.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Rigid Plastic Less Than or Equal to 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

Rigid Plastic Greater Than 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of empty container in a sanitary landfill or by other procedures allowed by State and local authorities.

Rigid Plastic Bulk and Mini-Bulk Containers: Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda Cropchem Limited or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda Cropchem Limited and Seller harmless for any claims relating to such factors.

Sharda Cropchem Limited warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda Cropchem Limited and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA CROPChem LIMITED MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda Cropchem Limited nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA CROPChem LIMITED AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA CROPChem LIMITED OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda Cropchem Limited and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda Cropchem Limited.

All trademarks are the property of their respective owners.

ACEQUINOCYL GROUP 20B ACARICIDE

Actor SC Miticide

For the Control of Mites on: Small Fruit Vine Climbing except Fuzzy Kiwifruit (Crop Subgroup 13-07F) and Tree Nut (Crop Group 14-12).

ACTIVE INGREDIENT:	WT. BY %
Acequinocyl: 3-dodecyl-1, 4-dihydro-1, 4-dioxo-2-naphthyl-acetate.	30.0%
OTHER INGREDIENTS:	70.0%
TOTAL:	100.0%

This product contains 2.5 lbs. a.i. per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by the poison control center or doctor. DO NOT give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBERS

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at **1-800-222-1222**. For general information about this product, contact the National Pesticides Information Center (NPIC) at **1-800-858-7378**, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Harmful if swallowed. Harmful if inhaled.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates such as Eastern oysters and mysid shrimp. **DO NOT** apply directly to water, or to areas where surface water is present or is intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate.

Surface Water Statement

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several days after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep tightly closed. Store in a cool dry place. Protect from excessive heat.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Rigid Plastic Less Than or Equal to 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

See label booklet for additional Precautionary Statements and Directions For Use.

Manufactured For: Sharda Cropchem Ltd

2nd Floor, Prime Business Park, Dashrathlal Joshi Rd. Vile Parle (West), Mumbai - 400056, India

EPA Reg. No. 82633-151

EPA Est. No. **AG** 72159-GA-001; **MA** 83411-MN-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: **2.5 Gallons**

OPEN HERE