

Sojourn 50 T&O

FUNGICIDE

Turfgrass and Ornamental Fungicide

A SYSTEMIC FUNGICIDE FOR CONTROL OF THE LISTED TURFGRASS AND ORNAMENTAL DISEASES

ACTIVE INGREDIENT:	WT. BY %
Metconazole: 5-[(4-chlorophenyl)methyl]-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol	50.0%
OTHER INGREDIENTS:	50.0%
TOTAL:	100.0%
Sojourn 50 T&O is a water dispersible granule containing 50% active ingredient.	

KEEP OUT OF REACH OF CHILDREN / MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand this label, find someone to explain it to you in detail.)

See inside label booklet for complete First Aid, Precautionary Statements,
Directions For Use, and Storage and Disposal.

EPA Reg. No. 82633-122-83529

EPA Est. No. (AG) 72159-GA-001; (MC) 89332-GA-001; (SC) 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Manufactured For:

Sharda USA LLC



7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

Net Contents: **5 lbs.**

FIRST AID	
IF SWALLOWED:	• Immediately call a poison control center or doctor. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION/PRECAUCIÓN

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin, and clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Coveralls, long-sleeved shirt, and long pants
- Chemical-resistant gloves of any waterproof material including barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber (includes natural rubber blends and laminates) ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils
- Shoes plus socks

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, mammals, fish, and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory

This product may impact surface water quality through spray and runoff of rain water. This product has a high potential for runoff for several months or more after application. Poorly draining soils or soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied, and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. **DO NOT** discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls, long-sleeved shirt, and long pants
- Chemical-resistant gloves made of any waterproof material including barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber (includes natural rubber blends and laminates) ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils
- Socks and shoes

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to the uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, greenhouses, or sod farms.

- Keep all unprotected persons out of operating areas or vicinity where there may be drift. **DO NOT** enter or allow others to enter treated areas until sprays have dried.

PRODUCT INFORMATION

Sojourn 50 T&O is formulated as 50% water dispersible granular (WDG). The active ingredient in **Sojourn 50 T&O** is metconazole, a broad-spectrum triazole fungicide that works by inhibiting demethylation and other processes in sterol biosynthesis.

Sojourn 50 T&O is systemic and is quickly absorbed into plant tissue.

Sojourn 50 T&O displays preventive and curative properties.

Sojourn 50 T&O may be used for disease control in turfgrass on golf courses, lawns and landscape areas around residential, institutional, public, commercial, and industrial buildings, parks, recreational areas, athletic fields, and sod farms.

Sojourn 50 T&O may be used for disease control on ornamental plants in commercial indoor and outdoor nurseries, greenhouses and on ornamental plants found in landscaped areas around residential, institutional, public, commercial, and industrial buildings, parks, recreational areas, and athletic fields.

Sojourn 50 T&O provides optimal disease control when applied in combination or as part of a rotation in a regularly scheduled spray program with other effective fungicides that have different modes of action (i.e., non-Group 3 fungicides).

MODE OF ACTION

The active ingredient in **Sojourn 50 T&O**, metconazole, belongs to the sterol biosynthesis inhibitor group of fungicides as classified by the U.S. EPA and Canada PMRA as a target site of action Group 3 fungicide.

RESISTANCE MANAGEMENT

For resistance management, **Sojourn 50 T&O** contains a Group 3 fungicide/bactericide. Any fungal/bacterial population may contain individuals naturally resistant to **Sojourn 50 T&O** and other Group 3 fungicides/bactericides. A gradual or total loss of pest control may occur over time if these fungicides/bactericides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of **Sojourn 50 T&O** or other Group 3 fungicides/bactericides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological, and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For information or to report suspected resistance contact *Sharda USA LLC* at <https://shardausa.com>. You can also contact your pesticide distributor or university extension specialist to report resistance.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND SOJOURN 50 T&O

A jar test should be performed before mixing commercial quantities of **Sojourn 50 T&O**, when using this product for the first time, when using new adjuvants, when using new tank mixes, or when using a new water source.

1. Add 1 pt. of the water to a quart jar. Use water from the same source and temperature as water that will be used in the spray tank mixing operation.
2. Add about 1 teaspoon (2.4 grams) of **Sojourn 50 T&O** to the quart jar; gently mix until product goes into suspension.
3. Add appropriate amount of each tank mix adjuvant or partner.
4. Place cap on jar, invert 10 times, let stand for 15 minutes and evaluate.
5. An ideal tank mix combination will be uniform and free of large suspended particles. If any of the following conditions are observed, reevaluate the choice of the adjuvant:
 - a. Layer of oil or globules on the mixture's surface.
 - b. Flocculation: fine particles in suspension or as a layer on the bottom of the jar.
 - c. Clabbering: Thickening texture (coagulated) like gelatin.

TANK MIX PARTNERS

Sojourn 50 T&O is compatible with most herbicides, insecticides, fungicides, and other types of tank mix components. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. **DO NOT** tank mix with any product which has a prohibition on tank mixing. When evaluating potential tank mix partners, always apply to a small area of target plants and then monitor for adverse effects for at least two weeks before making large-scale applications.

SPRAYER PREPARATION

Before applying **Sojourn 50 T&O**, start with clean, well maintained application equipment. The spray tank hoses and booms must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply.

Sojourn 50 T&O. Check nozzles frequently for accuracy. If two or more products were tank mixed prior to **Sojourn 50 T&O** application, follow the most restrictive cleanup procedure.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, slowly add the **Sojourn 50 T&O** to the spray tank. A rippling or rolling action on the water surface will occur as the result of the agitation.
3. If tank mixing **Sojourn 50 T&O** with other labeled pesticides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions.
4. Add any required adjuvants.
5. Fill spray tank to desired level with water. Continue to agitate until all spray solution has been applied.
6. Mix only the amount of spray solution that can be applied the day of mixing. Apply **Sojourn 50 T&O** within 12 hours of mixing.

SPRAYER CLEANUP

Clean the spray equipment each day following **Sojourn 50 T&O** application. After **Sojourn 50 T&O** is applied, follow the steps below to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens, and nozzles.
3. Drain tank completely.
4. Remove all nozzles and screens and rinse them in clean water.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S572 and S641).
- If the windspeed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 - 15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed-wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to select the nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to select the nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572) for all applications.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Boomless Ground Applications

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

TURFGRASS

- Use **Sojourn 50 T&O** for disease control on golf courses, lawns and landscape areas around residential, institutional, public, commercial, and industrial buildings, parks, recreational areas, athletic fields, and sod farms.
- Use **Sojourn 50 T&O** in conjunction with turf management practices that promote good plant health and optimum disease control. Proper identification of disease infestations is essential for providing control. Consult diagnostic guides, kits, or other means of identification to determine the best control measures.
- **Sojourn 50 T&O** is more effective when used preventively or early in disease development. Length of disease control will vary according to local environmental conditions, disease pressure and turfgrass management practices.
- Apply to dry turf or after mowing. For control of foliar diseases, allow turf to dry before irrigation. For control of soilborne diseases, **Sojourn 50 T&O** can be watered in after application.
- **Sojourn 50 T&O** may be tank mixed with other registered turfgrass fungicides. Perform a jar test as describe above to determine compatibility before making applications. Directions for use of **Sojourn 50 T&O** have been developed using no adjuvant in the spray solution.

- **Sojourn 50 T&O** can injure golf course greens that are under high heat (temperatures exceeding 90°F) and/or drought stress. Under high heat and/or drought stress conditions (mid to late summer), **DO NOT** apply more than 16 oz. (0.5 lb. a.i./A) per acre (0.12 oz. (0.004 lb. a.i./A) per 1,000 sq. ft.) every 30 days.
- Important: Under conditions of optimal disease development (such as high humidity and temperature), use higher labeled rates and shorter application intervals. When applications are made curatively, use higher specified rates of **Sojourn 50 T&O** and shorter specified application intervals. Use the higher specified application rate in areas with a history of severe disease pressure.
- Failure to follow label directions may result in turfgrass injury and or inferior disease control.

APPLICATION DIRECTIONS

Apply **Sojourn 50 T&O** using standard, low-pressure spray equipment designed to deliver coarse to medium size spray droplets (**DO NOT** use nozzles designed to deliver fine droplets). Apply in a sufficient volume of water to provide thorough spray coverage and a uniform spray pattern. **DO NOT** position the spray boom more than 20 inches above the turf. For foliar disease, apply 2 gallons of spray solution per 1,000 sq. ft. For crown and root diseases, apply 2 to 4 gallons of spray solution per 1,000 sq. ft. For spot treatment, mix 0.1 to 0.2 oz. (0.003 to 0.006 lb. a.i./A) of **Sojourn 50 T&O** per gallon of water.

TURFGRASS TOLERANCE

Due to the large number of turfgrass types, possible tank mixes, and application techniques, it has not been determined if **Sojourn 50 T&O** can be safely used on all turfgrasses under all conditions. Before making a large-scale application, determine safety in a small test area. Monitor the test area for at least 2 weeks to determine safety under local growing conditions.

TURFGRASS SPECIES

- Annual Bluegrass
- Bentgrass
- Bermudagrass
- Centipedegrass
- Fescue
- Kentucky Bluegrass
- Perennial Ryegrass
- St. Augustinegrass
- Seashore Paspalum
- Zoysia

TURFGRASS RESTRICTIONS

- **DO NOT** apply more than 4 lbs. (2 lbs. a.i.) of **Sojourn 50 T&O** per acre per year. See Table 1 for specific application rates.
- **DO NOT** apply directly to bermudagrass greens.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply by air.
- **DO NOT** graze animals on treated areas for one year after application.
- **DO NOT** feed clippings from treated areas to livestock or poultry.
- **DO NOT** exceed 0.2 oz. (0.006 lb. a.i./A) product per gallon of water for spot treatment.
- **DO NOT** make repeat applications less than 14 days apart.
- Up to 8 applications can be made not exceeding 2 lbs. a.i./A (regardless of Metconazole source).

Table 1 - TURFGRASS – Specific Diseases, Rates, Timings and Application Information

Diseases (Pathogens)	Application Rates		Application Interval/ Timing	Special Instructions
	oz./ 1,000 sq. ft.	oz./A		
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	0.18 to 0.37 (0.005 to 0.012 lb. a.i./A)	8 to 16 (0.25 to 0.5 lb. a.i./A)	14 to 21 days	Use preventively. Begin application when conditions favor disease development.
Brown Patch (<i>Rhizoctonia solani</i>) Anthraxnose (<i>Colletotrichum graminicola</i>)	0.28 to 0.37 (0.009 to 0.012 lb. a.i./A)	12 to 16 (0.375 to 0.5 lb. a.i./A)	14 to 21 days	Begin applications when conditions favor disease development.
Gray Leaf Spot (<i>Pyricularia grisea</i>) Necrotic Ring Spot (<i>Ophiostoma korrae</i>) Red Thread (<i>Laetisaria fuciformis</i>) Rhizoctonia Large Patch (<i>Rhizoctonia solani</i>) Rust Diseases (<i>Puccinia</i> spp.)	0.37 (0.012 lb. a.i./A)	16 (0.5 lb. a.i./A)	14 days	Begin applications when conditions favor disease development or when the disease first appears.

(continued)

Table 1 - TURFGRASS – Specific Diseases, Rates, Timings and Application Information (continued)

Diseases (Pathogens)	Application Rates		Application Interval/ Timing	Special Instructions
	oz./ 1,000 sq. ft.	oz./A		
Summer Patch <i>(Magnaporthe poae)</i> Take-all Patch <i>(Gaeumannomyces graminis)</i> Waitea Patch Brown Ring Patch <i>(Waitea circinata)</i> Zoysia Patch <i>(Rhizoctonia solani)</i>	0.37 (0.012 lb. a.i./A)	16 (0.5 lb. a.i./A)	14 days	Begin applications when conditions favor disease development or when the disease first appears.
Fairy Ring <i>(Various Basidiomycetes)</i>	0.37 (0.012 lb. a.i./A)	16 (0.5 lb. a.i./A)	21 days	Apply as soon as possible after first symptoms appear. Apply in a volume of 4 gals. per 1,000 sq. ft. For optimal control reapply after 21 days. Symptoms may take several weeks to disappear following application.
Pink Snow Mold Fusarium Patch <i>(Microdochium nivale)</i> Gray Snow Mold <i>(Typhula spp.)</i> Yellow Patch <i>(Rhizoctonia cerealis)</i>	0.37 to 0.44 (0.012 to 0.014 lb. a.i./A)	16 to 19.2 (0.5 to 0.6 lb. a.i./A)	Late Fall	Make 1 to 2 applications in combination with PCNB, chlorothalonil or other fungicides active on snow mold in late fall before snow cover and if possible, one application during a mid-winter thaw.

Sojourn 50 T&O Rate Conversion Chart for Turfgrass

oz. product/ 1000 sq. ft.	oz. product/acre	lb. product/acre	lb. a.i./acre
0.18	8	0.5	0.25
0.28	12	0.75	0.38
0.37	16	1.0	0.50
0.44	19.2	1.2	0.60

1 level teaspoon contains 2.4 grams, and one 8 fl. oz. (0.25 lb. a.i./A) cup contains 4.1 oz. (0.128 lb. a.i./A) of **Sojourn 50 T&O**.

ORNAMENTAL PLANTS

Not registered for use by California.

- **Sojourn 50 T&O** provides disease control on ornamental plants in commercial outdoor nurseries and ornamental plants found in landscaped areas around residential, institutional, public, commercial, and industrial buildings, parks, recreational areas, and athletic fields.
- **Sojourn 50 T&O** provides disease control on ornamental plants in indoor greenhouses and nurseries.
- Use **Sojourn 50 T&O** as part of a preventive disease control program.
- **Sojourn 50 T&O** is more effective when used preventively or early in disease development. Length of disease control will vary according to local environmental conditions, disease pressure, and management practices.
- Follow all use directions on this label. Failure to follow label directions may result in injury and or inferior disease control.

ORNAMENTAL PLANT TOLERANCE

IMPORTANT: **Sojourn 50 T&O** may cause growth regulation to some ornamental plant species. The large number of existing ornamental species and their varieties and cultivars coupled with the constant introduction of new varieties makes it impossible to field test **Sojourn 50 T&O** in every locale where it is sold or in all of the combinations of location and plant variety. Differences include the soil or media type, pH, moisture or fertility, environmental conditions such as temperature, lighting or degree-days and horticultural practice and the manner of use and application of this product. To ensure that **Sojourn 50 T&O** is compatible with the ornamental plant variety or cultivar under your specific conditions, test this product on a small test area and observe for phytotoxicity or other unintended effects for at least two weeks before making large-scale applications.

APPLICATION DIRECTIONS

Foliar Applications

- Apply **Sojourn 50 T&O** at 1 to 4 oz. (0.03125 to 0.125. a.i./A) of product per 100 gallons of water for disease control in ornamental plants.
- Apply as a full coverage spray to the point of run-off.
- Apply as needed on a 14-to-28-day interval.
- Under conditions of optimal disease development (such as high humidity and temperature), use higher specified rates and shorter specified application intervals.
- When applications are made curatively, use higher specified rates of **Sojourn 50 T&O** and shorter specified application intervals. Use the higher specified application rate in areas with a history of severe disease pressure.
- For general disease control in landscapes, apply 1 to 4 oz. (0.03125 to 0.125. a.i./A) of product per 100 gallons of water on a 21- to 28-day interval.

Drench Applications

- To control soilborne, seedling and crown diseases, apply **Sojourn 50 T&O** as a preventive drench treatment prior to disease development.
- Apply at 1 to 4 oz. (0.03125 to 0.125. a.i./A) per 100 gallon of water in a volume of 1 to 2 pt. of solution per square foot of surface area on a 14-to-28-day interval.
- Roots must be healthy to ensure systemic uptake, translocation, and disease protection.

COMPATIBILITY

Sojourn 50 T&O is compatible with most commonly used insecticides, fungicides and spray adjuvants used in the production of ornamental plants. When using **Sojourn 50 T&O** in tank mixes with other pesticides, observe all directions for use and precautions on the label of tank mix partner. Before applying a tank mix for the first time, check physical compatibility of the mixture. **DO NOT** tank mix with any product which contains a prohibition on tank mixing.

APPLICATION EQUIPMENT

Apply **Sojourn 50 T&O** using suitable hand or power operated application equipment such as portable, pump-up, backpack, hydraulic or boom sprayers in a manner to provide complete and uniform coverage.

ORNAMENTAL RESTRICTIONS

- **DO NOT** apply more than 4 lbs. (2 lbs. a.i.) of **Sojourn 50 T&O** per acre per year. See Table 2 for specific application rates.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply by air.
- **DO NOT** graze animals on treated areas of one year after application.

- **DO NOT** feed plants from treated areas to livestock or poultry.
- **DO NOT** apply more than 4 oz. (0.125 a.i./A) in a single application.
- **DO NOT** make repeat applications less than 14 days apart.
- Up to 32 applications can be made not exceeding 2 lbs. a.i./A (regardless of Metconazole source).

Not registered for use by California.

Table 2. DIRECTIONS FOR USE ON ORNAMENTALS

Diseases*	Pathogen	Application Rates (By Weight)	Special Instructions
Anthracnose	<i>Apiognomon</i> spp. <i>Colletotrichum</i> spp. <i>Cryptocline</i> spp. <i>Discula</i> spp. <i>Gleosporium</i> spp.	1 to 4 oz. (0.03125 to 0.125 a.i./A)/ 100 gals. on a 14- to 28-day interval	When using a typical high-volume sprayer use 100 gals. of spray mix to treat 20,000 sq. ft. of area. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
Botrytis	<i>Botrytis</i> spp.		
Blossom Rot/Brown Rot	<i>Monilinia</i> spp.		
Boxwood Blight	<i>Calonectria pseudonaviculatum</i> <i>Cylindrocladium buxicola</i> <i>Cylindrocladium pseudonaviculatum</i>	4 oz. (0.125 a.i./A)/ 100 gals. on a 14- to 28-day interval	
Conifer Blight	<i>Diplodia pinea</i> <i>Phomopsis</i> spp. <i>Rhabdocline</i> spp. <i>Rhizosphaera</i> spp. <i>Sphaeropsis</i> spp.	2 to 4 oz. (0.0625 to 0.125 a.i./A)/ 100 gals. on a 14- to 28-day interval	
Damping-Off/Stem and Root Rots	<i>Rhizoctonia solani</i>	1 to 4 oz. (0.03125 to 0.125 a.i./A)/ 100 gals. on a 14- to 28-day interval	
Eastern Filbert Blight	<i>Anisogramma anomala</i>		
Fusarium Root Rot and Damping Off	<i>Fusarium oxysporum</i> <i>Fusarium solani</i>	4 oz. (0.125 a.i./A)/ 100 gals. on a 14- to 28-day interval	

(continued)

Table 2. DIRECTIONS FOR USE ON ORNAMENTALS (continued)

Diseases*	Pathogen	Application Rates (By Weight)	Special Instructions
Leaf Spots	<i>Alternaria</i> spp. <i>Cercospora</i> spp. <i>Diplocarpon</i> spp. <i>Mycosphaerella</i> spp. <i>Myrothecium</i> spp. <i>Phyllosticta</i> spp. <i>Septoria</i> spp.	1 to 4 oz. (0.03125 to 0.125 a.i./A)/ 100 gals. on a 14- to 28-day interval	When using a typical high-volume sprayer use 100 gals. of spray mix to treat 20,000 sq. ft. of area. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
Powdery Mildew	<i>Erysiphe</i> spp. <i>Microsphaera</i> spp. <i>Oidium</i> spp. <i>Phyllactinia</i> spp. <i>Podosphaera</i> spp. <i>Sphaerotheca</i> spp.		
Rusts	<i>Chrysomyxa</i> spp. <i>Coleosporium</i> spp. <i>Gymnosporangium</i> spp. <i>Puccinia</i> spp. <i>Uredo</i> spp. <i>Uromyces</i> spp.		
Scab	<i>Cladosporium</i> spp. <i>Sphaceloma</i> spp. <i>Venturia</i> spp.	2 to 4 oz. . (0.0625 to 0.125 a.i./A)/ 100 gals. on a 14- to 28-day interval	
Shot Hole	<i>Wilsonomyces carpophilus</i> <i>Blumeriella</i> spp.		
White Mold	<i>Sclerotinia sclerotium</i>		

*If symptoms are visible of a labeled disease, use a registered contact, protectant fungicide.

Table 3. ORNAMENTAL PLANT LIST

Note: **Sojourn 50 T&O** has been tested on the following plants with no crop injury.

Flowering and Foliage Plants	Herbaceous Ornamental Plants	Woody Ornamental Plants	Non-Bearing Tree, Fruit, Nut and Vine
Geranium Petunia Poinsettia Snapdragon	Carnation Chrysanthemum Daylily Dianthus Hydrangea Iris Lantana Liriope Marigold Phlox Verbena	Amelanchier Ash Azalea Bayberry Beech Boxwood Birch Camellia Conifers Cotoneaster Crabapple Crape Myrtle Dogwood Elm Euonymus Fir Hawthorn Hemlock Holly Juniper Laurel Lilac Linden London Plane Magnolia Maple Mountain Ash Mountain Laurel Oaks Pine Poplar Photinia Pyracantha	Apple Apricot Cherry Citrus Filbert Grape Peach Pear Pecan Pistachio Plum Walnut

(continued)

Table 3. ORNAMENTAL PLANT LIST (continued)

Flowering and Foliage Plants	Herbaceous Ornamental Plants	Woody Ornamental Plants	Non-Bearing Tree, Fruit, Nut and Vine
		Rhododendron Rose Sassafras Spirea Spruce Sweetgum Sycamore	

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: **DO NOT** store at extreme temperatures; store in a cool dry place. Store in a dry place away from heat or open flame. Keep pesticide in original container. Keep container closed when not in use. **DO NOT** put dilute into food or drink containers. **DO NOT** store in or around the home.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Non-Refillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for plastic containers, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures approved by State and local authorities. For metal containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

(continued)

STORAGE AND DISPOSAL *(continued)*

CONTAINER HANDLING: *(continued)*

Non-Refillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for plastic containers, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures approved by State and local authorities. For metal containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Refillable Fiber Drums with Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with this fungicide only. **DO NOT** reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: **DO NOT** reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by State and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill or by incineration, or by other procedures approved by State and local authorities.

(continued)

STORAGE AND DISPOSAL *(continued)*

CONTAINER HANDLING: *(continued)*

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with this pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. Check for leaks after refilling and before transporting. Disposing of Container: **DO NOT** reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom, and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration, and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for plastic containers, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures approved by State and local authorities. For metal containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

PRIMEROS AUXILIOS	
SI SE INGIERE:	• Llame de inmediato a un centro de control de envenenamientos o a un médico para consejo de tratamiento. • Si la persona puede tragar, haga que tome pequeños sorbos de agua. • NO induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. • NO administre nada por boca a una persona que haya perdido el conocimiento.
SI ENTRA EN CONTACTO CON LOS OJOS:	• Mantenga el ojo abierto y enjuáguelo lenta y cuidadosamente con agua, durante 15 a 20 minutos. • Si utiliza lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
SI CAE EN LA PIEL O LA ROPA:	• Quítese la ropa contaminada. • Enjuague la piel inmediatamente con bastante agua por 15 - 20 minutos. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
SI ES INHALADO:	• Traslade a la persona al aire fresco. • Si la persona no está respirando llame al 911 o a una ambulancia, luego dé respiración artificial, preferiblemente de boca a boca, si es posible. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
NÚMERO DE TELÉFONO DIRECTO	
Cuando llame a un centro de control de envenenamientos, o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. Para información de emergencia sobre este producto, llame al centro de control de envenenamientos a 1-800-222-1222.	

DECLARACIONES PREVENTIVAS

RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS

PRECAUCIÓN

Daño si se ingiere. Causa irritación moderada en los ojos. Evite el contacto con los ojos, la piel o la ropa. Lávese bien con agua y jabón después de manipular y antes de comer, beber, masticar chicle, usar tabaco o usar el baño. Quítese y lave la ropa contaminada antes de reutilizarla.

EQUIPO DE PROTECCIÓN PERSONAL (PPE, POR SUS SIGLAS EN INGLÉS)

Los aplicadores y otros manipuladores deberán usar:

- Overol (mameluco), camisa de manga larga y pantalones largos
- Guantes resistentes a productos químicos fabricados con laminado de barrera, goma de butilo ≥ 14 mils, goma de nitrilo ≥ 14 mils, goma de neopreno ≥ 14 mils, goma natural ≥ 14 mils, polietileno, cloruro de polivinilo (PVC) ≥ 14 mils, o Viton ≥ 14 mils
- Zapatos y calcetines

RECOMENDACIONES DE SEGURIDAD DEL USUARIO

Los usuarios deben:

- Quitarse inmediatamente la ropa/equipo de protección personal (PPE) si el pesticida entra en contacto con la ropa o el PPE por dentro. Luego deben lavarse muy bien y ponerse ropa limpia.
- Qútese equipo de protección personal (PPE) inmediatamente después de manipular este producto. Lávese bien y póngase ropa limpia lo antes posible.

RIESGOS AMBIENTALES

Este pesticida es tóxico para las aves, los mamíferos, los peces y los invertebrados acuáticos. **NO** aplique directamente al agua, en áreas donde haya aguas superficiales, o en áreas intermareales que se encuentren bajo el nivel de altura máxima promedio del agua. **NO** contamine el agua cuando disponga de las aguas de lavado del equipo o aguas de enjuague. La deriva y la escorrentía desde las áreas tratadas pueden ser peligrosos para los organismos acuáticos en las aguas adyacentes a dichas zonas.

Advertencia Sobre Aguas Subterráneas

Este producto químico tiene propiedades y características relacionadas con productos químicos detectados en aguas subterráneas. El uso de este producto químico en zonas con suelos permeables, especialmente donde el nivel freático es poco profundo, puede provocar la contaminación de las aguas subterráneas.

Aviso de deriva de pulverización

Este producto puede afectar la calidad del agua superficial debido a la dispersión de las pulverizaciones y la escorrentía del agua de lluvia. Este producto tiene un alto potencial de escorrentía durante varios meses o más después de la aplicación. Los suelos con drenaje deficiente y los suelos con niveles freáticos poco profundos son más propensos a generar escorrentía que contiene este producto. Una franja vegetativa de protección, nivelada y bien mantenida entre las áreas donde este producto es utilizado y cuerpos de agua superficiales tales como estanques, arroyos y manantiales, reducirá el potencial de contaminación del agua por escorrentía de lluvia. La escorrentía de este producto puede reducirse al evitar aplicaciones cuando se pronostique lluvia dentro de 48 horas. Las buenas prácticas de control de la erosión reducirán la contribución de este producto a la contaminación de las aguas superficiales.

NO descargue los efluentes que contengan este producto en lagos, arroyos, estanques, estuarios, océanos u otras aguas a menos que sea de acuerdo con los requisitos de un permiso del Sistema Nacional de Eliminación de Descargas de Contaminantes (NPDES, por sus siglas en inglés) y la autoridad otorgante haya sido notificada por escrito antes de la descarga. **NO** descargue efluentes que contengan este producto a los sistemas de alcantarillado sin notificar previamente a la autoridad local de la planta de tratamiento de aguas residuales. Para obtener orientación, comuníquese con la Junta Estatal de Agua o la Oficina Regional de la EPA.

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal.

ALMACENAMIENTO Y DESECHO

NO contamine el agua, los alimentos ni los forrajes durante el almacenamiento y el desecho.

ALMACENAMIENTO DE PESTICIDAS: **NO** almacene a temperaturas extremas; almacene en un lugar fresco y seco. Almacene en un lugar seco, lejos del calor o de llamas abiertas. Mantenga el pesticida en su envase original. Mantenga el envase cerrado cuando no lo use. **NO** vierta el producto diluido en envases de alimentos o bebidas. **NO** lo almacene dentro ni en los alrededores de la vivienda.

DESECHO DE PESTICIDAS: Los residuos resultantes del uso de este producto pueden eliminarse en el lugar o en una instalación de eliminación de residuos autorizada. Si estos residuos no pueden eliminarse de acuerdo con las instrucciones de la etiqueta, contacte la agencia estatal de pesticidas o de control ambiental de su estado, o el representante para el Manejo de Residuos Peligrosos de la Oficina Regional de la EPA más cercana para obtener instrucciones sobre métodos adecuados de eliminación.

(continuado)

ALMACENAMIENTO Y DESECHO *(continuado)*

MANIPULACIÓN DE ENVASES:

Envases de plástico y metal no rellenables (Capacidad igual o menos de 50 libras): Envase no rellenable. **NO** reutilice ni rellene este envase. Enjuague tres veces el envase (o equivalente) inmediatamente después de vaciarlo. Enjuague tres veces siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezcla. Llene el envase hasta 1/4 de su capacidad con agua y tape de nuevo. Agite por 10 segundos. Vierta las aguas de enjuague en el equipo de aplicación o tanque de mezcla, o almacene aguas de enjuague para uso posterior o eliminación. Drene por 10 segundos después de que el flujo empiece a gotear. Repita este procedimiento dos veces más. Luego, para envases de plástico, ofrézcalo para reciclaje si está disponible o perforo y desecho en un relleno sanitario, o mediante incineración, o por otros procedimientos aprobados por las autoridades estatales y locales. Para envases de metal, ofrézcalo para reciclaje si está disponible o reacondiciónelo si es apropiado, o perforo y desecho en un relleno sanitario o por otros procedimientos aprobados por las autoridades estatales y locales.

Envases de plástico y metal no rellenables (Capacidad más de 50 libras): Envase no rellenable. **NO** reutilice ni rellene este envase. Enjuague tres veces el envase (o equivalente) inmediatamente después de vaciarlo. Enjuague tres veces siguiendo estas indicaciones: vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezcla. Llene el envase hasta 1/4 de su capacidad con agua. Reemplace y asegure los cierres. Vuelque el envase al lado y hágalo rodar por 30 segundos, asegurando por lo menos una revolución completa. Ponga el envase vertical e inclínelo hacia adelante y hacia atrás varias veces. Dé vuelta al envase hacia el lado opuesto e inclínelo hacia adelante y hacia atrás varias veces. Vierta las aguas de enjuague en el equipo de aplicación o tanque de mezcla, o almacene aguas de enjuague para uso posterior o eliminación. Repita este procedimiento dos veces más. Luego, para envases de plástico, ofrézcalo para reciclaje si está disponible o perforo y desecho en un relleno sanitario, o mediante incineración, o por otros procedimientos aprobados por las autoridades estatales y locales. Para envases de metal, ofrézcalo para reciclaje si está disponible o reacondiciónelo si es apropiado, o perforo y desecho en un relleno sanitario o por otros procedimientos aprobados por las autoridades estatales y locales.

(continuado)

ALMACENAMIENTO Y DESECHO *(continuado)*

MANIPULACIÓN DE ENVASES: *(continuado)*

Tambores de fibra con bolsa rellenables: Envase rellenable (solo bidón de fibra). Rellenado del tambor de fibra: Rellene este tambor de fibra únicamente con este fungicida. **NO** reutilice este tambor de fibra para ningún otro propósito. La limpieza antes de rellenarlo es responsabilidad de la persona que lo rellena. Vacíe por completo la bolsa interior, sacudiéndola y golpeando sus lados y fondo para aflojar las partículas adheridas. Vacíe los residuos en el equipo de aplicación o de fabricación. Desecho del tambor de fibra y/o bolsa interior: **NO** reutilice este tambor de fibra para ningún otro propósito que no sea el de rellenar (véase más arriba). La limpieza del envase (bolsa interior y/o tambor de fibra) antes de su desecho final es responsabilidad de la persona que lo desecha. Ofrezca la bolsa interior para reciclaje, si el servicio está disponible, o deshágase de la misma en un relleno sanitario o mediante incineración. **NO** lo queme a menos que esté permitido por las ordenanzas estatales y locales. Si el tambor está contaminado y no se puede reutilizar, deséchelo de la misma manera requerida para sus bolsas interiores. Para limpiar el tambor de fibra antes de su desecho final, vacíe por completo el tambor de fibra, sacudiendo y golpeando los lados y el fondo para aflojar las partículas pegadas. Vacíe los residuos en el equipo de aplicación o industrial. Después, ofrezca el tambor de fibra para reciclaje si el servicio está disponible o deséchelo en un vertedero o mediante incineración o mediante otros procedimientos aprobados por las autoridades estatales y locales.

(continuado)

ALMACENAMIENTO Y DESECHO *(continuado)*

MANIPULACIÓN DE ENVASES: *(continuado)*

Todos los demás envases rellenables: Envase rellenable. Rellene este envase solo con pesticida. **NO** reutilice este envase para ningún otro propósito. La limpieza antes de relleno es responsabilidad de la persona que lo rellena. Antes de rellenar, inspeccione cuidadosamente en busca de daños tales como grietas, perforaciones, abrasiones, roscas desgastadas y dispositivos de cierre. Compruebe que no haya fugas después del rellenado y antes de transportar. Desecho del envase: **NO** reutilice este envase para ningún otro propósito que no sea su relleno (vea la sección anterior). La limpieza del envase antes de su eliminación final es la responsabilidad de la persona que se deshace del envase. Para limpiar el envase antes de su desecho final, utilice el siguiente procedimiento de enjuague a presión. Inserte una lanza con boquilla apropiada para la limpieza de tanques en el envase, y asegúrese de que el agua asperjada cubra completamente la parte superior, parte inferior y los lados del interior del envase. El fabricante de las boquillas generalmente proporciona instrucciones acerca de la presión de aspersión adecuada, la duración de la aspersión y/o volumen de aspersión. Si las instrucciones del fabricante no están disponibles, enjuague el envase a presión por lo menos durante 60 segundos usando una presión mínima de 30 PSI (libras por pulgada cuadrada), con un volumen de enjuague mínimo del 10% del volumen del envase. Drene, vierta o bombee las aguas de enjuague hacia dentro del equipo de aplicación o en un sistema recolector de aguas de enjuague. Repita este procedimiento de enjuague a presión dos veces más. Luego, para envases de plástico, ofrézcalo para reciclaje si está disponible o perforo y deseche en un relleno sanitario, o mediante incineración, o por otros procedimientos aprobados por las autoridades estatales y locales. Para envases de metal, ofrézcalo para reciclaje si está disponible o reacondiciónelo si es apropiado, o perforo y deseche en un relleno sanitario o por otros procedimientos aprobados por las autoridades estatales y locales.

¡EL ENVASE NO ES SEGURO PARA ALIMENTOS, FORRAJES, NI AGUA POTABLE!

METCONAZOLE GROUP 3 FUNGICIDE

OPEN HERE

Sojourn 50 T&O

Turfgrass and Ornamental Fungicide
A SYSTEMIC FUNGICIDE FOR CONTROL OF THE LISTED TURFGRASS AND ORNAMENTAL DISEASES

ACTIVE INGREDIENT:	WT. BY %
Metconazole: 5-[(4-chlorophenyl)methyl]-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol	50.0%
OTHER INGREDIENTS:	50.0%
TOTAL:	100.0%

Sojourn 50 T&O is a water dispersible granule containing 50% active ingredient.

KEEP OUT OF REACH OF CHILDREN /
MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand this label, find someone to explain it to you in detail.)

**See label booklet for complete First Aid, Precautionary Statements,
Directions For Use, and Storage and Disposal.**

EPA Reg. No. 82633-122-83529

EPA Est. No. (AG) 72159-GA-001; (MC) 89332-GA-001; (SC) 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

Net Contents: **5 lbs.**