

Companera 21.50% WDG

HERBICIDE

For postemergent control of annual grass and broadleaf weeds
in spring and winter wheat (including durum), and triticale.

ACTIVE INGREDIENT:	WT. BY %
pyroxsulam: N-(5,7-dimethoxy[1,2,4]triazolo [1,5-a]pyrimidin-2-yl)-2-methoxy-4-(trifluoromethyl)-3-pyridinesulfonamide	21.5%
OTHER INGREDIENTS:	78.5%
TOTAL:	100.0%
Contains 0.215 lb. of active ingredient per pound of product.	

KEEP OUT OF REACH OF CHILDREN / MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give anything to an unconscious person.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378 , Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	

PRIMEROS AUXILIOS	
SI SE INGIERE:	• Llame de inmediato a un centro de control de envenenamientos o a un médico para consejo de tratamiento. • Si la persona puede tragar, haga que tome pequeños sorbos de agua. • NO induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. • NO administre nada por boca a una persona que haya perdido el conocimiento.
SI ENTRA EN CONTACTO CON LOS OJOS:	• Mantenga el ojo abierto y enjuáguelo lenta y cuidadosamente con agua, durante 15 a 20 minutos. • Si utiliza lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
NÚMERO DE TELÉFONO DIRECTO	
Cuando llame a un centro de control de envenenamientos, o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. Para información de emergencia sobre este producto, llame al centro de control de envenenamientos a 1-800-222-1222 . Para información general sobre este producto, contacte al Centro Nacional de Información de Pesticidas (NPIC, por sus siglas en inglés) a 1-800-858-7378 , Lunes a Viernes, 8 AM a 12 PM PST (hora del Pacífico), o a http://npic.orst.edu .	

See label booklet for complete Precautionary Statements,
Directions For Use and Storage and Disposal.

Manufactured For:
Sharda USA LLC



7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 82633-141-83529

EPA Est. No. **AG** 72159-GA-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Weight: **10 lbs.**

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION/PRECAUCIÓN

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear protective eyewear, long sleeved shirt and long pants, socks and shoes.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

- Users should:**
- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
 - Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

GROUNDWATER ADVISORY: This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

SURFACE WATER ADVISORY: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this chemical from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

PRODUCT INFORMATION

Use **Companera 21.50% WDG** as a postemergence herbicide for the control of annual grass and broadleaf weeds in spring and winter wheat (including durum), and triticale. **Companera 21.50% WDG** rapidly stops growth of susceptible weeds. However, typical symptoms (discoloration) of controlled or suppressed weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent upon weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

When applying this product in tank mix combination, follow all applicable use directions, precautions, and limitations on each manufacturer's label.

Use Restrictions

- **Chemigation:** **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply **Companera 21.50% WDG** directly to, or otherwise permit it to come into direct contact with, susceptible crops or desirable plants including alfalfa, barley, canola, beans, cotton, flowers, grapes, lettuce, lentils, mustard, oats, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes, vegetables, or other desirable broadleaf crops or ornamental plants. **DO NOT** permit spray mists containing **Companera 21.50% WDG** to drift onto such plants.
- **DO NOT** apply to crops underseeded with legumes.

MANDATORY SPRAY DRIFT MANAGEMENT

- Aerial Applications:**
- **DO NOT** release spray at a height greater than 10 ft. above the ground or vegetative canopy unless a greater application height is necessary for pilot safety.
 - Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
 - If the windspeed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 - 15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
 - **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
 - **DO NOT** apply during temperature inversions.
- Ground Boom Applications:**
- User must only apply with the release height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
 - Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
 - **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
 - **DO NOT** apply during temperature inversions.

SPRAY DRIFT MANAGEMENT ADVISORY

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume:** Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure:** Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle:** Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- **Adjust Nozzles:** Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

Avoiding Injurious Spray Drift

This product can affect broadleaf plants directly through foliage and indirectly by root uptake from treated soil. **DO NOT** apply **Companera 21.50% WDG** directly to, or allow spray drift to come into contact with, broadleaf crops including alfalfa, barley, canola, beans, cotton, flowers, grapes, lettuce, lentils, mustard, oats, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes, vegetables, or other desirable broadleaf crops or ornamental plants or soil where sensitive crops will be planted the same season. (See Crop Rotation Intervals section.)

Make applications only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure crops, whether dormant or actively growing. When applying **Companera 21.50% WDG**, use low pressure equipment capable of producing sprays of uniform droplet size with a minimum of fine spray droplets. Under adverse weather conditions, fine spray droplets that **DO NOT** settle rapidly onto target vegetation may be carried a considerable distance from the treatment area. A drift control or spray thickening agent may be used with this product to improve spray deposition and minimize the potential for spray drift. If used, follow all use directions and precautions on the product label.

Ground Applications: To minimize spray drift, apply **Companera 21.50% WDG** in a total spray volume of 10 gallons or more per acre using spray equipment designed to produce large droplet, low pressure sprays. Refer to the spray equipment manufacturer’s directions for detailed information on nozzle types, arrangement, spacing and operating height and pressure. Apply spot treatments only with a calibrated boom to prevent over application. Operate equipment at spray pressures no greater than is necessary to produce a uniform spray pattern. Operate the spray boom no higher than is necessary to produce a uniformly overlapping pattern between spray nozzles. **DO NOT** apply with hollow cone-type insecticide nozzles or other nozzles that produce a fine-droplet spray.

Aerial Application: To minimize spray drift, apply **Companera 21.50% WDG** in a total spray volume of 5 gallons or more per acre. Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Avoid applications below 2 mph due to variable wind direction and high potential for temperature inversion. Minimize spray drift from aerial applications by applying a coarse spray at spray boom pressure no greater than 30 psi; by using straight-stream nozzles directed straight back; and by using a spray boom no longer than 3/4 of the rotor or wingspan of the aircraft. Evaluate spray pattern and droplet size distribution by applying sprays containing a water-soluble dye marker or appropriate drift control agents over a paper tape (adding machine tape). Mechanical flagging devices may also be used.

Sensitive Areas: Apply the pesticide only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

California Application Requirements for Protection of Sensitive Crops

In addition to precautions in the **Aerial Drift Reduction Advisory** section, the following drift management requirements must be followed to minimize the potential for exposure of sensitive crops in California.

Determine the prevailing wind speed and direction before application. Choose nozzles that will give a coarse droplet size spectrum (volume median diameter (VMD) of 350 - 400 microns) and minimize droplets that are less than 200 microns.

Buffer Zones

The following buffer zones between the treated area and sensitive crops are required when these sensitive crops are downwind of the application site.

Sensitive Crop	Downwind Buffer Zone – Feet	
	Ground	Air
Carrot	120	500
Onions	10	100
Oat and ryegrass	50	250
All other broadleaf annual crops	20	160
Tree and vine crops	20	160
Dormant tree and vine crops	No buffer zone is required	

CROP ROTATION INTERVALS

The following rotational crops may be planted at the indicated interval following application of **Companera 21.50% WDG**.

Crop Rotation Intervals for All States Except Arizona, California, Idaho, Oregon, and Washington Superscripted numbers refer to Crop Specific Rotation Information.	
Crop	Rotation Interval (Months) ¹
Wheat, triticale	1
Soybean ²	5
Barley, field corn, grasses, millet, oats, popcorn, seed corn, sweet corn, grain sorghum ⁴ , sunflower ⁴	9
Alfalfa, camelina, canola, chickpea, cotton ³ , dry bean, pea (dry and succulent), flax, lentil, mustard, peanuts, potato, safflower, sugar beet, sunflower	9
Other crops not listed	12

Crop Specific Rotation Information

¹ Minimum number of months that must elapse before planting other crops after application of **Companera 21.50% WDG**.
² As a rotation crop, soybeans may be planted 3 months following an application of **Companera 21.50% WDG** in February or later in the following states: Alabama, Arkansas, Delaware, Georgia, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maryland, Missouri, Mississippi, North Carolina, Nebraska, New Jersey, Ohio, Oklahoma, Pennsylvania, South Carolina, Tennessee, Texas, and Virginia. However, to ensure adequate crop safety, avoid planting soybeans prior to April 30 following an application of **Companera 21.50% WDG** made before February. All other states not listed require a minimum rotation interval of 5 months after an application of **Companera 21.50% WDG**.
³ As a rotation crop, cotton may be planted 3 months following an application of **Companera 21.50% WDG** in February or later in the following states: Alabama, Arkansas, Georgia, Kansas, Kentucky, Louisiana, Missouri, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, and Virginia. However, to ensure adequate crop safety, avoid planting cotton prior to April 30 following an application of **Companera 21.50% WDG** made before February. All other states not listed require a minimum rotation interval of 9 months after an application of **Companera 21.50% WDG**.
⁴ As a rotation crop, grain sorghum and sunflowers may be planted 3 months following an application of **Companera 21.50% WDG** in February or later in the following states: Alabama, Arkansas, Colorado, Delaware, Georgia, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maryland, Missouri, Mississippi, North Carolina, Nebraska, New Jersey, Ohio, Oklahoma, Pennsylvania, South Carolina, Tennessee, Texas, and Virginia. However, to ensure adequate crop safety, avoid planting sunflowers and grain sorghum prior to April 30 following an application of **Companera 21.50% WDG** made before February. All other states not listed require a minimum rotation interval of 9 months after an application of **Companera 21.50% WDG**.

Crop Rotation Intervals for Arizona and California Superscripted numbers refer to Crop Specific Rotation Information.	
Crop	Rotation Interval (Months) ¹
Triticale, wheat	1
Corn (field, silage), cotton, dry beans, melons, sorghum (grain, silage), sudangrass	3
Alfalfa, broccoli, cabbage, cauliflower, lettuce, onion	5
Barley, grasses, millet, oats, popcorn, seed corn, sweet corn, camelina, canola, chickpea, flax, lentil, mustard, pea (dry and succulent), potato, safflower, soybean, sugar beet, sunflower, tomato	9
Other crops not listed	12
Crop Specific Rotation Information	
¹ Minimum number of months that must elapse before planting other crops after application of Companera 21.50% WDG .	

Crop Rotation Intervals for Idaho, Oregon, and Washington Superscripted numbers refer to Crop Specific Rotation Information.		
Crop	Soil pH >6 and Rainfall ² >16 Inches	Soil pH <6 or Rainfall ² <16 Inches
Wheat, triticale	1	1
Barley, field corn, grasses, millet, oats, popcorn, seed corn, sweet corn, grain sorghum	10	10
Alfalfa, camelina, canola, cotton, dry bean, flax, mustard, peanuts, safflower, soybean, sugar beet, sunflower	10	10
Pulse crops ³ including chickpea, lentil, and pea (dry and succulent), potato ³	10	18
Other crops not listed	12	18
Crop Specific Rotation Information		
¹ Minimum number of months that must elapse before planting other crops after application of Companera 21.50% WDG .		
² Including irrigation.		
³ Pulse crops, including chickpea, lentil, and pea (dry and succulent), and potatoes may be planted 10 months after application if the soil pH is uniformly 6 or greater AND total rainfall (including irrigation) during the interval is greater than 16 inches. If the soil pH is less than 6 OR total rainfall (including irrigation) is less than 16 inches, then the rotation interval is 18 months.		

Note: Companera 21.50% WDG is degraded primarily by microbial activity and breaks down more rapidly under favorable soil moisture and temperature conditions. Correspondingly, the rate of degradation may be slower under extreme conditions of drought or cold temperatures. When soil moisture conditions are abnormally dry during the interval between an application of **Companera 21.50% WDG** and planting the next crop, conduct a field bioassay by planting test strips of the desired rotational crop. Monitor the test strips during germination and emergence for any abnormal growth to determine if the rotational crop can be grown successfully.

Mixing Directions

Companera 21.50% WDG – Alone

1. Fill the tank with 1/2 of the total amount of water and begin agitation (If using a liquid nitrogen fertilizer solution in place of water, see **Application Directions** section for additional details).
2. Add a water conditioning agent, if needed.
3. Add the required amount of **Companera 21.50% WDG**.
4. Add the required amount of adjuvant (refer to **Adjuvants** section).
5. Continue agitation while filling the spray tank to the required volume.
6. To ensure a uniform spray mixture, continuous agitation is required during application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

Companera 21.50% WDG – Tank Mix

If a broader spectrum of weed control is needed, **Companera 21.50% WDG** may be tank mixed with labeled rates of other herbicides provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product.

Tank Mixing:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- **DO NOT** mix with products containing dicamba or amine formulations of 2,4-D or MCPA as these products may reduce grass control provided by **Companera 21.50% WDG**.
- **DO NOT** tank mix with organophosphate insecticides as these mixtures may result in unacceptable crop injury.
- **DO NOT** exceed specified application rates for respective products or maximum allowable application rates for any active ingredient in the tank mix.
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.
- Incompatible mixtures may lead to difficult-to-clean herbicide residues in the spray tank, which if not properly cleaned, could cause damage to sensitive crops.

Tank Mix Compatibility Testing: Perform a jar test prior to tank mixing to ensure compatibility of **Companera 21.50% WDG** and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Vigorous, continuous agitation during mixing, filling and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

Mixing Order for Tank Mixes:

1. Fill the spray tank to 1/2 to 3/4 of the total spray volume required with water and begin agitation (If using a liquid nitrogen fertilizer solution in place of water, see **Application Directions** section for additional details).
2. Add a water conditioning agent, if needed.
3. Add **Companera 21.50% WDG** and agitate for 2 to 3 minutes.
4. After adding **Companera 21.50% WDG**, add different formulation types in the following order: (1) dry flowables; (2) wettable powders; (3) aqueous suspensions, flowables and liquids. Maintain agitation and add: (4) emulsifiable concentrates; (5) solutions; and (6) adjuvants. Allow time for complete mixing and dispersion after each addition.
5. Finish filling the spray tank. Maintain continuous agitation during mixing and throughout application. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying. Apply mixture immediately after it is prepared.

If application or agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

Clean-Out Procedures for Spray Equipment

1. Completely drain the spray system, including pump, lines, and spray boom.
2. Fill the spray tank with clean water to at least 10% of the total tank volume and circulate the solution through the entire system so that all internal surfaces are contacted for at least 15 minutes to complete the first rinse of the application equipment. Spray the solution out of the spray tank through the boom.
3. Completely drain the spray system, including lines and spray boom; remove and clean filters and strainers.
4. During the second rinse, fill the container half full with clean water and then add a commercial tank cleaner at the manufacturer’s recommended rates. Circulate the cleaning solution through the entire system for at least 20 minutes. Let the solution stand for several hours. Again, circulate and flush the solution through the lines and boom.
5. Completely drain and flush the spray system, including lines and spray boom.
6. Fill the container with clean water to at least 10% of the total tank volume and circulate the solution through the entire system so that all internal surfaces are contacted for at least 15 minutes to complete the third rinse of the application equipment. Spray the solution out of the spray tank through the boom.

Note: Rinsate may be disposed of on site according to label use directions or at an approved waste disposal facility.

Weeds Controlled (C) or Suppressed (S)

Best results are obtained when grass weeds are treated at the 2-leaf to 2-tiller stage of growth and before broadleaf weeds are larger than 2 inches tall or 2 inches in diameter. Best control is achieved when applications are made to actively growing weeds. Control may be reduced when weeds are exposed to drought or extreme temperatures. **Companera 21.50% WDG** will not control known ALS (Group 2) resistant biotypes of labeled weeds.

GRASS WEEDS		
Common Name	Scientific Name	Weeds Controlled (C) or Suppressed (S)
barley, foxtail	<i>Hordeum jubatum</i>	S
Barnyardgrass*	<i>Echinochloa crus-galli</i>	C
blackgrass	<i>Alopecurus myosuroides</i>	C
bluegrass, bulbous*	<i>Poa bulbosa</i>	C
brome, downy	<i>Bromus tectorum</i>	S

(continued)

GRASS WEEDS (continued)		
Common Name	Scientific Name	Weeds Controlled (C) or Suppressed (S)
brome, Japanese	<i>Bromus japonicus</i>	C
brome, ripgut	<i>Bromus diandrus</i>	C
canarygrass, hood	<i>Phalaris paradoxa</i>	S
canarygrass, littleseed	<i>Phalaris minor</i>	S
cheat	<i>Bromus secalinus</i>	C
chess, hairy	<i>Bromus commutatus</i>	C
corn, volunteer*	<i>Zea mays</i>	C
darnel, Persian*	<i>Lolium persicum</i>	C ⁴
fescue, rattail	<i>Vulpia myuros</i>	S
foxtail, green	<i>Setaria viridis</i>	S
foxtail, yellow	<i>Setaria pumila</i>	C ⁴
oat, wild	<i>Avena fatua</i>	C
quackgrass	<i>Elymus repens</i>	S
rescuegrass	<i>Bromus catharticus</i>	S
ryegrass, Italian	<i>Lolium perenne</i>	C
windgrass	<i>Apera spica-venti</i>	C
BROADLEAF WEEDS		
Common Name	Scientific Name	Weeds Controlled (C) or Suppressed (S)
bedstraw, catchweed (cleavers)	<i>Galium aparine</i>	C
bittercress, hairy*	<i>Cardamine hirsuta</i>	C
buckwheat, wild	<i>Polygonum convolvulus</i>	S
burclover, spotted*	<i>Medicago arabica</i>	C
buttercup, smallflower*	<i>Ranunculus abortivus</i>	C
canola, volunteer (wild turnip) ²	<i>Brassica rapa, B. napus</i>	C
chickweed, common	<i>Stellaria media</i>	C
chickweed, mouseear	<i>Cerastium fontanum</i>	C
clover, white*	<i>Trifolium repens</i>	C
coreopsis, plains*	<i>Coreopsis tinctoria</i> Nutt.	S
evening-primrose, cutleaf*	<i>Oenothera laciniata</i> Hill	S
falseflax, smallseed ¹	<i>Camelina microcarpa</i>	C
fiddleneck, coast	<i>Amsinckia menziesii</i> var. <i>intermedia</i>	C
flixweed ¹	<i>Descurainia sophia</i>	C
geranium, Carolina*	<i>Geranium carolinianum</i>	C
gromwell, corn	<i>Buglossoides arvensis</i>	C
hempnettle, common	<i>Galeopsis tetrahit</i>	C
henbit	<i>Lamium amplexicaule</i>	S
lambsquartars, common	<i>Chenopodium album</i>	C ³
mustard, black	<i>Brassica nigra</i>	C
mustard, blue ¹	<i>Chorispora tenella</i>	C
mustard, tumble ¹	<i>Sisymbrium altissimum</i>	C
mustard, wild	<i>Sinapis arvensis</i>	C
mustard, wormseed ¹	<i>Erysimum cheiranthoides</i>	C
pennycress, field ¹	<i>Thlaspi arvense</i>	C
pepperweed, Virginia*	<i>Lepidium virginicum</i>	C
pigweed, redroot	<i>Amaranthus retroflexus</i>	C
shepherd's-purse ¹	<i>Capsella bursa-pastoris</i>	C
smartweed, annual*	<i>Polygonum sp.</i>	C
tansymustard, pinnate ¹	<i>Descurainia pinnata</i>	C
thistle, Russian	<i>Salsola tragus</i>	C ³
vetch, hairy*	<i>Vicia villosa</i> Roth	C
wallflower, bushy ¹	<i>Erysimum repandum</i>	C

¹ Control may be reduced when application is made after bolting.
² Including herbicide-tolerant canola varieties except Clearfield (imidazolinone-tolerant) canola.
³ Less than 2 inches tall. For control of lambsquartars over 2 inches tall, tank mix with 0.25 lb. a.e. 2,4-D ester or MCPA ester. For control of Russian thistle over 2 inches tall, tank mix with 0.25 lb. a.e. 2,4-D ester.
⁴ One to four-leaf stage of growth.
 * Not registered for use in California.

PYROXSULAM

GROUP 2

HERBICIDE

HERBICIDE RESISTANCE MANAGEMENT

For resistance management, **Companera 21.50% WDG** is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to **Companera 21.50% WDG** and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, consider the below best practices for resistance management:

- Rotate the use of **Companera 21.50% WDG** or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Fields should be scouted prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective. Scout after herbicide application to verify that the treatment was effective and monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact a Sharda USA, LLC representative at <https://www.shardausa.com/>.

APPLICATION DIRECTIONS

Application Timing

Apply **Companera 21.50% WDG** postemergence to the main flush of actively growing weeds according to the target weed stage shown in the above table. Extreme growing conditions such as drought, temperatures near or below freezing prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.

Warm, moist growing conditions promote active weed growth and enhance the activity of **Companera 21.50% WDG** by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds.

If foliage is wet at the time of application, control may be decreased. Applications of **Companera 21.50% WDG** are rainfast within 4 hours after application.

Spray Coverage

Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. **DO NOT** broadcast apply in less than 5 gallons of total spray volume per acre. For best results and to minimize spray drift, apply in a spray volume of 10 gallons or more per acre. As vegetative canopy and weed density increase, increase spray volume to obtain equivalent weed control. Use only nozzle types and spray equipment designed for herbicide application. To reduce spray drift, follow precautions under **Avoiding Injurious Spray Drift**.

Surfactants and Adjuvants

When **Companera 21.50% WDG** is applied alone, use one of the following surfactants or adjuvants:

- Non-ionic surfactant with at least 80% active ingredient at 0.25% to 0.50% v/v (1 to 2 quarts per 100 gallons of spray solution); for best results under dry or low humidity environments, use a rate of 0.50% v/v. Addition of spray quality urea ammonium nitrogen fertilizer (28-0-0 to 32-0-0 at 1 to 2 quarts per acre) or ammonium sulfate fertilizer (21-0-0-24 at 1.5 to 3 lbs. per acre) may be added to non-ionic surfactant to enhance control.
- Crop oil concentrate adjuvant at 1.0 to 1.25% v/v (1 to 1.25 gallons per 100 gallons of spray solution).
- Methylated seed oil adjuvant at 1.0% v/v (1 gallon per 100 gallons of spray solution).

Potential for crop response is increased with the use of oil adjuvants versus non-ionic surfactants. **DO NOT** use oil adjuvants with spray solutions containing nitrogen fertilizer.

When **Companera 21.50% WDG** is applied in combination with emulsifiable concentrate (EC) formulations, such as 2,4-D ester or MCPA ester products, a non-ionic surfactant may be added to the mixture at 1/2 to 1 quart per 100 gallons of spray solution (0.125 to 0.25% v/v). Use the lower amount of surfactant if the total amount of EC product rate/acre exceeds 6 fluid ounces/acre.

DO NOT use additives that lower the spray solution below a pH of 6.0.

Sharda USA, LLC recommends the use of a Chemical Producers and Distributors Association certified surfactant or adjuvant when used with **Companera 21.50% WDG**.

Application in Fluid Fertilizer

Companera 21.50% WDG may be applied in spray solutions containing liquid nitrogen fertilizer. The spray solution should not be composed of more than 50% liquid nitrogen fertilizer and should not exceed 30 lbs. of actual nitrogen per acre. When **Companera 21.50% WDG** is applied in spray solutions containing liquid nitrogen fertilizer, use a non-ionic surfactant at a maximum of 0.25% v/v instead of crop oil concentrate or methylated seed oil.

Temporary crop injury may result when liquid nitrogen fertilizer is used as the spray carrier. Foliar applied liquid nitrogen fertilizer may cause foliar leaf burn, yellowing or reduced growth due to the activity of the liquid fertilizer on the crop.

Spring and Winter Wheat (including Durum), and Triticale

Apply 1 oz. of **Companera 21.50% WDG** per acre (0.013 lb./a.i./A) in spring to actively growing spring or winter wheat and triticale from the 3-leaf to jointing stage (Zadoks scale 31) according to the application timings shown in the table entitled **Weeds Controlled (C) or Suppressed (S)**. Treat after the majority of weeds have emerged. Best results are obtained when application is made to weeds that are actively growing.

Occasionally, slight yellowing or height reduction may be observed in the treated crop. These transient symptoms disappear within 14 days with no reduction to yield. **DO NOT** apply to crops suffering from drought, water-logged soils, nutrient deficiency or exposed to frost or other agronomic factors affecting plant growth. **DO NOT** use on wheat or triticale varieties that are sensitive to acetolactate synthase (ALS) herbicides.

An independent liquid ammonium nitrogen fertilizer application made 7 days before or after an application of Companera 21.50% WDG may result in transient leaf burn or stunting. DO NOT make a liquid fertilizer application during this period unless the risk of crop response is acceptable.

Tank Mixtures: **Companera 21.50% WDG** may be applied in tank mix combination with labeled rates of other products registered for postemergence application in spring and winter wheat or triticale. See **Tank Mixing Precautions** under **Mixing Directions**. When tank mixing, **DO NOT** exceed specified application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels.

Crop Specific Use Restrictions:

- **Preharvest Interval:** **DO NOT** apply within 60 days of harvest.
- **DO NOT** apply more than 1 oz. of **Companera 21.50% WDG** per acre (0.013 lb. a.i./A) per year.
- **DO NOT** graze the treated crop within 7 days following application.
- **DO NOT** cut the treated crop for hay within 28 days following application.
- **DO NOT** apply a product containing organophosphates for five days before or five days after an application of **Companera 21.50% WDG**.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site according to label use directions or at an approved waste disposal facility.

CONTAINER HANDLING:

Non-Refillable Plastic Containers (Capacity Equal to or Less Than 50 lbs.): Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

CONTAINER HANDLING:

Non-Refillable Plastic Containers (Capacity Greater Than 50 lbs.): Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

CONTAINER HANDLING:

For Bulk and Mini-Bulk Containers

Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

WARRANTY AND DISCLAIMER STATEMENT

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sharda USA, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA, LLC MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Sharda USA, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA, LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SHARDA USA, LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

DECLARACIONES PREVENTIVAS
RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS
PRECAUCIÓN

Dañino si se ingiere. Causa irritación moderada en los ojos. Evite el contacto con los ojos o la ropa. Lávese bien con agua y jabón después de manipular y antes de comer, beber, masticar chicle, usar tabaco o usar el baño. Quítese y lave la ropa contaminada antes de reutilizarla. Use gafas protectoras, camisa de manga larga y pantalones largos, calcetines y zapatos.

EQUIPO DE PROTECCIÓN PERSONAL (PPE, POR SUS SIGLAS EN INGLÉS)

Todos los manipuladores deben usar:

- Camisa de manga larga y pantalones largos
- Zapatos y calcetines

Siga las instrucciones del fabricante para la limpieza y el mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE por separado del resto de la ropa.

DECLARACIÓN DE CONTROL TÉCNICO

Cuando los manipuladores de pesticidas usen sistemas cerrados, cabinas cerradas o aeronaves, de manera que cumplan con los requisitos enumerados en la Norma de Protección al Trabajador (WPS, por sus siglas en inglés) [40 CFR 170.240 (d)(4-6)], para pesticidas agrícolas, los requisitos de equipo de protección personal (PPE) para el manipulador de pesticidas pueden reducirse o modificarse como está especificado en la WPS.

RECOMENDACIONES DE SEGURIDAD DEL USUARIO

Los usuarios deben:

- Lávese bien las manos con agua y jabón después de manipular y antes de comer, beber, masticar chicle, usar tabaco o usar el baño.
- Quitarse inmediatamente la ropa/equipo de protección personal (PPE) si el pesticida entra en contacto con la ropa o el PPE por dentro. Luego deben lavarse muy bien y ponerse ropa limpia.
- Quítese equipo de protección personal (PPE) inmediatamente después de manipular este producto. Lávese bien y póngase ropa limpia lo antes posible.

RIESGOS AMBIENTALES

NO aplique directamente al agua, en áreas donde haya aguas superficiales, o en áreas intermareales que se encuentren bajo el nivel de altura máxima promedio del agua. **NO** contamine el agua al desechar las aguas de lavado del equipo ni las aguas de enjuague.

ADVERTENCIA SOBRE AGUAS SUBTERRÁNEAS: Este producto químico tiene propiedades y características relacionadas con productos químicos detectados en aguas subterráneas. Se sabe que este producto químico se filtra a través del suelo hacia las aguas subterráneas en determinadas condiciones como resultado del uso indicado en la etiqueta. Este producto químico puede filtrarse a las aguas subterráneas si se utiliza en zonas con suelos permeables, especialmente donde la capa freática es poco profunda.

ADVERTENCIA REFERENTE A LAS AGUAS SUPERFICIALES: Este producto puede afectar la calidad del agua superficial debido a la escorrentía de aguas pluviales. Esto es especialmente cierto en suelos con mal drenaje y suelos con niveles freáticos poco profundos.

Este producto está clasificado como de alto potencial para alcanzar las aguas superficiales por escorrentía durante varias semanas después de su aplicación.

Una franja de protección vegetal nivelada y bien mantenida entre las áreas de aplicación y elementos de aguas superficiales, como estanques, arroyos y manantiales, reducirá la posible carga de este producto químico proveniente de la escorrentía y los sedimentos. La escorrentía de este producto puede reducirse al evitar aplicaciones cuando se pronostiquen lluvias o riego en las próximas 48 horas.

DECLARACIÓN CONSULTIVA SOBRE ORGANISMOS NO OBJETIVO: Este producto es tóxico para las plantas y puede afectar negativamente al forraje y al hábitat de organismos no objetivo, incluidos los polinizadores, en zonas adyacentes al lugar tratado. Proteja el forraje y el hábitat de los organismos no objetivo siguiendo las instrucciones de la etiqueta destinadas a minimizar la deriva de la aspersión.

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal.

ALMACENAMIENTO Y DESECHO

NO contamine el agua, los alimentos ni los forrajes durante el almacenamiento o el desecho.

ALMACENAMIENTO DE PESTICIDAS: Almacene únicamente en el envase original.

DESECHO DE PESTICIDAS: Los residuos resultantes del uso de este producto deben eliminarse en el lugar de origen o en una instalación de eliminación de residuos autorizada.

MANIPULACIÓN DE ENVASES:

Envases de plástico no rellenables (Capacidad igual o menos de 50 libras): Envase no rellenable. **NO** reutilice ni rellene este envase. Ofrézcalo para reciclaje si está disponible o perfore y deseche en un relleno sanitario, o mediante incineración, o por otros procedimientos aprobados por las autoridades estatales y locales. Enjuague tres veces el envase (o equivalente) inmediatamente después de vaciarlo. **Enjuague tres veces siguiendo estas indicaciones:** vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezcla. Llene el envase hasta 1/4 de su capacidad con agua y tape de nuevo. Agite por 10 segundos. Vierta las aguas de enjuague en el equipo de aplicación o un tanque de mezcla, o almacene aguas de enjuague para uso posterior o eliminación. Drene por 10 segundos después de que el flujo empiece a gotear. Repita este procedimiento dos veces más. **Enjuague a presión siguiendo estas indicaciones:** vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezcla, y drene por 10 segundos después de que el flujo empiece a gotear. Mantenga el envase boca abajo sobre el equipo de aplicación o un tanque de mezcla, o recolecte aguas de enjuague para uso más tarde o la eliminación. Inserte la boquilla de enjuague a presión en el lado del envase y enjuague a aproximadamente 40 PSI por al menos 30 segundos. Drene por 10 segundos después de que el flujo empiece a gotear.

MANIPULACIÓN DE ENVASES:

Envases de plástico no rellenables (Capacidad más de 50 libras): Envase no rellenable. **NO** reutilice ni rellene este envase. Ofrézcalo para reciclaje si está disponible o perfore y deseche en un relleno sanitario, o mediante incineración, o por otros procedimientos aprobados por las autoridades estatales y locales. Enjuague tres veces el envase (o equivalente) inmediatamente después de vaciarlo. **Enjuague tres veces siguiendo estas indicaciones:** vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezcla. Llene el envase hasta 1/4 de su capacidad con agua. Reemplace y asegure los cierres. Vuelque el envase al lado y hágalo rodar por 30 segundos, asegurando por lo menos una revolución completa. Ponga el envase vertical e inclínelo hacia adelante y hacia atrás varias veces. Dé vuelta al envase hacia el lado opuesto e inclínelo hacia adelante y hacia atrás varias veces. Vierta las aguas de enjuague en el equipo de aplicación o un tanque de mezcla, o almacene aguas de enjuague para uso posterior o eliminación. Repita este procedimiento dos veces más. **Enjuague a presión siguiendo estas indicaciones:** vacíe los contenidos restantes en el equipo de aplicación o un tanque de mezcla, y drene por 10 segundos después de que el flujo empiece a gotear. Mantenga el envase boca abajo sobre el equipo de aplicación o un tanque de mezcla, o recolecte aguas de enjuague para uso más tarde o la eliminación. Inserte la boquilla de enjuague a presión en el lado del envase y enjuague a aproximadamente 40 PSI por al menos 30 segundos. Drene por 10 segundos después de que el flujo empiece a gotear.

MANIPULACIÓN DE ENVASES:

Para contenedores y mini-contenedores a granel

Envase rellenable. Rellene este envase solo con pesticida. **NO** reutilice este envase para ningún otro propósito. La limpieza del envase antes de su eliminación final es la responsabilidad de la persona que se deshace del envase. La limpieza antes de rellenarlo es responsabilidad de la persona que lo rellena. Para limpiar el envase antes de su eliminación final, vacíe el contenido restante de este envase en el equipo de aplicación o en el tanque de mezcla. Llene el envase con un 10% de agua. Agite vigorosamente o haga recircular el agua con la bomba durante 2 minutos. Vierta o bombee las aguas de enjuague en el equipo de aplicación o en el sistema de recolección de aguas de enjuague. Repita este procedimiento dos veces más. Luego ofrézcalo para reciclaje si está disponible o perfore y deseche en un relleno sanitario, o mediante incineración, o por otros procedimientos aprobados por las autoridades estatales y locales.

¡EL ENVASE NO ES SEGURO PARA ALIMENTOS, FORRAJES, NI AGUA POTABLE!

Companera 21.50% WDG

For postemergent control of annual grass and broadleaf weeds
in spring and winter wheat (including durum), and triticale.

ACTIVE INGREDIENT:	WT. BY %
pyroxsulam: N-(5,7-dimethoxy[1,2,4]triazolo [1,5-a]pyrimidin-2-yl)-2-methoxy-4-(trifluoromethyl)-3-pyridinesulfonamide	21.5%
OTHER INGREDIENTS:	78.5%
TOTAL:	100.0%

Contains 0.215 lb. of active ingredient per pound of product.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give anything to an unconscious person.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378 , Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	

See label booklet for complete Precautionary Statements,
Directions For Use and Storage and Disposal.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION/PRECAUCIÓN

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear protective eye-wear, long sleeved shirt and long pants, socks and shoes.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

GROUNDWATER ADVISORY: This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

SURFACE WATER ADVISORY: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having high potential for reaching surface water via runoff for several weeks after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this chemical from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site according to label use directions or at an approved waste disposal facility.

CONTAINER HANDLING:

Non-Refillable Plastic Containers (Capacity Equal to or Less Than 50 lbs.): Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Manufactured For:

Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 82633-141-83529 EPA Est. No. (AG) 72159-GA-001; (MC) 89332-GA-001; (SC) 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Weight: **10 lbs.**